



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:59 AM GMT

PDB ID : 3CX8
Title : Crystal Structure of PDZRhGEF rgRGS Domain in a Complex with Galpha-13 Bound to GTP-gamma-S
Authors : Sprang, S.R.; Chen, Z.
Deposited on : 2008-04-23
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	338	
2	B	203	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	A	378	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4513 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Guanine nucleotide-binding protein alpha-13 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	0	0
			2635	1678	467	480	10			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	40	GLY	-	EXPRESSION TAG	UNP P27601

- Molecule 2 is a protein called Rho guanine nucleotide exchange factor 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	193	Total	C	N	O	S	0	0	0
			1547	977	260	303	7			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	306	GLY	-	EXPRESSION TAG	UNP Q9ES67

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is 5'-GUANOSINE-DIPHOSPHATE-MONOTHIOPHOSPHATE (three-letter code: GSP) (formula: C₁₀H₁₆N₅O₁₃P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	S	0	0
			32	10	5	13	3	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	220	Total O 220 220	0	0
5	B	78	Total O 78 78	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Guanine nucleotide-binding protein alpha-13 subunit



- Molecule 2: Rho guanine nucleotide exchange factor 11



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.11Å 66.12Å 152.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.00 40.30 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.2 (40.00-2.00) 98.2 (40.30-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.00Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.223 , 0.253 0.241 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	26.4	Xtriage
Anisotropy	0.085	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 59.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 39805 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4513	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GSP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/2687	0.58	0/3618
2	B	0.30	0/1574	0.54	0/2127
All	All	0.34	0/4261	0.56	0/5745

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2635	0	2631	56	0
2	B	1547	0	1524	46	0
3	A	1	0	0	0	0
4	A	32	0	12	2	0
5	A	220	0	0	5	0
5	B	78	0	0	5	0
All	All	4513	0	4167	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:307:LEU:HB2	5:B:274:HOH:O	1.83	0.79
2:B:395:LYS:HE2	2:B:395:LYS:HA	1.64	0.78
2:B:330:ASP:HB3	2:B:333:LYS:HD2	1.69	0.75
1:A:302:VAL:HG23	1:A:303:VAL:HG23	1.70	0.71
2:B:307:LEU:HB3	5:B:25:HOH:O	1.91	0.69
2:B:330:ASP:HB3	2:B:333:LYS:CD	2.25	0.65
1:A:259:ASP:O	1:A:261:GLN:N	2.30	0.65
2:B:364:GLU:O	2:B:368:GLN:HG3	1.96	0.65
2:B:347:TYR:CE1	2:B:351:GLN:HG3	2.33	0.64
1:A:191:PRO:HB2	1:A:196:ILE:CD1	2.28	0.64
1:A:191:PRO:HB2	1:A:196:ILE:HD11	1.83	0.61
2:B:307:LEU:N	2:B:307:LEU:HD12	2.14	0.61
1:A:146:LEU:HB3	1:A:147:PRO:HD3	1.82	0.61
1:A:188:ASP:HB2	5:A:948:HOH:O	2.00	0.60
1:A:278:ASN:HB2	2:B:351:GLN:HE22	1.68	0.58
1:A:100:VAL:CG2	1:A:123:MET:SD	2.92	0.58
2:B:466:MET:HG3	2:B:470:GLN:NE2	2.18	0.58
1:A:127:THR:HG22	5:A:767:HOH:O	2.03	0.58
1:A:117:LEU:N	1:A:117:LEU:HD12	2.20	0.57
2:B:469:LYS:HE2	5:B:266:HOH:O	2.05	0.56
2:B:338:PRO:O	2:B:342:VAL:HG23	2.05	0.56
1:A:85:ARG:HB3	1:A:86:PRO:HD3	1.87	0.56
2:B:412:GLU:O	2:B:414:PRO:HD3	2.06	0.55
1:A:176:TYR:CD2	1:A:198:LEU:HD13	2.42	0.55
2:B:335:LYS:HE3	2:B:364:GLU:OE2	2.06	0.55
1:A:341:PRO:HD3	5:A:942:HOH:O	2.06	0.55
2:B:395:LYS:HE2	2:B:395:LYS:CA	2.36	0.55
2:B:466:MET:HE2	2:B:466:MET:O	2.06	0.55
1:A:96:MET:HG3	1:A:122:LEU:HD22	1.89	0.54
2:B:468:GLU:O	2:B:468:GLU:HG3	2.08	0.54
2:B:352:ALA:HB1	5:B:90:HOH:O	2.08	0.53
2:B:482:GLU:HB2	2:B:484:ASP:OD1	2.08	0.53
1:A:176:TYR:CG	1:A:198:LEU:HD13	2.44	0.53
1:A:248:SER:O	1:A:251:GLU:HG2	2.08	0.53
1:A:172:GLU:H	1:A:172:GLU:CD	2.11	0.52
1:A:204:LYS:O	1:A:207:HIS:HE1	1.93	0.52
1:A:117:LEU:H	1:A:117:LEU:HD12	1.74	0.51
2:B:427:LEU:HD13	2:B:427:LEU:C	2.30	0.51
1:A:143:LEU:HD11	5:A:951:HOH:O	2.10	0.51
2:B:346:ARG:HG3	2:B:477:ILE:CD1	2.40	0.51
1:A:270:ASN:O	1:A:274:THR:HG22	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:382:TRP:CH2	2:B:405:ASP:HB2	2.47	0.50
2:B:346:ARG:HG3	2:B:477:ILE:HD11	1.93	0.50
1:A:100:VAL:HG22	1:A:123:MET:SD	2.51	0.49
1:A:278:ASN:CB	2:B:351:GLN:HE22	2.26	0.49
1:A:292:LYS:HD3	1:A:295:LEU:HD12	1.95	0.49
2:B:466:MET:HG3	2:B:470:GLN:HE21	1.76	0.49
1:A:310:LEU:H	1:A:310:LEU:HD12	1.78	0.48
2:B:307:LEU:O	2:B:308:ILE:HB	2.14	0.47
1:A:166:ARG:HH21	1:A:303:VAL:HG21	1.80	0.47
1:A:84:PHE:CD1	1:A:196:ILE:HG13	2.51	0.46
2:B:345:LEU:O	2:B:349:LEU:HD13	2.15	0.46
2:B:457:ASP:O	2:B:459:ASP:N	2.49	0.46
1:A:84:PHE:CG	1:A:196:ILE:HG13	2.51	0.46
2:B:466:MET:CE	2:B:469:LYS:HB3	2.46	0.46
1:A:248:SER:OG	1:A:292:LYS:HD2	2.16	0.45
1:A:114:LYS:O	1:A:117:LEU:HD13	2.17	0.45
1:A:165:ARG:HA	1:A:168:PHE:CZ	2.51	0.45
1:A:310:LEU:N	1:A:310:LEU:HD12	2.31	0.45
2:B:462:ARG:O	2:B:465:GLN:HB3	2.17	0.45
1:A:165:ARG:HA	1:A:168:PHE:CE2	2.52	0.44
1:A:273:GLU:OE1	2:B:439:LYS:NZ	2.46	0.44
1:A:347:THR:HG22	1:A:358:VAL:HG21	2.00	0.44
1:A:117:LEU:H	1:A:117:LEU:CD1	2.30	0.44
1:A:292:LYS:HG2	4:A:755:GSP:C6	2.52	0.44
2:B:366:TYR:CE1	2:B:374:SER:HB2	2.53	0.44
1:A:334:ARG:HH22	1:A:341:PRO:N	2.16	0.44
1:A:94:LYS:HE2	1:A:127:THR:OG1	2.17	0.44
1:A:198:LEU:HA	4:A:755:GSP:O2'	2.18	0.44
2:B:406:LEU:HD23	2:B:406:LEU:C	2.39	0.43
2:B:307:LEU:O	2:B:308:ILE:CB	2.66	0.43
1:A:122:LEU:HD13	1:A:152:LEU:CD1	2.48	0.43
2:B:462:ARG:HD2	2:B:462:ARG:O	2.18	0.43
1:A:237:PHE:HA	1:A:240:VAL:CG2	2.49	0.43
1:A:146:LEU:HG	1:A:150:ARG:HH12	1.83	0.42
2:B:338:PRO:HB2	2:B:467:ALA:HB2	2.02	0.42
1:A:99:LEU:HD13	1:A:170:LEU:HD22	2.02	0.42
2:B:456:LEU:HD23	2:B:462:ARG:NE	2.34	0.42
1:A:127:THR:HG23	2:B:313:GLU:OE1	2.20	0.42
2:B:399:MET:O	2:B:403:GLU:HG3	2.19	0.42
1:A:78:GLN:O	1:A:82:GLU:HG3	2.20	0.42
2:B:308:ILE:N	5:B:274:HOH:O	2.33	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:406:LEU:HD23	2:B:406:LEU:O	2.20	0.41
2:B:466:MET:HE3	2:B:469:LYS:HB3	2.03	0.41
1:A:324:LYS:O	1:A:328:GLU:HG3	2.20	0.41
1:A:184:LEU:HD22	1:A:189:TYR:CZ	2.56	0.41
1:A:237:PHE:HA	1:A:240:VAL:HG21	2.01	0.41
2:B:365:VAL:HG13	2:B:499:HIS:CD2	2.55	0.41
1:A:310:LEU:CD1	1:A:310:LEU:H	2.33	0.41
2:B:461:LEU:HA	2:B:464:ARG:HH11	1.85	0.41
1:A:150:ARG:O	1:A:154:GLU:HG3	2.21	0.41
1:A:103:ARG:NE	5:A:956:HOH:O	2.21	0.41
1:A:290:LEU:HD12	1:A:346:PHE:CE2	2.56	0.41
2:B:403:GLU:O	2:B:407:ARG:HG3	2.21	0.40
1:A:279:ARG:O	2:B:480:LYS:HE2	2.21	0.40
1:A:334:ARG:NH1	1:A:342:LEU:HB2	2.37	0.40
1:A:315:ASP:HA	1:A:316:PRO:HD2	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/338 (93%)	306 (97%)	8 (2%)	1 (0%)	46	41
2	B	189/203 (93%)	180 (95%)	5 (3%)	4 (2%)	9	3
All	All	504/541 (93%)	486 (96%)	13 (3%)	5 (1%)	19	11

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	260	ARG
2	B	469	LYS
2	B	308	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	455	GLY
2	B	458	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/309 (94%)	285 (98%)	6 (2%)	61	63
2	B	170/179 (95%)	163 (96%)	7 (4%)	37	32
All	All	461/488 (94%)	448 (97%)	13 (3%)	51	50

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	LEU
1	A	99	LEU
1	A	100	VAL
1	A	114	LYS
1	A	122	LEU
1	A	274	THR
2	B	307	LEU
2	B	323	GLU
2	B	333	LYS
2	B	401	GLN
2	B	418	LEU
2	B	457	ASP
2	B	469	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	GLN
1	A	160	ASN
1	A	207	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	254	GLN
1	A	283	ASN
1	A	368	HIS
2	B	351	GLN
2	B	383	ASN
2	B	416	ASN
2	B	430	GLN
2	B	434	ASN
2	B	465	GLN
2	B	470	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GSP	A	755	3	25,34,34	1.71	5 (20%)	31,54,54	1.94	7 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GSP	A	755	3	-	0/15/38/38	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	755	GSP	PG-S1G	-3.26	1.84	1.90
4	A	755	GSP	PG-O2G	-2.40	1.46	1.55
4	A	755	GSP	PB-O2B	-2.25	1.45	1.54
4	A	755	GSP	C6-N1	3.59	1.39	1.33
4	A	755	GSP	O4'-C1'	4.44	1.46	1.41

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	755	GSP	N3-C2-N1	-5.28	119.40	127.44
4	A	755	GSP	C6-C5-C4	-3.69	116.49	120.90
4	A	755	GSP	C4-C5-N7	-2.95	106.76	109.48
4	A	755	GSP	O4'-C1'-N9	-2.76	102.32	108.10
4	A	755	GSP	C5-C6-N1	-2.76	119.81	123.59
4	A	755	GSP	N2-C2-N1	2.21	120.86	117.20
4	A	755	GSP	C6-N1-C2	4.01	121.50	115.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	755	GSP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	319/338 (94%)	0.33	10 (3%) 52 53	11, 25, 45, 60	0
2	B	193/203 (95%)	1.00	34 (17%) 2 2	20, 40, 60, 66	0
All	All	512/541 (94%)	0.58	44 (8%) 13 14	11, 29, 58, 66	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	457	ASP	6.0
2	B	307	LEU	5.7
2	B	462	ARG	5.2
2	B	458	GLY	5.2
2	B	454	LEU	4.6
2	B	456	LEU	4.2
2	B	465	GLN	4.0
2	B	472	ALA	4.0
2	B	461	LEU	3.7
1	A	301	GLN	3.5
2	B	476	ASP	3.4
2	B	455	GLY	3.3
1	A	313	GLU	3.3
2	B	463	GLU	3.3
2	B	370	ASN	3.3
1	A	215	ASN	3.1
2	B	451	ASN	3.1
1	A	310	LEU	3.0
2	B	375	ARG	3.0
2	B	332	GLU	3.0
2	B	453	LEU	2.9
2	B	482	GLU	2.8
2	B	388	LYS	2.8
2	B	468	GLU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	467	ALA	2.7
1	A	214	LYS	2.6
2	B	452	ASP	2.6
2	B	372	LYS	2.6
1	A	368	HIS	2.6
1	A	188	ASP	2.5
1	A	336	ASP	2.5
2	B	466	MET	2.5
2	B	464	ARG	2.5
2	B	469	LYS	2.4
2	B	471	LEU	2.4
2	B	374	SER	2.4
1	A	304	SER	2.3
2	B	411	ASN	2.3
1	A	316	PRO	2.3
2	B	459	ASP	2.2
2	B	484	ASP	2.2
2	B	366	TYR	2.2
2	B	394	VAL	2.1
2	B	322	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MG	A	378	1/1	0.96	0.25	2.30	22,22,22,22	0
4	GSP	A	755	32/32	0.99	0.17	0.74	14,17,19,21	0

6.5 Other polymers

There are no such residues in this entry.