



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 10:23 PM GMT

PDB ID : 5CZ2
Title : Crystal structure of a two-domain fragment of MMTV integrase
Authors : Cook, N.; Ballandras-Colas, A.; Engelman, A.; Cherepanov, P.
Deposited on : 2015-07-31
Resolution : 2.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2439 (2.74-2.70)
Clashscore	102246	2771 (2.74-2.70)
Ramachandran outliers	100387	2726 (2.74-2.70)
Sidechain outliers	100360	2727 (2.74-2.70)
RSRZ outliers	91569	2443 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	210	<div> <div>2%</div> <div>61% 11% 28%</div> </div>
1	B	210	<div> <div>3%</div> <div>59% 9% 32%</div> </div>
1	C	210	<div> <div>2%</div> <div>62% 10% 28%</div> </div>
1	D	210	<div> <div>6%</div> <div>59% 7% 34%</div> </div>
1	E	210	<div> <div>5%</div> <div>67% 6% 27%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	210	<p>8% 49% 6% 45%</p>
1	G	210	<p>20% 80%</p>
1	H	210	<p>2% 19% 81%</p>
1	I	210	<p>1% 19% 81%</p>
1	J	210	<p>3% 19% 80%</p>
1	K	210	<p>3% 17% 83%</p>
1	L	210	<p>19% 80%</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8591 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pol polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	151	Total	C	N	O	S	0	0	0
			1214	775	225	211	3			
1	B	142	Total	C	N	O	S	0	0	0
			1150	737	209	201	3			
1	C	152	Total	C	N	O	S	0	0	0
			1207	770	224	210	3			
1	D	139	Total	C	N	O	S	0	0	0
			1098	701	200	194	3			
1	E	153	Total	C	N	O	S	0	0	0
			1215	774	227	211	3			
1	F	115	Total	C	N	O	S	0	0	0
			895	574	163	155	3			
1	G	41	Total	C	N	O	S	0	0	0
			323	200	65	56	2			
1	H	40	Total	C	N	O	S	0	0	0
			291	181	57	51	2			
1	I	39	Total	C	N	O	S	0	0	0
			282	176	56	48	2			
1	J	41	Total	C	N	O	S	0	0	0
			315	195	64	54	2			
1	K	36	Total	C	N	O	S	0	0	0
			270	167	56	45	2			
1	L	41	Total	C	N	O	S	0	0	0
			319	198	65	54	2			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total	Mg	0	0
			1	1		
2	E	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total 1	Mg 1	0	0
2	C	1	Total 1	Mg 1	0	0
2	A	1	Total 1	Mg 1	0	0
2	F	1	Total 1	Mg 1	0	0


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	1	Total 1	Zn 1	0	0
3	J	1	Total 1	Zn 1	0	0
3	K	1	Total 1	Zn 1	0	0
3	H	1	Total 1	Zn 1	0	0
3	I	1	Total 1	Zn 1	0	0
3	L	1	Total 1	Zn 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Pol polypeptide




• Molecule 1: Pol polypeptide




• Molecule 1: Pol polypeptide




• Molecule 1: Pol polypeptide









- Molecule 1: Pol polypeptide




- Molecule 1: Pol polypeptide




- Molecule 1: Pol polypeptide



- Molecule 1: Pol polypeptide







4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.37Å 83.15Å 141.14Å 90.00° 90.19° 90.00°	Depositor
Resolution (Å)	70.57 – 2.72 70.57 – 2.72	Depositor EDS
% Data completeness (in resolution range)	99.3 (70.57-2.72) 98.3 (70.57-2.72)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.14 (at 2.73Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.241 , 0.280 0.240 , 0.278	Depositor DCC
R_{free} test set	1654 reflections (4.90%)	DCC
Wilson B-factor (Å ²)	54.4	Xtriage
Anisotropy	0.082	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.8	EDS
Estimated twinning fraction	0.865 for H, K, L 0.135 for -H, -K, L 0.116 for h,-k,-l	Xtriage
Reported twinning fraction	0.865 for H, K, L 0.135 for -H, -K, L	Depositor
L-test for twinning ²	$\langle L \rangle = 0.37$, $\langle L^2 \rangle = 0.20$	Xtriage
Outliers	0 of 33771 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	8591	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.86 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.2301e-03.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/1247	0.39	0/1692
1	B	0.23	0/1181	0.39	0/1602
1	C	0.23	0/1239	0.39	0/1682
1	D	0.23	0/1128	0.38	0/1535
1	E	0.23	0/1247	0.39	0/1693
1	F	0.22	0/916	0.38	0/1245
1	G	0.22	0/330	0.37	0/446
1	H	0.23	0/298	0.36	0/407
1	I	0.23	0/289	0.36	0/395
1	J	0.22	0/322	0.37	0/437
1	K	0.23	0/276	0.36	0/374
1	L	0.22	0/326	0.37	0/441
All	All	0.23	0/8799	0.38	0/11949

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1214	0	1189	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1150	0	1117	10	0
1	C	1207	0	1183	12	0
1	D	1098	0	1032	9	0
1	E	1215	0	1187	6	0
1	F	895	0	847	6	0
1	G	323	0	313	0	0
1	H	291	0	255	0	0
1	I	282	0	249	0	0
1	J	315	0	298	2	0
1	K	270	0	235	0	0
1	L	319	0	309	2	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
3	I	1	0	0	0	0
3	J	1	0	0	0	0
3	K	1	0	0	0	0
3	L	1	0	0	0	0
All	All	8591	0	8214	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:115:ILE:HD12	1:E:141:ILE:HD11	1.79	0.65
1:C:115:ILE:HD12	1:C:141:ILE:HD11	1.83	0.61
1:A:115:ILE:HD12	1:A:141:ILE:HD11	1.82	0.61
1:E:54:ARG:NH2	1:E:85:TYR:O	2.40	0.55
1:A:143:HIS:O	1:L:20:ARG:NH2	2.39	0.54
1:D:92:THR:OG1	1:D:105:HIS:ND1	2.37	0.53
1:C:111:ALA:O	1:D:193:HIS:NE2	2.42	0.52
1:C:92:THR:OG1	1:C:105:HIS:ND1	2.37	0.52
1:A:92:THR:OG1	1:A:105:HIS:ND1	2.42	0.49
1:A:133:GLN:HB3	1:L:26:THR:HG23	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:94:ARG:HE	1:D:105:HIS:HB2	1.79	0.48
1:C:75:LEU:HD12	1:C:180:PRO:HG2	1.96	0.48
1:C:143:HIS:O	1:J:20:ARG:NH2	2.42	0.47
1:A:75:LEU:HD12	1:A:180:PRO:HG2	1.97	0.47
1:B:115:ILE:HD12	1:B:141:ILE:HD11	1.97	0.47
1:E:196:MET:HB2	1:E:196:MET:HE2	1.71	0.46
1:E:86:SER:O	1:E:206:ARG:NH2	2.48	0.46
1:E:196:MET:HG2	1:E:202:THR:HG22	1.98	0.46
1:B:94:ARG:HE	1:B:105:HIS:HB2	1.81	0.46
1:B:196:MET:HB2	1:B:196:MET:HE3	1.79	0.45
1:D:196:MET:HG2	1:D:202:THR:HG22	1.99	0.45
1:F:196:MET:HB2	1:F:196:MET:HE2	1.80	0.44
1:E:75:LEU:HD12	1:E:180:PRO:HG2	1.99	0.44
1:C:207:HIS:HD2	1:C:208:TRP:CE2	2.36	0.43
1:C:196:MET:HG2	1:C:202:THR:HG22	2.00	0.43
1:A:54:ARG:NH2	1:A:85:TYR:O	2.51	0.43
1:D:196:MET:HE2	1:D:196:MET:HB2	1.76	0.43
1:B:88:PHE:CD1	1:B:113:MET:HG2	2.53	0.43
1:D:88:PHE:CD1	1:D:113:MET:HG2	2.53	0.43
1:B:99:THR:HG21	1:B:131:SER:HB2	2.01	0.43
1:C:196:MET:HB2	1:C:196:MET:HE2	1.62	0.43
1:F:92:THR:OG1	1:F:105:HIS:ND1	2.46	0.43
1:B:177:TYR:CD2	1:B:183:LEU:HG	2.53	0.43
1:A:72:PHE:CD1	1:A:180:PRO:HB3	2.53	0.42
1:A:166:ALA:O	1:A:170:LYS:HG3	2.19	0.42
1:A:196:MET:HG2	1:A:202:THR:HG22	2.02	0.42
1:D:166:ALA:O	1:D:170:LYS:HG3	2.20	0.42
1:D:207:HIS:HD2	1:D:208:TRP:CE2	2.37	0.42
1:C:88:PHE:CD1	1:C:113:MET:HG2	2.55	0.41
1:A:115:ILE:HA	1:A:115:ILE:HD13	1.90	0.41
1:F:207:HIS:HD2	1:F:208:TRP:CE2	2.38	0.41
1:A:88:PHE:CD1	1:A:113:MET:HG2	2.55	0.41
1:B:166:ALA:O	1:B:170:LYS:HG3	2.20	0.41
1:F:88:PHE:CD1	1:F:113:MET:HG2	2.56	0.41
1:B:92:THR:OG1	1:B:105:HIS:ND1	2.46	0.41
1:A:52:ASN:OD1	1:A:155:ALA:N	2.54	0.41
1:F:115:ILE:HD13	1:F:115:ILE:HA	1.93	0.41
1:B:72:PHE:CD1	1:B:180:PRO:HB3	2.55	0.41
1:F:166:ALA:O	1:F:170:LYS:HG3	2.21	0.40
1:C:133:GLN:HB3	1:J:26:THR:HG23	2.03	0.40
1:B:154:GLN:O	1:B:158:GLU:HG2	2.21	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:193:HIS:NE2	1:D:111:ALA:O	2.51	0.40
1:C:94:ARG:HE	1:C:105:HIS:HB2	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	147/210 (70%)	142 (97%)	5 (3%)	0	100	100
1	B	136/210 (65%)	134 (98%)	2 (2%)	0	100	100
1	C	148/210 (70%)	144 (97%)	4 (3%)	0	100	100
1	D	133/210 (63%)	131 (98%)	2 (2%)	0	100	100
1	E	149/210 (71%)	144 (97%)	5 (3%)	0	100	100
1	F	105/210 (50%)	104 (99%)	1 (1%)	0	100	100
1	G	39/210 (19%)	38 (97%)	1 (3%)	0	100	100
1	H	38/210 (18%)	37 (97%)	1 (3%)	0	100	100
1	I	37/210 (18%)	36 (97%)	1 (3%)	0	100	100
1	J	39/210 (19%)	38 (97%)	1 (3%)	0	100	100
1	K	32/210 (15%)	31 (97%)	1 (3%)	0	100	100
1	L	39/210 (19%)	38 (97%)	1 (3%)	0	100	100
All	All	1042/2520 (41%)	1017 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/176 (72%)	122 (97%)	4 (3%)	46	76
1	B	120/176 (68%)	118 (98%)	2 (2%)	68	89
1	C	125/176 (71%)	120 (96%)	5 (4%)	38	68
1	D	111/176 (63%)	109 (98%)	2 (2%)	66	88
1	E	125/176 (71%)	122 (98%)	3 (2%)	57	84
1	F	90/176 (51%)	87 (97%)	3 (3%)	45	75
1	G	33/176 (19%)	33 (100%)	0	100	100
1	H	27/176 (15%)	27 (100%)	0	100	100
1	I	26/176 (15%)	26 (100%)	0	100	100
1	J	31/176 (18%)	31 (100%)	0	100	100
1	K	26/176 (15%)	26 (100%)	0	100	100
1	L	32/176 (18%)	32 (100%)	0	100	100
All	All	872/2112 (41%)	853 (98%)	19 (2%)	60	86

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	139	TRP
1	A	172	GLN
1	A	173	LYS
1	A	183	LEU
1	B	172	GLN
1	B	183	LEU
1	C	108	GLN
1	C	139	TRP
1	C	172	GLN
1	C	173	LYS
1	C	183	LEU
1	D	172	GLN
1	D	183	LEU
1	E	139	TRP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	172	GLN
1	E	183	LEU
1	F	172	GLN
1	F	173	LYS
1	F	183	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	104	GLN
1	I	39	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	151/210 (71%)	0.39	5 (3%)	50	50	30, 53, 100, 129	0
1	B	142/210 (67%)	0.50	6 (4%)	40	40	32, 57, 87, 102	0
1	C	152/210 (72%)	0.42	5 (3%)	50	50	33, 57, 91, 106	0
1	D	139/210 (66%)	0.69	12 (8%)	13	11	41, 73, 112, 155	0
1	E	153/210 (72%)	0.56	11 (7%)	18	17	35, 61, 105, 137	0
1	F	115/210 (54%)	0.99	17 (14%)	3	2	52, 74, 116, 150	0
1	G	41/210 (19%)	0.31	1 (2%)	62	62	44, 53, 81, 120	0
1	H	40/210 (19%)	0.91	4 (10%)	9	7	52, 83, 129, 166	0
1	I	39/210 (18%)	0.77	2 (5%)	32	31	50, 70, 115, 133	0
1	J	41/210 (19%)	0.89	6 (14%)	3	2	65, 79, 107, 133	0
1	K	36/210 (17%)	0.85	6 (16%)	2	2	72, 96, 113, 131	0
1	L	41/210 (19%)	0.16	1 (2%)	62	62	40, 57, 81, 91	0
All	All	1090/2520 (43%)	0.59	76 (6%)	19	18	30, 64, 108, 166	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	34	VAL	5.2
1	A	174	ALA	5.2
1	K	34	VAL	5.1
1	F	179	THR	5.0
1	F	191	LEU	4.9
1	I	33	ILE	4.7
1	F	182	HIS	4.6
1	D	128	VAL	4.3
1	H	33	ILE	4.3
1	H	16	ALA	4.2
1	F	106	LEU	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	141	ILE	3.7
1	F	128	VAL	3.7
1	F	180	PRO	3.5
1	K	38	PRO	3.4
1	J	38	PRO	3.3
1	F	183	LEU	3.3
1	H	34	VAL	3.3
1	J	36	LEU	3.2
1	K	33	ILE	3.2
1	F	104	GLN	3.1
1	F	177	TYR	3.1
1	H	17	ALA	3.1
1	D	95	THR	3.0
1	E	174	ALA	3.0
1	J	34	VAL	3.0
1	E	176	LYS	2.9
1	A	171	LEU	2.9
1	E	177	TYR	2.8
1	D	200	GLY	2.8
1	D	191	LEU	2.7
1	K	20	ARG	2.7
1	F	119	ILE	2.7
1	J	27	ARG	2.7
1	E	89	THR	2.7
1	C	52	ASN	2.7
1	F	80	VAL	2.7
1	A	154	GLN	2.6
1	E	189	PHE	2.6
1	D	80	VAL	2.6
1	J	29	GLN	2.5
1	F	61	LEU	2.5
1	B	141	ILE	2.5
1	K	26	THR	2.5
1	F	127	TYR	2.5
1	C	129	SER	2.5
1	D	84	THR	2.5
1	A	177	TYR	2.5
1	C	74	LYS	2.5
1	A	124	ALA	2.5
1	D	136	LEU	2.5
1	D	72	PHE	2.4
1	L	36	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	38	PRO	2.4
1	K	30	ALA	2.3
1	F	63	GLN	2.3
1	B	191	LEU	2.2
1	F	84	THR	2.2
1	E	156	ILE	2.2
1	F	91	ALA	2.2
1	E	132	ILE	2.2
1	J	37	CYS	2.2
1	E	173	LYS	2.2
1	B	136	LEU	2.2
1	D	119	ILE	2.1
1	C	209	GLY	2.1
1	D	194	VAL	2.1
1	E	171	LEU	2.1
1	F	165	LYS	2.1
1	B	178	TYR	2.1
1	D	178	TYR	2.1
1	E	178	TYR	2.0
1	C	50	GLY	2.0
1	B	69	VAL	2.0
1	B	144	VAL	2.0
1	E	180	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ZN	L	301	1/1	0.94	0.17	0.26	55,55,55,55	0
3	ZN	G	301	1/1	0.89	0.14	-1.37	55,55,55,55	0
3	ZN	I	301	1/1	0.89	0.10	-1.46	83,83,83,83	0
3	ZN	J	301	1/1	0.95	0.08	-2.23	57,57,57,57	0
3	ZN	H	301	1/1	0.94	0.11	-2.49	86,86,86,86	0
2	MG	B	301	1/1	0.78	0.11	-	52,52,52,52	0
2	MG	E	301	1/1	0.91	0.21	-	46,46,46,46	0
2	MG	C	301	1/1	0.82	0.25	-	65,65,65,65	0
2	MG	A	301	1/1	0.88	0.16	-	52,52,52,52	0
3	ZN	K	301	1/1	0.98	0.07	-	64,64,64,64	0
2	MG	F	301	1/1	0.86	0.06	-	53,53,53,53	0
2	MG	D	301	1/1	0.78	0.17	-	81,81,81,81	0

6.5 Other polymers [i](#)

There are no such residues in this entry.