



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:27 AM GMT

PDB ID : 2D2M
Title : Structure of an extracellular giant hemoglobin of the gutless beard worm *Oligobrachia mashikoi*
Authors : Numoto, N.; Nakagawa, T.; Kita, A.; Sasayama, Y.; Fukumori, Y.; Miki, K.
Deposited on : 2005-09-12
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2499 (2.90-2.82)
Ramachandran outliers	100387	2439 (2.90-2.82)
Sidechain outliers	100360	2442 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	140	
2	B	142	
3	C	147	
4	D	145	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4457 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Giant hemoglobin, A1(b) globin chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	140	Total	C	N	O	S	0	0	0
			1065	656	196	206	7			

- Molecule 2 is a protein called Giant hemoglobin, A2(a5) globin chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	142	Total	C	N	O	S	0	0	0
			1075	662	198	209	6			

- Molecule 3 is a protein called Giant hemoglobin, B2(c) globin chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	146	Total	C	N	O	S	0	0	0
			1088	668	201	210	9			

- Molecule 4 is a protein called Giant hemoglobin, B1(d) globin chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	145	Total	C	N	O	S	0	0	0
			1036	642	178	209	7			

- Molecule 5 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
5	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
5	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
5	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 6 is OXYGEN MOLECULE (three-letter code: OXY) (formula: O_2).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O 2 2	0	0
6	B	1	Total O 2 2	0	0
6	C	1	Total O 2 2	0	0
6	D	1	Total O 2 2	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	6	Total O 6 6	0	0
7	B	1	Total O 1 1	0	0
7	C	3	Total O 3 3	0	0
7	D	3	Total O 3 3	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: Giant hemoglobin, A1(b) globin chain




- Molecule 2: Giant hemoglobin, A2(a5) globin chain



- Molecule 3: Giant hemoglobin, B2(c) globin chain



- Molecule 4: Giant hemoglobin, B1(d) globin chain



V102	A106	V107	T108	H109	L110	S111	Q112	A113	I114	S115	S116	V117	V118	V121	L122	P123	S124	A125	H126	I127	D128	A129	W130	M134	I137	G142	L145
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	111.50Å 111.50Å 276.83Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.85	Depositor
% Data completeness (in resolution range)	97.3 (50.00-2.85)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.236 , 0.277	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4457	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, OXY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/1083	0.55	0/1464
2	B	0.32	0/1093	0.53	0/1481
3	C	0.32	0/1110	0.51	0/1507
4	D	0.30	0/1055	0.52	0/1436
All	All	0.31	0/4341	0.53	0/5888

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1065	0	1015	74	0
2	B	1075	0	1038	71	0
3	C	1088	0	1013	64	0
4	D	1036	0	987	52	0
5	A	43	0	30	4	0
5	B	43	0	30	4	0
5	C	43	0	30	0	0
5	D	43	0	30	2	0
6	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	2	0	0	0	0
6	C	2	0	0	0	0
6	D	2	0	0	0	0
7	A	6	0	0	0	0
7	B	1	0	0	0	0
7	C	3	0	0	0	0
7	D	3	0	0	0	0
All	All	4457	0	4173	245	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 245 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:25:SER:HA	3:C:28:ARG:HD3	1.42	0.98
1:A:14:TRP:HE1	1:A:71:ASN:ND2	1.65	0.95
1:A:46:PHE:HB2	1:A:51:VAL:HG11	1.50	0.93
2:B:9:LEU:HD21	2:B:13:GLN:HE21	1.34	0.93
3:C:19:ALA:HB1	3:C:123:VAL:HG12	1.54	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/140 (99%)	122 (88%)	10 (7%)	6 (4%)	3	11
2	B	140/142 (99%)	109 (78%)	27 (19%)	4 (3%)	6	20
3	C	144/147 (98%)	120 (83%)	18 (12%)	6 (4%)	3	11
4	D	143/145 (99%)	118 (82%)	21 (15%)	4 (3%)	6	21

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	565/574 (98%)	469 (83%)	76 (14%)	20 (4%)	4	16

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	49	VAL
3	C	104	VAL
2	B	18	TYR
2	B	54	ILE
3	C	54	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	110/110 (100%)	108 (98%)	2 (2%)	66	89
2	B	113/113 (100%)	105 (93%)	8 (7%)	18	44
3	C	110/111 (99%)	106 (96%)	4 (4%)	42	75
4	D	108/108 (100%)	104 (96%)	4 (4%)	41	74
All	All	441/442 (100%)	423 (96%)	18 (4%)	37	71

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	95	ASP
2	B	127	ASP
4	D	28	GLU
2	B	89	ARG
2	B	91	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	84	ASN
3	C	47	GLN
4	D	17	GLN
2	B	134	ASN
3	C	11	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	HEM	A	200	1,6	30,50,50	2.21	7 (23%)	24,82,82	2.42	8 (33%)
6	OXY	A	201	5	1,1,1	0.12	0	0,0,0	0.00	-
6	OXY	B	1201	5	1,1,1	0.13	0	0,0,0	0.00	-
5	HEM	B	200	2,6	30,50,50	2.22	6 (20%)	24,82,82	2.41	8 (33%)
5	HEM	C	200	3,6	30,50,50	2.15	6 (20%)	24,82,82	2.38	8 (33%)
6	OXY	C	2201	5	1,1,1	0.13	0	0,0,0	0.00	-
5	HEM	D	200	4,6	30,50,50	2.14	5 (16%)	24,82,82	2.44	8 (33%)
6	OXY	D	3201	5	1,1,1	0.13	0	0,0,0	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEM	A	200	1,6	-	0/10/54/54	0/0/8/8
6	OXY	A	201	5	-	0/0/0/0	0/0/0/0
6	OXY	B	1201	5	-	0/0/0/0	0/0/0/0
5	HEM	B	200	2,6	-	0/10/54/54	0/0/8/8
5	HEM	C	200	3,6	-	0/10/54/54	0/0/8/8
6	OXY	C	2201	5	-	0/0/0/0	0/0/0/0
5	HEM	D	200	4,6	-	0/10/54/54	0/0/8/8
6	OXY	D	3201	5	-	0/0/0/0	0/0/0/0

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	200	HEM	C3B-C4B	-6.82	1.45	1.51
5	B	200	HEM	C3B-C4B	-6.77	1.45	1.51
5	C	200	HEM	C3B-C4B	-6.61	1.46	1.51
5	D	200	HEM	C3B-C4B	-6.36	1.46	1.51
5	B	200	HEM	C2D-C3D	-5.35	1.38	1.54

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	200	HEM	C3B-CAB-CBB	2.04	127.58	124.46
5	C	200	HEM	C3B-C4B-CHC	2.11	126.13	123.16
5	C	200	HEM	C3C-CAC-CBC	2.16	127.76	124.46
5	B	200	HEM	C3B-CAB-CBB	2.16	127.78	124.46
5	B	200	HEM	C3B-C4B-CHC	2.20	126.26	123.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	200	HEM	4	0
5	B	200	HEM	4	0
5	D	200	HEM	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates

EDS was not executed - this section will therefore be empty.

6.4 Ligands

EDS was not executed - this section will therefore be empty.

6.5 Other polymers

EDS was not executed - this section will therefore be empty.