



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2016 – 04:44 AM EDT

PDB ID : 5D29
Title : X-ray structure of human glutamate carboxypeptidase II (GCP II) in complex with a hydroxamate inhibitor JHU241
Authors : Barinka, C.; Novakova, Z.; Pavlicek, J.
Deposited on : 2015-08-05
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027674
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	695	<div> <div>4%</div> <div>88%</div> <div>10% ..</div> </div>

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 6297 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate carboxypeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	686	Total	C	N	O	S	0	40	0
			5743	3681	969	1072	21			

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is BETA-D-MANNOSE (three-letter code: BMA) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		


- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Cl	0	0
			1	1		

- Molecule 7 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is 4-[(2 {S})-2-carboxy-5-(oxidanylamino)-5-oxidanylidene-pentyl]benzoic acid (three-letter code: 5Q1) (formula: C₁₃H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			20	13	1	6		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	364	Total	O	0	0
			364	364		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glutamate carboxypeptidase 2



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	100.19 Å 130.46 Å 157.20 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.82 – 1.80 44.82 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (44.82-1.80) 99.7 (44.82-1.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 1.79 Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.174 , 0.205 0.175 , 0.205	Depositor DCC
R_{free} test set	4755 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	23.4	Xtriage
Anisotropy	0.577	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 49.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6297	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BMA, NAG, CL, CA, ACT, 5Q1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.02	4/5921 (0.1%)	0.83	3/8018 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	697[A]	HIS	C-N	14.98	1.68	1.34
1	A	697[B]	HIS	C-N	14.98	1.68	1.34
1	A	522	GLU	CD-OE2	-6.29	1.18	1.25
1	A	557	GLU	CB-CG	-5.20	1.42	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	697[A]	HIS	C-N-CA	-6.01	106.67	121.70
1	A	697[B]	HIS	C-N-CA	-6.01	106.67	121.70
1	A	280	ARG	NE-CZ-NH1	-5.20	117.70	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5743	0	5562	61	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	140	0	126	4	0
3	A	22	0	19	0	0
4	A	2	0	0	0	0
5	A	1	0	0	0	0
6	A	1	0	0	0	0
7	A	4	0	3	0	0
8	A	20	0	0	0	0
9	A	364	0	0	7	0
All	All	6297	0	5710	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:697[A]:HIS:C	1:A:698:ASN:N	1.68	1.43
1:A:693[B]:ALA:HB2	9:A:1056:HOH:O	1.31	1.22
1:A:465[A]:ASP:OD1	1:A:536[A]:ARG:NH1	1.90	1.04
1:A:684:ARG:NH2	1:A:694[B]:PRO:O	1.96	0.96
1:A:177:VAL:HG12	1:A:188:LEU:HD11	1.58	0.85
1:A:494:TYR:O	1:A:498[A]:THR:HG23	1.77	0.84
2:A:808:NAG:H83	9:A:1187:HOH:O	1.78	0.82
1:A:684:ARG:NH1	1:A:694[B]:PRO:O	2.13	0.80
1:A:697[A]:HIS:C	1:A:698:ASN:CA	2.52	0.78
1:A:684:ARG:CZ	1:A:694[B]:PRO:O	2.34	0.76
1:A:517:SER:OG	1:A:522:GLU:OE2	2.04	0.74
1:A:681:LEU:HD11	1:A:693[B]:ALA:HB3	1.74	0.69
1:A:87[A]:GLU:CD	1:A:87[A]:GLU:H	1.96	0.68
1:A:697[B]:HIS:ND1	9:A:902:HOH:O	2.28	0.65
1:A:517:SER:OG	1:A:518:GLY:N	2.30	0.64
1:A:659:ILE:O	1:A:663[A]:MET:HG3	1.98	0.63
1:A:517:SER:HB2	1:A:694[A]:PRO:HG3	1.81	0.61
1:A:58:MET:HE1	1:A:586:GLU:HG2	1.84	0.60
1:A:58:MET:CE	1:A:586:GLU:HG2	2.32	0.60
1:A:400:ARG:O	1:A:404:THR:HG23	2.02	0.59
1:A:131:ILE:HG22	1:A:137:GLU:HG2	1.85	0.58
1:A:697[A]:HIS:CA	1:A:698:ASN:N	2.63	0.57
1:A:681:LEU:CD1	1:A:693[B]:ALA:HB3	2.37	0.55
1:A:695[B]:SER:C	1:A:697[B]:HIS:H	2.11	0.52
1:A:612:TYR:CZ	1:A:616:MET:HG3	2.45	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:ILE:HD12	1:A:88:GLN:HG2	1.91	0.52
1:A:174:LEU:HD22	1:A:342:VAL:HG21	1.91	0.51
1:A:307:GLU:HA	1:A:325:VAL:HG22	1.93	0.51
1:A:263:GLY:HA2	1:A:522:GLU:OE1	2.11	0.51
1:A:498[A]:THR:HG22	1:A:510:PRO:HD3	1.92	0.51
1:A:517:SER:CB	1:A:694[A]:PRO:HG3	2.41	0.51
1:A:681:LEU:HD11	1:A:693[B]:ALA:CB	2.42	0.50
1:A:687:TYR:CZ	1:A:694[A]:PRO:HG2	2.46	0.50
1:A:693[B]:ALA:CB	9:A:1056:HOH:O	2.13	0.50
1:A:246:TRP:CD1	2:A:806:NAG:H83	2.48	0.48
1:A:681:LEU:CD1	1:A:693[B]:ALA:CB	2.92	0.47
1:A:70:ILE:HG21	1:A:573:HIS:HB3	1.97	0.46
1:A:90:PHE:CE2	1:A:94:LYS:HE2	2.51	0.46
1:A:177:VAL:HG12	1:A:188:LEU:CD1	2.38	0.46
1:A:258:ILE:CD1	1:A:290:PRO:HG3	2.45	0.46
1:A:695[B]:SER:O	1:A:697[B]:HIS:N	2.48	0.46
1:A:61:PHE:HB2	1:A:405[A]:LEU:HD11	1.97	0.45
1:A:347:HIS:HE1	9:A:1221:HOH:O	1.98	0.45
1:A:691:ILE:O	1:A:704:SER:HA	2.16	0.45
1:A:695[B]:SER:C	1:A:697[B]:HIS:N	2.70	0.44
1:A:169:MET:HA	1:A:344:MET:O	2.18	0.44
1:A:349:THR:HG21	2:A:803:NAG:H81	1.98	0.44
1:A:437:GLU:OE1	9:A:901:HOH:O	2.21	0.44
1:A:498[A]:THR:HG22	1:A:510:PRO:CD	2.48	0.44
1:A:349:THR:HG21	2:A:803:NAG:C8	2.48	0.43
1:A:721:PRO:HD2	9:A:940:HOH:O	2.19	0.43
1:A:258:ILE:HD12	1:A:290:PRO:HG3	2.00	0.43
1:A:489:GLU:H	1:A:489:GLU:CD	2.22	0.42
1:A:662[B]:ARG:HD3	1:A:662[B]:ARG:HA	1.93	0.42
1:A:131:ILE:CG2	1:A:137:GLU:HG2	2.49	0.41
1:A:215:LYS:HD2	1:A:260[B]:ASN:OD1	2.20	0.41
1:A:188:LEU:HD21	1:A:329:VAL:HG11	2.02	0.41
1:A:444:GLU:HA	1:A:444:GLU:OE1	2.21	0.41
1:A:663[A]:MET:HB3	1:A:663[A]:MET:HE3	1.74	0.41
1:A:188:LEU:CD2	1:A:329:VAL:HG11	2.50	0.41
1:A:660:VAL:O	1:A:664[A]:MET:HB2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	718/695 (103%)	697 (97%)	18 (2%)	3 (0%)	39 23

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	696[A]	SER
1	A	696[B]	SER
1	A	382	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	620/592 (105%)	604 (97%)	16 (3%)	54 37

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	134	ASP
1	A	155[A]	SER
1	A	155[B]	SER
1	A	187	LYS
1	A	239[A]	VAL
1	A	239[B]	VAL
1	A	388	PRO
1	A	465[A]	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	465[B]	ASP
1	A	536[A]	ARG
1	A	536[B]	ARG
1	A	537	TYR
1	A	600	TYR
1	A	664[A]	MET
1	A	664[B]	MET
1	A	673	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 4 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	801	1,2	14,14,15	0.61	0	15,19,21	1.37	2 (13%)
2	NAG	A	802	2	14,14,15	0.59	0	15,19,21	1.24	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	803	1	14,14,15	0.60	0	15,19,21	1.64	3 (20%)
2	NAG	A	804	1	14,14,15	0.76	0	15,19,21	0.86	0
2	NAG	A	805	1	14,14,15	0.73	1 (7%)	15,19,21	2.54	4 (26%)
2	NAG	A	806	1	14,14,15	0.71	0	15,19,21	1.76	3 (20%)
2	NAG	A	807	1,2	14,14,15	0.93	0	15,19,21	1.60	2 (13%)
2	NAG	A	808	2	14,14,15	0.82	0	15,19,21	1.42	3 (20%)
2	NAG	A	809	1,2	14,14,15	0.61	0	15,19,21	1.47	3 (20%)
2	NAG	A	810	3,2	14,14,15	0.83	1 (7%)	15,19,21	2.52	5 (33%)
3	BMA	A	811	3,2	11,11,12	0.63	0	15,15,17	1.83	3 (20%)
3	BMA	A	812	3	11,11,12	0.72	0	15,15,17	2.84	3 (20%)
7	ACT	A	817	-	0,3,3	0.00	-	0,3,3	0.00	-
8	5Q1	A	818	4	14,20,20	1.16	2 (14%)	19,26,26	1.87	8 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	801	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	802	2	-	0/6/23/26	0/1/1/1
2	NAG	A	803	1	-	0/6/23/26	0/1/1/1
2	NAG	A	804	1	-	0/6/23/26	0/1/1/1
2	NAG	A	805	1	-	0/6/23/26	0/1/1/1
2	NAG	A	806	1	-	0/6/23/26	0/1/1/1
2	NAG	A	807	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	808	2	-	0/6/23/26	0/1/1/1
2	NAG	A	809	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	810	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	811	3,2	-	0/2/19/22	0/1/1/1
3	BMA	A	812	3	-	0/2/19/22	0/1/1/1
7	ACT	A	817	-	-	0/0/0/0	0/0/0/0
8	5Q1	A	818	4	-	0/11/19/19	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	810	NAG	O5-C1	-2.03	1.40	1.43
2	A	805	NAG	C1-C2	2.15	1.55	1.52
8	A	818	5Q1	CAK-CAQ	2.23	1.55	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	818	5Q1	OAC-CAQ	2.35	1.28	1.23

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	812	BMA	C1-O5-C5	-8.88	99.08	112.14
2	A	810	NAG	C2-N2-C7	-7.22	113.71	123.11
2	A	807	NAG	O5-C5-C4	-4.38	102.88	110.13
3	A	812	BMA	C1-C2-C3	-3.94	104.77	109.55
2	A	809	NAG	C2-N2-C7	-3.71	118.28	123.11
2	A	806	NAG	C2-N2-C7	-3.28	118.84	123.11
3	A	811	BMA	O3-C3-C4	-3.25	103.03	110.36
2	A	801	NAG	O4-C4-C3	-3.05	103.48	110.36
8	A	818	5Q1	CAM-CAT-CAP	-3.01	105.03	111.02
2	A	808	NAG	O7-C7-C8	-2.91	116.71	122.07
2	A	806	NAG	O3-C3-C4	-2.88	103.85	110.36
2	A	807	NAG	O4-C4-C3	-2.84	103.95	110.36
8	A	818	5Q1	CAJ-CAS-CAO	-2.83	116.71	120.43
8	A	818	5Q1	CAJ-CAH-CAR	-2.74	117.17	121.02
2	A	803	NAG	O3-C3-C4	-2.65	104.39	110.36
8	A	818	5Q1	CAG-CAI-CAS	-2.52	117.60	121.15
2	A	801	NAG	C3-C4-C5	-2.44	105.87	110.23
2	A	810	NAG	C4-C3-C2	-2.42	107.59	111.34
2	A	810	NAG	O7-C7-C8	-2.32	117.80	122.07
2	A	808	NAG	C4-C3-C2	-2.27	107.82	111.34
8	A	818	5Q1	OAC-CAQ-NAN	-2.26	121.03	123.60
2	A	809	NAG	C6-C5-C4	-2.04	107.89	112.99
2	A	805	NAG	O3-C3-C2	2.03	113.71	109.37
2	A	803	NAG	C3-C4-C5	2.26	114.25	110.23
3	A	811	BMA	O2-C2-C3	2.33	114.88	110.19
2	A	805	NAG	O5-C5-C6	2.41	112.50	107.34
8	A	818	5Q1	CAJ-CAS-CAI	2.42	122.28	117.56
2	A	810	NAG	O4-C4-C3	2.55	116.10	110.36
8	A	818	5Q1	CAH-CAR-CAG	2.74	122.68	118.15
2	A	805	NAG	C2-N2-C7	2.83	126.78	123.11
2	A	802	NAG	C2-N2-C7	2.95	126.94	123.11
2	A	806	NAG	C1-O5-C5	2.98	116.52	112.14
8	A	818	5Q1	OAC-CAQ-CAK	3.01	127.19	121.97
2	A	809	NAG	C1-O5-C5	3.21	116.86	112.14
2	A	808	NAG	C8-C7-N2	3.24	122.30	116.10
3	A	812	BMA	O5-C5-C6	3.81	115.49	107.34
2	A	810	NAG	C1-O5-C5	3.88	117.84	112.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	803	NAG	C1-O5-C5	3.89	117.86	112.14
3	A	811	BMA	O3-C3-C2	5.03	119.22	110.01
2	A	805	NAG	C1-O5-C5	7.98	123.87	112.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	803	NAG	2	0
2	A	806	NAG	1	0
2	A	808	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	697[B]:HIS	C	698:ASN	N	2.90
1	A	697[A]:HIS	C	698:ASN	N	1.68

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	686/695 (98%)	-0.10	28 (4%) 41 35	16, 29, 53, 71	5 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	700[A]	TYR	4.8
1	A	699[A]	LYS	4.5
1	A	155[A]	SER	4.0
1	A	505	GLU	3.3
1	A	719	VAL	3.2
1	A	701	ALA	3.2
1	A	123	THR	3.0
1	A	506	PHE	3.0
1	A	337	PHE	2.9
1	A	335	GLY	2.9
1	A	124	HIS	2.8
1	A	702	GLY	2.8
1	A	698	ASN	2.8
1	A	695[A]	SER	2.8
1	A	134	ASP	2.8
1	A	487	GLY	2.8
1	A	507	SER	2.7
1	A	333	PHE	2.6
1	A	153	ASN	2.6
1	A	133	GLU	2.6
1	A	122	LYS	2.6
1	A	336	ASN	2.4
1	A	56	HIS	2.3
1	A	312[A]	SER	2.3
1	A	341	LYS	2.2
1	A	334	THR	2.2
1	A	338	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	194	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	A	803	14/15	0.74	0.28	1.57	52,59,66,67	0
3	BMA	A	812	11/12	0.86	0.14	1.47	58,61,65,67	0
2	NAG	A	807	14/15	0.93	0.09	1.46	30,34,38,39	0
2	NAG	A	809	14/15	0.96	0.08	0.72	22,30,42,47	0
8	5Q1	A	818	20/20	0.96	0.14	0.14	19,24,29,33	0
7	ACT	A	817	4/4	0.95	0.12	-0.18	29,30,30,31	0
5	CA	A	815	1/1	1.00	0.09	-0.32	17,17,17,17	0
2	NAG	A	804	14/15	0.85	0.11	-0.73	43,46,50,51	0
6	CL	A	816	1/1	0.98	0.10	-1.34	26,26,26,26	0
4	ZN	A	813	1/1	1.00	0.09	-1.50	21,21,21,21	0
4	ZN	A	814	1/1	1.00	0.10	-1.62	21,21,21,21	0
2	NAG	A	801	14/15	0.94	0.12	-	34,44,52,54	0
2	NAG	A	805	14/15	0.71	0.30	-	70,78,79,80	0
2	NAG	A	808	14/15	0.90	0.17	-	41,47,51,53	0
3	BMA	A	811	11/12	0.89	0.08	-	45,50,55,55	0
2	NAG	A	806	14/15	0.84	0.12	-	38,49,54,56	0
2	NAG	A	802	14/15	0.82	0.28	-	50,59,63,64	0
2	NAG	A	810	14/15	0.90	0.13	-	42,48,56,57	0

6.5 Other polymers [i](#)

There are no such residues in this entry.