



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 06:54 PM GMT

PDB ID : 1D39  
Title : COVALENT MODIFICATION OF GUANINE BASES IN DOUBLE STRANDED DNA: THE 1.2 ANGSTROMS Z-DNA STRUCTURE OF D(CGCGCG) IN THE PRESENCE OF CUCL2  
Authors : Kagawa, T.F.; Geierstanger, B.H.; Wang, A.H.-J.; Ho, P.S.  
Deposited on : 1991-05-07  
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1607 (1.26-1.14)
RSRZ outliers	91569	1500 (1.26-1.14)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	6	 100%
1	B	6	 67% 33%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 313 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(\*CP\*(CU)GP\*CP\*(CU)GP\*CP\*(CU)G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	6	Total	C	N	O	P	0	0	0
			120	57	24	34	5			
1	B	6	Total	C	N	O	P	0	0	0
			120	57	24	34	5			

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	3	Total	Cu	0	0
			3	3		
2	A	3	Total	Cu	0	0
			3	3		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	38	Total	O	0	0
			38	38		
4	B	28	Total	O	0	0
			28	28		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(\*CP\*(CU)GP\*CP\*(CU)GP\*CP\*(CU)G)-3')

Chain A:  100%

C1 C2 C3 C4 C5 C6

- Molecule 1: DNA (5'-D(\*CP\*(CU)GP\*CP\*(CU)GP\*CP\*(CU)G)-3')

Chain B:  67% 33%

C7 C8 C9 C10 C11 C12

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	18.01Å 31.03Å 44.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 1.20 31.03 – 1.26	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-1.20) 83.8 (31.03-1.26)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.70 (at 1.26Å)	Xtriage
Refinement program	NUCLSQ	Depositor
R, $R_{free}$	0.198 , (Not available) 0.249 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	7.8	Xtriage
Anisotropy	0.259	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.25 , 62.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	4 of 7081 reflections (0.056%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	313	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.53 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 7.8767e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	4.11	33/134 (24.6%)	5.07	50/205 (24.4%)
1	B	4.06	26/134 (19.4%)	6.14	81/205 (39.5%)
All	All	4.08	59/268 (22.0%)	5.63	131/410 (32.0%)

All (59) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	8	DG	C4'-C3'	11.81	1.65	1.53
1	B	12	DG	C3'-C2'	-11.42	1.38	1.52
1	A	2	DG	N1-C2	-10.22	1.29	1.37
1	A	6	DG	C2'-C1'	10.06	1.62	1.52
1	A	5	DC	N1-C6	9.99	1.43	1.37
1	B	7	DC	N3-C4	9.95	1.41	1.33
1	A	1	DC	N3-C4	9.83	1.40	1.33
1	B	12	DG	C2-N2	-9.78	1.24	1.34
1	A	2	DG	N9-C8	-9.68	1.31	1.37
1	B	10	DG	C2-N2	-9.54	1.25	1.34
1	B	7	DC	N1-C6	9.47	1.42	1.37
1	A	4	DG	C2-N2	-8.74	1.25	1.34
1	A	2	DG	C2-N2	-8.73	1.25	1.34
1	A	2	DG	C4'-C3'	8.68	1.62	1.53
1	B	10	DG	C2-N3	8.61	1.39	1.32
1	B	8	DG	C2'-C1'	8.48	1.60	1.52
1	B	12	DG	C8-N7	8.40	1.35	1.30
1	A	3	DC	C4-C5	-8.15	1.36	1.43
1	A	6	DG	C2-N2	-8.01	1.26	1.34
1	B	12	DG	C2'-C1'	7.56	1.59	1.52
1	A	3	DC	P-O5'	7.43	1.67	1.59
1	A	2	DG	C5-C4	-7.28	1.33	1.38
1	A	4	DG	N1-C2	-7.24	1.31	1.37
1	B	12	DG	N9-C8	-7.03	1.32	1.37

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	2	DG	C2-N3	6.89	1.38	1.32
1	A	1	DC	C2'-C1'	6.62	1.58	1.52
1	B	11	DC	N3-C4	6.46	1.38	1.33
1	A	5	DC	O4'-C1'	6.43	1.50	1.42
1	A	2	DG	C1'-N9	-6.32	1.38	1.47
1	A	2	DG	N9-C4	6.25	1.43	1.38
1	B	11	DC	N1-C6	6.15	1.40	1.37
1	A	6	DG	C6-N1	-6.06	1.35	1.39
1	A	2	DG	C8-N7	6.03	1.34	1.30
1	A	6	DG	C6-O6	5.98	1.29	1.24
1	A	6	DG	N1-C2	-5.96	1.32	1.37
1	B	7	DC	N1-C2	-5.95	1.34	1.40
1	B	10	DG	C6-N1	-5.94	1.35	1.39
1	B	11	DC	P-OP2	5.72	1.58	1.49
1	B	7	DC	O5'-C5'	-5.66	1.28	1.42
1	B	8	DG	C6-O6	5.64	1.29	1.24
1	B	11	DC	P-O5'	5.60	1.65	1.59
1	B	8	DG	C8-N7	5.59	1.34	1.30
1	A	4	DG	C4'-O4'	-5.57	1.39	1.45
1	A	6	DG	C4'-C3'	5.57	1.58	1.53
1	A	3	DC	C3'-C2'	5.56	1.58	1.52
1	B	11	DC	C3'-C2'	5.51	1.58	1.52
1	A	2	DG	P-OP1	-5.50	1.39	1.49
1	B	10	DG	C4'-C3'	5.49	1.58	1.53
1	A	4	DG	C1'-N9	-5.41	1.39	1.47
1	B	12	DG	C1'-N9	-5.38	1.39	1.47
1	A	5	DC	N3-C4	5.31	1.37	1.33
1	A	6	DG	N9-C8	-5.28	1.34	1.37
1	B	10	DG	C1'-N9	-5.27	1.39	1.47
1	A	2	DG	C3'-C2'	-5.24	1.46	1.52
1	A	4	DG	P-OP2	-5.21	1.40	1.49
1	B	10	DG	N9-C8	-5.13	1.34	1.37
1	B	11	DC	C2-N3	5.06	1.39	1.35
1	A	5	DC	P-O5'	5.03	1.64	1.59
1	A	2	DG	O4'-C1'	5.02	1.48	1.42

All (131) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	10	DG	N3-C2-N2	-18.80	106.74	119.90
1	B	8	DG	C6-N1-C2	-17.65	114.51	125.10
1	A	4	DG	N3-C2-N2	-17.28	107.80	119.90

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	9	DC	N3-C4-C5	-17.06	115.08	121.90
1	B	7	DC	O4'-C4'-C3'	-16.59	96.04	106.00
1	B	9	DC	C2-N3-C4	16.48	128.14	119.90
1	A	1	DC	C2-N3-C4	16.17	127.98	119.90
1	A	6	DG	N3-C4-C5	-15.27	120.97	128.60
1	B	10	DG	C6-N1-C2	-14.77	116.23	125.10
1	A	1	DC	N3-C4-C5	-14.42	116.13	121.90
1	B	8	DG	N1-C2-N3	14.06	132.34	123.90
1	B	8	DG	C5-C6-N1	14.04	118.52	111.50
1	B	10	DG	N1-C2-N3	13.86	132.22	123.90
1	A	6	DG	N3-C4-N9	13.83	134.30	126.00
1	A	3	DC	N3-C4-C5	-13.40	116.54	121.90
1	B	12	DG	C6-N1-C2	-13.40	117.06	125.10
1	A	1	DC	C6-N1-C2	13.39	125.66	120.30
1	B	8	DG	N3-C2-N2	-13.26	110.62	119.90
1	B	10	DG	O4'-C4'-C3'	-13.02	98.19	106.00
1	A	3	DC	C2-N3-C4	12.91	126.35	119.90
1	A	1	DC	N1-C2-O2	12.70	126.52	118.90
1	B	12	DG	C5-C6-N1	12.30	117.65	111.50
1	A	3	DC	O5'-P-OP1	-12.08	94.83	105.70
1	A	6	DG	N7-C8-N9	-12.02	107.09	113.10
1	B	10	DG	O3'-P-O5'	11.89	126.60	104.00
1	B	11	DC	O4'-C4'-C3'	-11.81	98.91	106.00
1	B	12	DG	N1-C6-O6	-11.45	113.03	119.90
1	B	10	DG	C2-N3-C4	-11.26	106.27	111.90
1	A	6	DG	N3-C2-N2	-11.14	112.10	119.90
1	A	6	DG	C8-N9-C4	11.09	110.83	106.40
1	A	6	DG	C2-N3-C4	11.08	117.44	111.90
1	A	1	DC	N1-C2-N3	-10.98	111.51	119.20
1	B	10	DG	P-O3'-C3'	10.83	132.70	119.70
1	B	10	DG	C5-C6-N1	10.76	116.88	111.50
1	B	12	DG	O4'-C1'-N9	-10.56	100.61	108.00
1	B	7	DC	C2-N3-C4	10.30	125.05	119.90
1	A	6	DG	C6-N1-C2	-9.96	119.12	125.10
1	A	5	DC	N3-C4-C5	-9.87	117.95	121.90
1	B	12	DG	N3-C4-N9	9.75	131.85	126.00
1	B	9	DC	N3-C4-N4	9.60	124.72	118.00
1	B	12	DG	C8-N9-C4	9.57	110.23	106.40
1	B	10	DG	O4'-C1'-N9	-9.46	101.38	108.00
1	B	7	DC	C5-C6-N1	9.29	125.65	121.00
1	A	6	DG	C5-N7-C8	9.18	108.89	104.30
1	A	6	DG	C5-C6-N1	9.15	116.07	111.50

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	7	DC	N3-C4-C5	-9.13	118.25	121.90
1	A	2	DG	O4'-C4'-C3'	-9.02	100.59	106.00
1	B	12	DG	N7-C8-N9	-8.93	108.63	113.10
1	A	2	DG	C8-N9-C4	8.87	109.95	106.40
1	B	8	DG	C5-C6-O6	-8.85	123.29	128.60
1	B	12	DG	N3-C4-C5	-8.79	124.20	128.60
1	A	3	DC	C4-C5-C6	8.78	121.79	117.40
1	B	12	DG	N3-C2-N2	-8.65	113.85	119.90
1	B	9	DC	O4'-C4'-C3'	-8.59	100.85	106.00
1	A	5	DC	C2-N3-C4	8.55	124.18	119.90
1	B	9	DC	O4'-C1'-C2'	-8.46	99.13	105.90
1	B	10	DG	C5-C6-O6	-8.45	123.53	128.60
1	B	7	DC	C4'-C3'-C2'	8.40	110.66	103.10
1	A	4	DG	N1-C2-N2	8.28	123.65	116.20
1	A	3	DC	N1-C2-N3	-8.23	113.44	119.20
1	A	5	DC	P-O3'-C3'	8.16	129.49	119.70
1	B	7	DC	N1-C2-O2	8.07	123.74	118.90
1	A	3	DC	O4'-C1'-N1	-8.04	102.37	108.00
1	B	9	DC	O4'-C1'-N1	8.02	113.61	108.00
1	B	11	DC	O5'-P-OP1	-7.98	98.52	105.70
1	B	9	DC	C1'-O4'-C4'	7.86	117.96	110.10
1	A	5	DC	C4'-C3'-C2'	7.80	110.12	103.10
1	A	4	DG	N1-C2-N3	7.68	128.51	123.90
1	A	2	DG	P-O3'-C3'	7.50	128.70	119.70
1	B	12	DG	C1'-O4'-C4'	-7.50	102.60	110.10
1	B	11	DC	C6-N1-C2	-7.33	117.37	120.30
1	B	7	DC	C3'-C2'-C1'	-7.28	93.77	102.50
1	B	7	DC	C5'-C4'-C3'	7.22	127.09	114.10
1	B	12	DG	P-O5'-C5'	-7.19	109.39	120.90
1	B	10	DG	C5-N7-C8	-7.11	100.74	104.30
1	A	1	DC	O4'-C1'-N1	7.04	112.93	108.00
1	A	2	DG	N9-C4-C5	-7.01	102.60	105.40
1	B	11	DC	C4-C5-C6	6.95	120.87	117.40
1	B	10	DG	O5'-P-OP1	-6.94	99.46	105.70
1	B	11	DC	C1'-O4'-C4'	6.83	116.93	110.10
1	B	11	DC	N3-C4-C5	-6.79	119.19	121.90
1	A	6	DG	O5'-P-OP1	-6.73	99.64	105.70
1	B	8	DG	O5'-P-OP1	-6.73	99.64	105.70
1	B	8	DG	P-O5'-C5'	-6.73	110.13	120.90
1	B	8	DG	N3-C4-N9	6.70	130.02	126.00
1	B	11	DC	N3-C2-O2	-6.66	117.24	121.90
1	B	12	DG	O5'-P-OP1	-6.60	99.76	105.70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	10	DG	O4'-C1'-C2'	6.59	111.17	105.90
1	B	8	DG	C1'-O4'-C4'	6.57	116.67	110.10
1	B	8	DG	OP1-P-OP2	6.45	129.28	119.60
1	B	10	DG	C4'-C3'-C2'	6.44	108.90	103.10
1	A	3	DC	N1-C2-O2	6.33	122.70	118.90
1	B	11	DC	C5-C4-N4	6.33	124.63	120.20
1	B	11	DC	C4'-C3'-C2'	6.31	108.78	103.10
1	B	9	DC	N1-C2-N3	-6.29	114.80	119.20
1	B	10	DG	O5'-P-OP2	6.24	118.19	110.70
1	A	4	DG	C6-N1-C2	-6.12	121.43	125.10
1	A	2	DG	N7-C8-N9	-6.12	110.04	113.10
1	B	9	DC	N3-C2-O2	6.12	126.18	121.90
1	B	10	DG	C4-C5-N7	6.09	113.24	110.80
1	B	10	DG	C4-N9-C1'	6.06	134.38	126.50
1	B	7	DC	C1'-O4'-C4'	6.04	116.14	110.10
1	B	12	DG	C2-N3-C4	6.04	114.92	111.90
1	A	6	DG	N1-C2-N2	6.00	121.61	116.20
1	A	2	DG	N3-C2-N2	-5.98	115.72	119.90
1	A	6	DG	C4-C5-N7	-5.89	108.44	110.80
1	A	4	DG	C5-C6-N1	5.88	114.44	111.50
1	A	3	DC	C6-N1-C2	5.86	122.64	120.30
1	B	10	DG	C8-N9-C1'	-5.82	119.43	127.00
1	B	9	DC	C4-C5-C6	5.65	120.22	117.40
1	B	12	DG	N1-C2-N3	5.62	127.27	123.90
1	B	11	DC	P-O3'-C3'	5.58	126.40	119.70
1	A	3	DC	C5-C4-N4	5.52	124.06	120.20
1	A	2	DG	O5'-C5'-C4'	-5.51	97.21	111.00
1	A	3	DC	C3'-C2'-C1'	-5.46	95.95	102.50
1	B	8	DG	N3-C4-C5	-5.42	125.89	128.60
1	B	10	DG	N7-C8-N9	5.39	115.80	113.10
1	A	1	DC	C5-C4-N4	5.36	123.95	120.20
1	A	1	DC	O4'-C4'-C3'	5.35	109.21	106.00
1	B	7	DC	C5-C4-N4	5.33	123.93	120.20
1	B	7	DC	O4'-C1'-C2'	-5.32	101.64	105.90
1	B	10	DG	N1-C2-N2	5.30	120.97	116.20
1	A	2	DG	C5-C6-O6	-5.30	125.42	128.60
1	A	5	DC	C5-C4-N4	5.25	123.87	120.20
1	B	7	DC	N1-C2-N3	-5.19	115.57	119.20
1	B	12	DG	C5-N7-C8	5.19	106.89	104.30
1	A	1	DC	P-O3'-C3'	5.15	125.88	119.70
1	B	10	DG	P-O5'-C5'	-5.14	112.68	120.90
1	B	12	DG	C8-N9-C1'	-5.11	120.36	127.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2	DG	C4'-C3'-C2'	5.05	107.64	103.10
1	B	9	DC	C4'-C3'-C2'	5.01	107.61	103.10

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	120	0	61	0	0
1	B	120	0	64	0	2
2	A	3	0	0	0	0
2	B	3	0	0	0	2
3	B	1	0	0	0	0
4	A	38	0	0	0	0
4	B	28	0	0	0	7
All	All	313	0	125	0	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

All (7) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:15:CU:CU	4:B:79:HOH:O[4_466]	0.29	1.91
2:B:14:CU:CU	4:B:27:HOH:O[4_566]	0.59	1.61
4:B:27:HOH:O	4:B:76:HOH:O[4_466]	1.00	1.20
1:B:10:DG:N7	4:B:27:HOH:O[4_566]	1.83	0.37
4:B:27:HOH:O	4:B:77:HOH:O[4_466]	2.04	0.16
4:B:27:HOH:O	4:B:78:HOH:O[4_466]	2.07	0.13
1:B:12:DG:N7	4:B:79:HOH:O[4_466]	2.13	0.07

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	6/6 (100%)	0.19	0 100 100	8, 9, 11, 12	0
1	B	6/6 (100%)	0.31	0 100 100	8, 9, 11, 12	0
All	All	12/12 (100%)	0.25	0 100 100	8, 9, 12, 12	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	CU	B	14	1/1	0.93	0.18	-	39,39,39,39	1
2	CU	A	8	1/1	0.41	0.91	-	53,53,53,53	1
3	NA	B	16	1/1	0.80	0.34	-	35,35,35,35	0
2	CU	B	13	1/1	0.72	0.74	-	59,59,59,59	1
2	CU	A	9	1/1	0.99	0.18	-	19,19,19,19	1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CU	B	15	1/1	0.91	0.09	-	38,38,38,38	1
2	CU	A	7	1/1	0.11	1.48	-	75,75,75,75	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.