



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:27 AM GMT

PDB ID : 2D4Q  
Title : Crystal structure of the Sec-PH domain of the human neurofibromatosis type 1 protein  
Authors : D'angelo, I.; Welte, S.; Bonneau, F.; Scheffzek, K.  
Deposited on : 2005-10-22  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	<div> <div>5%</div> <div>79%</div> <div>18%</div> <div>.</div> </div>
1	B	257	<div> <div>6%</div> <div>69%</div> <div>26%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	OXN	A	701	-	-	-	X
2	OXN	B	702	-	-	-	X
3	POP	B	602	-	-	-	X

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4320 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neurofibromin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	89	0	0
			2082	1349	346	382	5			
1	B	255	Total	C	N	O	S	73	0	0
			2059	1336	340	378	5			

- Molecule 2 is OXTOXYNOL-10 (three-letter code: OXN) (formula: C<sub>34</sub>H<sub>62</sub>O<sub>11</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			24	20	4		
2	B	1	Total	C	O	0	0
			24	20	4		

- Molecule 3 is PYROPHOSPHATE 2- (three-letter code: POP) (formula: H<sub>2</sub>O<sub>7</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	P	0	0
			9	7	2		
3	B	1	Total	O	P	0	0
			9	7	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	58	Total	O	0	0
			58	58		
4	B	55	Total	O	0	0
			55	55		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Neurofibromin



#### • Molecule 1: Neurofibromin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	113.50 Å 113.50 Å 125.60 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.30 14.99 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.5 (15.00-2.30) 100.0 (14.99-2.35)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.48 (at 2.34 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.240 , 0.280 0.242 , 0.288	Depositor DCC
$R_{free}$ test set	1720 reflections (4.97%)	DCC
Wilson B-factor (Å <sup>2</sup> )	41.4	Xtriage
Anisotropy	0.259	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 49.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 34716 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4320	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OXN, POP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.55	0/2133	0.73	1/2892 (0.0%)
1	B	0.59	0/2109	0.77	0/2861
All	All	0.57	0/4242	0.75	1/5753 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	1720	HIS	N-CA-C	7.57	131.44	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2082	0	2082	39	1
1	B	2059	0	2050	52	1
2	A	24	0	30	1	0
2	B	24	0	30	1	0
3	B	18	0	0	2	0
4	A	58	0	0	4	0
4	B	55	0	0	4	0
All	All	4320	0	4192	91	1



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1575:ALA:HB1	1:B:1704:LYS:HD2	1.09	1.06
1:A:1605:HIS:O	1:A:1609:THR:HG22	1.54	1.06
1:B:1793:HIS:HD2	1:B:1795:GLU:H	1.18	0.90
1:B:1623:ASP:OD2	1:B:1625:THR:HG23	1.72	0.88
1:B:1793:HIS:CD2	1:B:1795:GLU:H	1.98	0.81
1:B:1575:ALA:CB	1:B:1704:LYS:HD2	2.04	0.80
1:A:1611:LYS:HB3	1:A:1612:PRO:HD3	1.65	0.78
1:B:1625:THR:HG21	1:B:1709:THR:HG23	1.66	0.77
1:A:1793:HIS:HD2	1:A:1795:GLU:H	1.33	0.76
1:B:1751:VAL:O	4:B:88:HOH:O	2.05	0.72
1:A:1662:ASN:ND2	1:A:1665:VAL:H	1.90	0.69
1:B:1604:TYR:CZ	1:B:1608:LEU:HD11	2.30	0.66
1:A:1662:ASN:C	1:A:1662:ASN:HD22	1.98	0.66
1:B:1566:LEU:O	1:B:1571:ILE:HD11	1.96	0.65
1:B:1687:PHE:CG	1:B:1756:VAL:HG11	2.32	0.65
1:B:1737:GLY:O	1:B:1806:ILE:HG12	1.97	0.64
1:B:1779:THR:HG22	1:B:1790:THR:HB	1.80	0.63
1:A:1598:ASN:HD21	1:A:1600:ASP:HB2	1.65	0.62
1:A:1666:ARG:O	1:A:1669:THR:HG22	2.01	0.61
1:B:1634:LYS:HA	3:B:602:POP:O2	2.02	0.60
1:A:1669:THR:HG23	4:A:81:HOH:O	2.02	0.59
1:B:1695:ALA:HA	1:B:1698:ILE:O	2.02	0.59
1:B:1720:HIS:O	1:B:1793:HIS:HE1	1.86	0.59
1:A:1695:ALA:O	1:A:1701:GLU:HA	2.03	0.59
1:A:1790:THR:HG23	4:A:29:HOH:O	2.02	0.59
1:B:1806:ILE:HD13	4:B:42:HOH:O	2.04	0.57
1:B:1623:ASP:OD2	1:B:1625:THR:CG2	2.50	0.56
1:B:1584:ILE:HD13	1:B:1698:ILE:HD13	1.88	0.56
1:A:1687:PHE:CG	1:A:1756:VAL:HG11	2.41	0.55
1:B:1569:LEU:O	1:B:1570:SER:HB2	2.07	0.55
1:B:1793:HIS:HD2	1:B:1795:GLU:N	1.98	0.55
1:A:1793:HIS:CD2	1:A:1795:GLU:H	2.21	0.55
1:B:1598:ASN:HD22	1:B:1601:LEU:H	1.55	0.55
1:A:1737:GLY:O	1:A:1806:ILE:HG12	2.06	0.55
1:A:1605:HIS:O	1:A:1609:THR:CG2	2.43	0.54
1:A:1669:THR:CG2	4:A:81:HOH:O	2.54	0.54
1:B:1656:VAL:HB	1:B:1685:LEU:HD23	1.90	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1592:PHE:CD1	1:A:1602:LEU:HD22	2.42	0.54
1:B:1682:SER:C	1:B:1684:ARG:H	2.12	0.54
1:A:1598:ASN:ND2	1:A:1601:LEU:H	2.06	0.53
1:B:1750:LYS:HA	1:B:1754:GLN:O	2.08	0.53
1:A:1624:LEU:HD13	1:A:1665:VAL:HG11	1.92	0.51
1:A:1695:ALA:HB1	1:A:1701:GLU:HA	1.92	0.51
1:A:1695:ALA:HB1	1:A:1701:GLU:CA	2.41	0.51
1:B:1593:LYS:HB3	1:B:1596:GLN:HG2	1.93	0.51
1:B:1778:PHE:HE1	1:B:1780:LEU:HD13	1.76	0.51
1:B:1606:VAL:HG21	1:B:1641:TRP:HH2	1.76	0.51
1:B:1748:ARG:HD3	1:B:1755:SER:HB3	1.92	0.51
1:A:1598:ASN:HD22	1:A:1601:LEU:H	1.59	0.51
1:A:1807:ARG:O	1:A:1811:GLU:HG3	2.11	0.51
1:B:1571:ILE:HD12	4:B:16:HOH:O	2.11	0.50
1:A:1574:GLN:HG2	1:A:1575:ALA:N	2.26	0.50
1:B:1598:ASN:HD21	1:B:1600:ASP:HB2	1.75	0.50
1:B:1776:ASN:HA	1:B:1793:HIS:O	2.10	0.50
1:A:1622:VAL:HG13	1:A:1658:ILE:HA	1.93	0.50
1:B:1598:ASN:HD22	1:B:1601:LEU:HG	1.76	0.49
1:A:1702:GLN:HE21	1:A:1702:GLN:HA	1.75	0.49
1:B:1797:GLU:O	1:B:1801:GLN:HG2	2.13	0.48
1:A:1594:THR:HG21	1:A:1632:ARG:HG2	1.95	0.48
1:B:1598:ASN:ND2	1:B:1601:LEU:H	2.12	0.48
1:B:1778:PHE:CE1	1:B:1780:LEU:HD13	2.49	0.48
1:B:1634:LYS:HG2	3:B:602:POP:O2	2.15	0.47
1:B:1604:TYR:CE1	1:B:1608:LEU:HD21	2.50	0.47
1:B:1604:TYR:CE2	1:B:1608:LEU:HD11	2.50	0.47
1:A:1691:PRO:HB3	1:A:1703:GLN:NE2	2.29	0.47
1:B:1674:ARG:HG3	4:B:11:HOH:O	2.16	0.46
1:A:1645:PHE:HB2	1:A:1650:TYR:CZ	2.51	0.46
1:B:1603:ILE:HD11	1:B:1637:PHE:CE1	2.50	0.46
1:A:1702:GLN:O	1:A:1702:GLN:HG3	2.15	0.46
1:B:1677:THR:HG23	1:B:1678:GLY:N	2.31	0.45
1:B:1611:LYS:HB3	1:B:1612:PRO:HD3	1.98	0.45
1:B:1720:HIS:O	1:B:1793:HIS:CE1	2.70	0.44
1:A:1799:ILE:O	1:A:1803:ILE:HG13	2.18	0.44
1:A:1660:ASN:ND2	1:A:1759:ASN:HD22	2.15	0.44
1:A:1662:ASN:HD21	1:A:1665:VAL:H	1.60	0.44
1:B:1638:LEU:HG	1:B:1668:TYR:OH	2.18	0.44
1:B:1624:LEU:HD13	1:B:1665:VAL:HG11	2.00	0.43
1:A:1645:PHE:HB2	1:A:1650:TYR:CE1	2.53	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1620:ILE:HD13	2:A:701:OXN:H191	2.01	0.43
1:A:1690:CYS:SG	1:A:1693:LYS:HG2	2.59	0.42
1:B:1602:LEU:HD13	1:B:1641:TRP:HZ2	1.84	0.42
1:A:1662:ASN:ND2	1:A:1662:ASN:C	2.71	0.42
1:B:1593:LYS:H	1:B:1597:ILE:HD12	1.84	0.42
1:A:1703:GLN:HE22	1:A:1710:LEU:HD21	1.85	0.42
1:B:1743:VAL:HG13	1:B:1762:TYR:HE1	1.85	0.41
1:B:1653:VAL:HG21	2:B:702:OXN:H221	2.01	0.41
1:A:1699:GLU:O	1:A:1700:HIS:C	2.58	0.41
1:B:1675:LEU:C	1:B:1675:LEU:HD23	2.40	0.41
1:B:1765:SER:HB2	1:B:1810:TRP:CZ2	2.55	0.41
1:A:1779:THR:HG22	4:A:48:HOH:O	2.21	0.40
1:B:1566:LEU:O	1:B:1571:ILE:CD1	2.69	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1667:GLU:OE1	1:B:1667:GLU:OE2[3_455]	2.11	0.09

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/257 (99%)	239 (94%)	15 (6%)	1 (0%)	39	48
1	B	251/257 (98%)	240 (96%)	10 (4%)	1 (0%)	39	48
All	All	506/514 (98%)	479 (95%)	25 (5%)	2 (0%)	39	48

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1596	GLN
1	B	1683	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/227 (100%)	213 (94%)	14 (6%)	23	30
1	B	224/227 (99%)	204 (91%)	20 (9%)	12	14
All	All	451/454 (99%)	417 (92%)	34 (8%)	17	21

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1594	THR
1	A	1609	THR
1	A	1614	TYR
1	A	1639	SER
1	A	1660	ASN
1	A	1662	ASN
1	A	1669	THR
1	A	1699	GLU
1	A	1702	GLN
1	A	1716	LEU
1	A	1725	LEU
1	A	1790	THR
1	A	1801	GLN
1	A	1806	ILE
1	B	1571	ILE
1	B	1602	LEU
1	B	1614	TYR
1	B	1625	THR
1	B	1638	LEU
1	B	1660	ASN
1	B	1694	LEU
1	B	1701	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	1710	LEU
1	B	1715	ASP
1	B	1723	LEU
1	B	1730	THR
1	B	1749	THR
1	B	1776	ASN
1	B	1780	LEU
1	B	1789	LEU
1	B	1790	THR
1	B	1797	GLU
1	B	1802	SER
1	B	1806	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1596	GLN
1	A	1598	ASN
1	A	1660	ASN
1	A	1662	ASN
1	A	1672	HIS
1	A	1702	GLN
1	A	1785	GLN
1	A	1793	HIS
1	A	1801	GLN
1	B	1598	ASN
1	B	1672	HIS
1	B	1754	GLN
1	B	1759	ASN
1	B	1777	GLN
1	B	1793	HIS
1	B	1801	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	OXN	A	701	-	24,24,45	1.77	4 (16%)	29,30,53	2.12	5 (17%)
3	POP	B	601	-	8,8,8	1.78	2 (25%)	13,13,13	1.51	2 (15%)
3	POP	B	602	-	8,8,8	1.82	3 (37%)	13,13,13	1.55	1 (7%)
2	OXN	B	702	-	24,24,45	1.97	6 (25%)	29,30,53	2.08	5 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OXN	A	701	-	-	0/22/22/43	0/1/1/1
3	POP	B	601	-	-	0/6/6/6	0/0/0/0
3	POP	B	602	-	-	0/6/6/6	0/0/0/0
2	OXN	B	702	-	-	0/22/22/43	0/1/1/1

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	602	POP	P2-O6	2.13	1.57	1.50
2	B	702	OXN	C11-C10	2.28	1.42	1.38
3	B	602	POP	P2-O	2.41	1.66	1.61
2	A	701	OXN	C23-C22	2.65	1.62	1.48
2	A	701	OXN	C5-C1	2.75	1.64	1.53
2	B	702	OXN	C23-C22	2.79	1.63	1.48

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	702	OXN	C6-C9	3.00	1.58	1.53
2	B	702	OXN	C14-C9	3.09	1.44	1.39
3	B	601	POP	P1-O	3.11	1.68	1.61
3	B	601	POP	P2-O6	3.18	1.61	1.50
2	B	702	OXN	C5-C1	3.19	1.65	1.53
3	B	602	POP	P1-O	3.25	1.68	1.61
2	A	701	OXN	C14-C9	3.42	1.44	1.39
2	A	701	OXN	C10-C9	3.83	1.45	1.39
2	B	702	OXN	C10-C9	4.31	1.46	1.39

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	602	POP	O5-P2-O4	-3.95	96.34	110.85
2	A	701	OXN	C7-C6-C9	-3.86	100.48	110.29
3	B	601	POP	O5-P2-O4	-3.80	96.90	110.85
2	B	702	OXN	C7-C6-C9	-2.90	102.93	110.29
2	B	702	OXN	C2-C1-C4	-2.09	100.01	110.55
2	A	701	OXN	C2-C1-C4	-2.03	100.33	110.55
3	B	601	POP	O5-P2-O	2.32	115.33	104.65
2	A	701	OXN	O21-C22-C23	2.74	122.54	110.36
2	A	701	OXN	C22-O21-C20	2.76	125.15	113.31
2	B	702	OXN	C22-O21-C20	2.86	125.59	113.31
2	B	702	OXN	O21-C22-C23	2.88	123.16	110.36
2	B	702	OXN	C16-O15-C12	8.25	138.22	117.91
2	A	701	OXN	C16-O15-C12	8.32	138.38	117.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	OXN	1	0
3	B	602	POP	2	0
2	B	702	OXN	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	257/257 (100%)	0.25	14 (5%)	29 38	24, 43, 79, 99	25 (9%)
1	B	255/257 (99%)	0.13	16 (6%)	23 31	21, 37, 72, 92	19 (7%)
All	All	512/514 (99%)	0.19	30 (5%)	26 34	21, 41, 76, 99	44 (8%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1562	GLU	4.9
1	A	1681	GLY	4.7
1	B	1727	HIS	4.3
1	A	1698	ILE	4.2
1	A	1563	PHE	4.0
1	A	1775	GLU	3.9
1	A	1604	TYR	3.8
1	B	1678	GLY	3.7
1	A	1595	GLY	3.6
1	A	1700	HIS	3.5
1	B	1595	GLY	3.1
1	B	1692	GLY	3.1
1	B	1597	ILE	3.1
1	A	1727	HIS	3.0
1	B	1747	GLU	3.0
1	A	1699	GLU	2.9
1	B	1683	LYS	2.9
1	A	1618	TYR	2.5
1	B	1682	SER	2.5
1	B	1751	VAL	2.5
1	B	1561	GLU	2.4
1	B	1726	ALA	2.3
1	A	1614	TYR	2.2
1	A	1596	GLN	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	1568	THR	2.1
1	A	1560	LYS	2.1
1	B	1704	LYS	2.1
1	B	1701	GLU	2.0
1	B	1677	THR	2.0
1	B	1636	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	POP	B	602	9/9	0.91	0.24	4.70	55,76,86,95	0
2	OXN	A	701	24/45	0.72	0.23	2.22	58,78,88,89	0
2	OXN	B	702	24/45	0.74	0.21	2.03	37,66,83,86	0
3	POP	B	601	9/9	0.83	0.19	0.53	68,92,93,96	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.