



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 06:55 PM GMT

PDB ID : 1D53
Title : CRYSTAL STRUCTURE AT 1.5 ANGSTROMS RESOLUTION OF D(CGCICICG), AN OCTANUCLEOTIDE CONTAINING INOSINE, AND ITS COMPARISON WITH D(CGCG) AND D(CGCGCG) STRUCTURES
Authors : Kumar, V.D.; Harrison, R.W.; Andrews, L.C.; Weber, I.T.
Deposited on : 1992-11-05
Resolution : 1.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2274 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	2	 100%
1	B	2	 100%
2	C	4	 100%
2	D	4	 100%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 303 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*CP*GP*CP*IP*CP*IP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	2	Total	C	N	O	P	0	0	0
			38	19	8	10	1			
1	B	2	Total	C	N	O	P	0	0	0
			38	19	8	10	1			

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*GP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	4	Total	C	N	O	P	0	0	0
			79	38	16	22	3			
2	D	4	Total	C	N	O	P	0	0	0
			79	38	16	22	3			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	6	Total	O	0	0
			6	6		
3	B	19	Total	O	0	0
			19	19		
3	C	15	Total	O	0	0
			15	15		
3	D	29	Total	O	0	0
			29	29		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: DNA (5'-D(*CP*GP*CP*IP*CP*IP*CP*G)-3')

Chain A:  100%

C1
G2

- Molecule 1: DNA (5'-D(*CP*GP*CP*IP*CP*IP*CP*G)-3')

Chain B:  100%

C3
G4

- Molecule 2: DNA (5'-D(*CP*GP*CP*G)-3')

Chain C:  100%

C5
G6
C7
G8

- Molecule 2: DNA (5'-D(*CP*GP*CP*G)-3')

Chain D:  100%

C9
G10
C11
G12

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	31.00 Å 31.00 Å 43.70 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	5.00 – 1.50	Depositor
% Data completeness (in resolution range)	(Not available) (5.00-1.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR, NUCLSQ	Depositor
R, R_{free}	0.225 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	303	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	3.00	6/42 (14.3%)	6.92	21/63 (33.3%)
1	B	4.05	5/42 (11.9%)	5.97	19/63 (30.2%)
2	C	3.59	13/88 (14.8%)	6.93	61/134 (45.5%)
2	D	3.87	12/88 (13.6%)	6.03	42/134 (31.3%)
All	All	3.68	36/260 (13.8%)	6.48	143/394 (36.3%)

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	11	DC	P-O5'	17.02	1.76	1.59
1	B	4	DG	P-O5'	14.23	1.74	1.59
2	C	6	DG	P-O5'	-12.57	1.47	1.59
2	D	12	DG	N9-C8	-11.71	1.29	1.37
1	B	4	DG	P-OP1	11.57	1.68	1.49

The worst 5 of 143 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	3	DC	P-O3'-C3'	27.15	152.28	119.70
1	A	2	DG	O5'-P-OP2	22.48	137.67	110.70
1	A	2	DG	O5'-P-OP1	-22.28	83.97	110.70
2	C	8	DG	O5'-P-OP1	-21.16	85.31	110.70
2	C	5	DC	N3-C4-C5	-18.83	114.37	121.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	38	0	24	2	0
1	B	38	0	24	14	0
2	C	79	0	46	11	1
2	D	79	0	46	31	3
3	A	6	0	0	1	4
3	B	19	0	0	14	3
3	C	15	0	0	5	3
3	D	29	0	0	31	8
All	All	303	0	140	58	12

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 155.

The worst 5 of 58 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:4:DG:H1'	3:B:42:HOH:O	1.24	1.37
1:B:3:DC:C4	3:B:44:HOH:O	1.67	1.25
2:C:6:DG:H2'	2:C:6:DG:N3	1.51	1.10
2:D:10:DG:O5'	3:D:16:HOH:O	1.73	1.04
1:B:3:DC:C5	3:B:44:HOH:O	1.87	1.01

The worst 5 of 12 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:9:DC:O5'	3:D:77:HOH:O[3_665]	0.95	1.25
2:D:9:DC:C5'	3:D:77:HOH:O[3_665]	1.08	1.12
2:C:8:DG:O3'	3:C:66:HOH:O[3_665]	1.72	0.48
3:A:61:HOH:O	3:D:40:HOH:O[5_455]	1.87	0.33
3:C:39:HOH:O	3:D:50:HOH:O[6_654]	1.91	0.29

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.