



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:43 PM GMT

PDB ID : 4D6N
Title : THE CRYSTAL STRUCTURE OF I-DMOI IN COMPLEX WITH ITS TARGET DNA AT 10 DAYS INCUBATION IN 5MM MG (STATE 7)
Authors : Molina, R.; Stella, S.; Redondo, P.; Gomez, H.; Marcaida, M.J.; Orozco, M.; Prieto, J.; Montoya, G.
Deposited on : 2014-11-13
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1456 (2.38-2.34)
Ramachandran outliers	100387	1435 (2.38-2.34)
Sidechain outliers	100360	1436 (2.38-2.34)
RSRZ outliers	91569	1358 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	199	 2% 79% 12% 8%
1	F	199	 3% 79% 16% 2% 2%
1	K	199	 4% 76% 12% 11%
2	B	14	 79% 21%
2	G	14	 93% 7%
2	L	14	 86% 14%
3	C	11	 55% 36% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	11	
3	M	11	
4	D	15	
4	I	15	
4	N	15	
5	E	10	
5	J	10	
5	O	10	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	ACT	A	1191	-	-	-	X
7	ACT	F	1198	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 7821 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HOMING ENDONUCLEASE I-DMOI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	183	Total	C	N	O	S	0	1	0
			1509	975	274	257	3			
1	F	191	Total	C	N	O	S	0	3	0
			1590	1027	289	271	3			
1	K	177	Total	C	N	O	S	0	2	0
			1474	953	268	250	3			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ALA	-	EXPRESSION TAG	UNP P21505
A	189	ALA	-	EXPRESSION TAG	UNP P21505
A	190	ALA	-	EXPRESSION TAG	UNP P21505
A	191	ALA	-	EXPRESSION TAG	UNP P21505
A	192	LEU	-	EXPRESSION TAG	UNP P21505
A	193	GLU	-	EXPRESSION TAG	UNP P21505
A	194	HIS	-	EXPRESSION TAG	UNP P21505
A	195	HIS	-	EXPRESSION TAG	UNP P21505
A	196	HIS	-	EXPRESSION TAG	UNP P21505
A	197	HIS	-	EXPRESSION TAG	UNP P21505
A	198	HIS	-	EXPRESSION TAG	UNP P21505
A	199	HIS	-	EXPRESSION TAG	UNP P21505
F	1	ALA	-	EXPRESSION TAG	UNP P21505
F	189	ALA	-	EXPRESSION TAG	UNP P21505
F	190	ALA	-	EXPRESSION TAG	UNP P21505
F	191	ALA	-	EXPRESSION TAG	UNP P21505
F	192	LEU	-	EXPRESSION TAG	UNP P21505
F	193	GLU	-	EXPRESSION TAG	UNP P21505
F	194	HIS	-	EXPRESSION TAG	UNP P21505
F	195	HIS	-	EXPRESSION TAG	UNP P21505
F	196	HIS	-	EXPRESSION TAG	UNP P21505
F	197	HIS	-	EXPRESSION TAG	UNP P21505
F	198	HIS	-	EXPRESSION TAG	UNP P21505

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	199	HIS	-	EXPRESSION TAG	UNP P21505
K	1	ALA	-	EXPRESSION TAG	UNP P21505
K	189	ALA	-	EXPRESSION TAG	UNP P21505
K	190	ALA	-	EXPRESSION TAG	UNP P21505
K	191	ALA	-	EXPRESSION TAG	UNP P21505
K	192	LEU	-	EXPRESSION TAG	UNP P21505
K	193	GLU	-	EXPRESSION TAG	UNP P21505
K	194	HIS	-	EXPRESSION TAG	UNP P21505
K	195	HIS	-	EXPRESSION TAG	UNP P21505
K	196	HIS	-	EXPRESSION TAG	UNP P21505
K	197	HIS	-	EXPRESSION TAG	UNP P21505
K	198	HIS	-	EXPRESSION TAG	UNP P21505
K	199	HIS	-	EXPRESSION TAG	UNP P21505

- Molecule 2 is a DNA chain called 5'-D(*DGP*CP*CP*TP*TP*GP*CP*CP*GP*GP*GP*TP*AP*DAP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	14	Total	C	N	O	P	0	0	0
			285	136	53	83	13			
2	G	14	Total	C	N	O	P	0	0	0
			285	136	53	83	13			
2	L	14	Total	C	N	O	P	0	0	0
			285	136	53	83	13			

- Molecule 3 is a DNA chain called 5'-D(*DGP*TP*TP*CP*CP*GP*GP*CP*GP*CP*GP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	11	Total	C	N	O	P	0	0	0
			227	106	41	69	11			
3	H	11	Total	C	N	O	P	0	0	0
			227	106	41	69	11			
3	M	11	Total	C	N	O	P	0	0	0
			227	106	41	69	11			

- Molecule 4 is a DNA chain called 5'-D(*DCP*GP*CP*GP*CP*CP*GP*GP*AP*AP*CP*TP*TP*DAP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	15	Total	C	N	O	P	0	0	0
			302	144	57	87	14			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	I	15	Total	C	N	O	P	0	0	0
			302	144	57	87	14			
4	N	15	Total	C	N	O	P	0	0	0
			302	144	57	87	14			


- Molecule 5 is a DNA chain called 5'-D(*DCP*CP*GP*GP*CP*AP*AP*GP*GP*CP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	10	Total	C	N	O	P	0	0	0
			207	96	42	59	10			
5	J	10	Total	C	N	O	P	0	0	0
			207	96	42	59	10			
5	O	10	Total	C	N	O	P	0	0	0
			207	96	42	59	10			

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Mg	0	0
			2	2		
6	K	2	Total	Mg	0	0
			2	2		
6	F	2	Total	Mg	0	0
			2	2		

- Molecule 7 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 4 2 2	0	0
7	A	1	Total C O 4 2 2	0	0
7	F	1	Total C O 4 2 2	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	59	Total O 59 59	0	0
8	B	7	Total O 7 7	0	0
8	C	9	Total O 9 9	0	0
8	D	4	Total O 4 4	0	0
8	E	1	Total O 1 1	0	0
8	F	32	Total O 32 32	0	0
8	G	1	Total O 1 1	0	0
8	H	6	Total O 6 6	0	0
8	I	4	Total O 4 4	0	0
8	K	31	Total O 31 31	0	0
8	L	4	Total O 4 4	0	0
8	M	4	Total O 4 4	0	0
8	N	3	Total O 3 3	0	0
8	O	2	Total O 2 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HOMING ENDONUCLEASE I-DMOI




- Molecule 1: HOMING ENDONUCLEASE I-DMOI



- Molecule 1: HOMING ENDONUCLEASE I-DMOI



- Molecule 2: 5'-D(*DGP*CP*CP*TP*TP*GP*CP*CP*GP*GP*GP*TP*AP*DAP)-3'




- Molecule 2: 5'-D(*DGP*CP*CP*TP*TP*GP*CP*CP*GP*GP*GP*TP*AP*DAP)-3'

Chain G:  93% 7%




- Molecule 2: 5'-D(*DGP*CP*CP*TP*TP*GP*CP*CP*GP*GP*GP*TP*AP*DAP)-3'

Chain L:  86% 14%




- Molecule 3: 5'-D(*DGP*TP*TP*CP*CP*GP*GP*CP*GP*CP*GP)-3'

Chain C:  55% 36% 9%




- Molecule 3: 5'-D(*DGP*TP*TP*CP*CP*GP*GP*CP*GP*CP*GP)-3'

Chain H:  64% 27% 9%




- Molecule 3: 5'-D(*DGP*TP*TP*CP*CP*GP*GP*CP*GP*CP*GP)-3'

Chain M:  64% 27% 9%




- Molecule 4: 5'-D(*DCP*GP*CP*GP*CP*CP*GP*GP*AP*AP*CP*TP*TP*DAP*C)-3'

Chain D:  87% 13%



- Molecule 4: 5'-D(*DCP*GP*CP*GP*CP*CP*GP*GP*AP*AP*CP*TP*TP*DAP*C)-3'

Chain I:  67% 27% 7%




- Molecule 4: 5'-D(*DCP*GP*CP*GP*CP*CP*GP*GP*AP*AP*CP*TP*TP*DAP*C)-3'

Chain N:  73% 27%




- Molecule 5: 5'-D(*DCP*CP*GP*GP*CP*AP*AP*GP*GP*CP)-3'

Chain E:  90% 10%



- Molecule 5: 5'-D(*DCP*CP*GP*GP*CP*AP*AP*GP*GP*CP)-3'

Chain J:  90% 10%



- Molecule 5: 5'-D(*DCP*CP*GP*GP*CP*AP*AP*GP*GP*CP)-3'

Chain O:  60% 30% 10%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	106.31Å 70.27Å 107.17Å 90.00° 119.59° 90.00°	Depositor
Resolution (Å)	46.60 – 2.35 46.60 – 2.35	Depositor EDS
% Data completeness (in resolution range)	98.8 (46.60-2.35) 98.5 (46.60-2.35)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.99 (at 2.34Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.182 , 0.217 0.200 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	50.2	Xtriage
Anisotropy	0.696	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 42.1	EDS
Estimated twinning fraction	0.025 for -h-l,k,h 0.025 for l,k,-h-l 0.018 for h,-k,-h-l 0.017 for -h-l,-k,l 0.018 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 56740 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7821	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.48	0/1541	0.68	3/2076 (0.1%)
1	F	0.46	0/1631	0.60	1/2197 (0.0%)
1	K	0.44	0/1506	0.63	1/2023 (0.0%)
2	B	0.88	0/319	0.97	0/491
2	G	0.73	0/319	1.00	0/491
2	L	0.88	0/319	0.96	0/491
3	C	1.05	1/253 (0.4%)	1.02	1/387 (0.3%)
3	H	1.32	3/253 (1.2%)	1.13	0/387
3	M	1.10	1/253 (0.4%)	1.01	0/387
4	D	0.87	0/338	0.95	1/519 (0.2%)
4	I	1.17	2/338 (0.6%)	1.33	4/519 (0.8%)
4	N	0.91	0/338	0.99	0/519
5	E	1.06	1/232 (0.4%)	0.85	0/354
5	J	1.01	1/232 (0.4%)	0.88	0/354
5	O	0.97	1/232 (0.4%)	0.87	1/354 (0.3%)
All	All	0.74	10/8104 (0.1%)	0.83	12/11549 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	F	0	1
All	All	0	2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	15	DG	OP3-P	-10.64	1.48	1.61

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	M	15	DG	OP3-P	-10.29	1.48	1.61
5	J	16	DC	OP3-P	-10.26	1.48	1.61
3	C	15	DG	OP3-P	-10.10	1.49	1.61
5	E	16	DC	OP3-P	-9.74	1.49	1.61
5	O	16	DC	OP3-P	-9.10	1.50	1.61
4	I	14	DA	P-O5'	7.17	1.67	1.59
3	H	15	DG	N1-C2	-5.58	1.33	1.37
4	I	14	DA	N7-C5	5.50	1.42	1.39
3	H	15	DG	C8-N7	5.29	1.34	1.30

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	I	14	DA	O5'-P-OP2	-13.46	93.59	105.70
1	A	31	GLY	N-CA-C	-10.04	87.99	113.10
4	I	14	DA	O5'-P-OP1	-9.73	96.94	105.70
1	F	77	ARG	NE-CZ-NH2	6.71	123.66	120.30
1	K	31	GLY	N-CA-C	-6.15	97.72	113.10
4	I	13	DT	OP1-P-O3'	6.05	118.51	105.20
4	D	8	DG	C1'-O4'-C4'	-5.68	104.42	110.10
4	I	13	DT	OP2-P-O3'	5.62	117.56	105.20
5	O	18	DG	O4'-C1'-N9	5.61	111.93	108.00
1	A	77	ARG	NE-CZ-NH1	5.45	123.02	120.30
3	C	24	DC	O4'-C1'-N1	5.20	111.64	108.00
1	A	77	ARG	NE-CZ-NH2	-5.14	117.73	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	30	LYS	Peptide
1	F	182	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1509	0	1578	18	0
1	F	1590	0	1663	27	0
1	K	1474	0	1553	23	0
2	B	285	0	159	3	0
2	G	285	0	159	1	0
2	L	285	0	159	2	0
3	C	227	0	124	5	0
3	H	227	0	124	5	0
3	M	227	0	124	4	0
4	D	302	0	169	1	0
4	I	302	0	169	2	0
4	N	302	0	169	4	0
5	E	207	0	111	0	0
5	J	207	0	111	1	0
5	O	207	0	111	2	0
6	A	2	0	0	0	0
6	F	2	0	0	0	0
6	K	2	0	0	0	0
7	A	8	0	6	0	0
7	F	4	0	3	0	0
8	A	59	0	0	2	0
8	B	7	0	0	1	0
8	C	9	0	0	0	0
8	D	4	0	0	0	0
8	E	1	0	0	0	0
8	F	32	0	0	1	0
8	G	1	0	0	0	0
8	H	6	0	0	0	0
8	I	4	0	0	0	0
8	K	31	0	0	3	0
8	L	4	0	0	1	0
8	M	4	0	0	0	0
8	N	3	0	0	0	0
8	O	2	0	0	0	0
All	All	7821	0	6492	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:6:ASN:HA	1:K:7:VAL:HB	1.45	0.97
1:F:120:LYS:NZ	5:J:16:DC:OP3	1.97	0.94
1:K:30:LYS:NZ	3:M:20:DG:OP2	2.03	0.91
1:A:30:LYS:NZ	3:C:20:DG:OP2	2.08	0.87
1:F:108:ALA:HB2	1:F:183:ASN:ND2	1.97	0.78
1:K:104:ARG:NH1	1:K:182:LEU:O	2.17	0.77
1:K:43:LYS:NZ	2:L:13:DA:OP2	2.16	0.76
1:F:30:LYS:NZ	3:H:20:DG:OP2	2.19	0.73
1:K:33:ARG:NH1	3:M:21:DG:O6	2.22	0.73
1:K:120:LYS:NZ	5:O:16:DC:OP1	2.22	0.72
1:K:21:ASP:OD1	8:K:2004:HOH:O	2.06	0.72
1:F:157:ARG:NH2	2:G:9:DG:N7	2.39	0.71
1:A:104:ARG:NH2	8:A:2040:HOH:O	2.17	0.67
1:F:42:GLN:HG2	1:F:47:LEU:HD23	1.77	0.67
1:F:77:ARG:NH1	3:H:15:DG:O6	2.28	0.65
1:F:60[B]:ILE:HD11	1:F:87:LEU:HD22	1.79	0.64
1:F:33:ARG:NH1	3:H:21:DG:O6	2.30	0.64
4:N:7:DG:H2''	4:N:8:DG:H5''	1.81	0.63
3:M:19:DC:H2'	3:M:20:DG:C8	2.36	0.61
4:I:14:DA:H2''	4:I:15:DC:H5''	1.83	0.59
1:A:77:ARG:HG3	1:A:77:ARG:HH11	1.67	0.59
1:F:108:ALA:HB2	1:F:183:ASN:HD22	1.67	0.57
1:A:97:ARG:NH2	8:A:2034:HOH:O	2.26	0.56
8:K:2031:HOH:O	3:M:15:DG:OP1	2.16	0.56
1:K:30:LYS:O	1:K:33:ARG:NE	2.31	0.56
3:C:19:DC:H2''	3:C:20:DG:H5'	1.88	0.55
1:F:47:LEU:HD12	1:F:51:HIS:HB3	1.88	0.55
1:F:42:GLN:HB3	1:F:48:ILE:HG13	1.90	0.54
1:F:185:LEU:HB3	1:F:190:ALA:HB2	1.90	0.54
2:L:6:DG:OP2	8:L:2002:HOH:O	2.18	0.54
1:F:77:ARG:O	1:F:77:ARG:HG2	2.03	0.53
1:K:97:ARG:HH11	1:K:97:ARG:HG3	1.73	0.53
1:A:77:ARG:HG3	1:A:77:ARG:NH1	2.24	0.53
1:K:73:LYS:HD2	1:K:78:TYR:CZ	2.45	0.52
4:I:2:DG:H2''	4:I:3:DC:H5''	1.90	0.52
1:A:43:LYS:NZ	2:B:13:DA:OP2	2.41	0.52
1:A:117:GLU:OE2	1:A:129:ASN:ND2	2.41	0.51
1:K:158:HIS:ND1	4:N:14:DA:N7	2.57	0.51
1:A:26:LYS:HE2	1:A:28:LYS:NZ	2.25	0.51
1:F:111:LYS:NZ	1:F:180:SER:O	2.42	0.51
1:F:104:ARG:NE	1:F:182:LEU:HB3	2.27	0.50
1:F:172:LYS:O	1:F:176:THR:HB	2.12	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:11:DG:N3	8:B:2006:HOH:O	2.35	0.50
1:F:48:ILE:HD12	1:F:78:TYR:HB2	1.94	0.50
1:K:6:ASN:N	8:K:2001:HOH:O	2.46	0.48
3:H:19:DC:H2'	3:H:20:DG:C8	2.48	0.48
1:F:22:GLY:HA3	1:F:39:VAL:O	2.13	0.48
1:A:60:ILE:HG23	1:A:65:VAL:HB	1.95	0.48
5:O:20:DC:H2''	5:O:21:DA:C8	2.49	0.47
1:F:132:LYS:HD2	1:F:161:TYR:CE2	2.49	0.47
1:A:22:GLY:HA3	1:A:39:VAL:O	2.13	0.47
1:K:73:LYS:HD2	1:K:78:TYR:CE2	2.50	0.47
1:A:47:LEU:HA	1:A:51:HIS:HB2	1.96	0.47
1:K:22:GLY:HA3	1:K:39:VAL:O	2.15	0.47
1:F:182:LEU:HB2	1:F:183:ASN:OD1	2.16	0.46
1:A:30:LYS:O	1:A:30:LYS:HG2	2.15	0.46
1:F:41:THR:HB	1:F:77:ARG:HG3	1.97	0.46
1:F:111:LYS:HE2	1:F:178:LEU:O	2.16	0.46
4:N:8:DG:H2'	4:N:9:DA:C8	2.52	0.45
1:K:157:ARG:C	1:K:158:HIS:HD2	2.21	0.44
1:F:29:TYR:HB2	1:F:33:ARG:HB2	2.00	0.44
1:K:115:VAL:O	1:K:120:LYS:HE2	2.18	0.44
1:F:42:GLN:CG	1:F:47:LEU:HD23	2.47	0.44
1:K:102:ASN:O	1:K:106:GLN:HG3	2.19	0.43
4:N:8:DG:H2''	4:N:9:DA:H5'	2.00	0.43
1:A:106:GLN:HG2	1:A:141:TRP:CE2	2.53	0.43
3:C:19:DC:H2'	3:C:20:DG:C8	2.54	0.43
1:K:6:ASN:HA	1:K:7:VAL:CB	2.27	0.42
1:A:85:LYS:NZ	4:D:4:DG:OP1	2.47	0.42
1:K:48:ILE:HD12	1:K:78:TYR:HB2	2.02	0.42
2:B:14:DA:HO3'	3:C:15:DG:P	2.43	0.42
1:K:97:ARG:HH11	1:K:97:ARG:CG	2.34	0.41
1:A:132:LYS:HD2	1:A:161:TYR:CE2	2.55	0.41
1:A:47:LEU:O	1:A:51:HIS:HB2	2.21	0.41
1:F:65:VAL:HG12	1:F:67:SER:H	1.85	0.41
1:K:157:ARG:CB	1:K:158:HIS:HD2	2.34	0.41
1:F:21:ASP:OD1	8:F:2002:HOH:O	2.21	0.41
1:K:31:GLY:HA2	1:K:32:ASN:HA	1.87	0.41
1:A:33:ARG:NH1	3:C:21:DG:O6	2.53	0.41
1:K:97:ARG:NH1	1:K:97:ARG:CG	2.84	0.40
1:F:30:LYS:HZ1	3:H:20:DG:P	2.35	0.40
1:A:185:LEU:HA	1:A:186:PRO:HD3	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	182/199 (92%)	171 (94%)	11 (6%)	0	100	100
1	F	192/199 (96%)	186 (97%)	5 (3%)	1 (0%)	34	39
1	K	177/199 (89%)	171 (97%)	5 (3%)	1 (1%)	30	34
All	All	551/597 (92%)	528 (96%)	21 (4%)	2 (0%)	39	46

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	K	122	LEU
1	F	122	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/178 (93%)	164 (99%)	1 (1%)	90	96
1	F	174/178 (98%)	173 (99%)	1 (1%)	90	96
1	K	161/178 (90%)	159 (99%)	2 (1%)	78	89
All	All	500/534 (94%)	496 (99%)	4 (1%)	86	94

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	77	ARG
1	K	42	GLN
1	K	51	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	ACT	A	1190	-	1,3,3	0.94	0	0,3,3	0.00	-
7	ACT	A	1191	-	1,3,3	1.45	0	0,3,3	0.00	-
7	ACT	F	1198	-	1,3,3	1.24	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	ACT	A	1190	-	-	0/0/0/0	0/0/0/0
7	ACT	A	1191	-	-	0/0/0/0	0/0/0/0
7	ACT	F	1198	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	183/199 (91%)	0.36	4 (2%) 65 77	29, 43, 68, 89	0
1	F	191/199 (95%)	0.38	5 (2%) 59 71	29, 47, 69, 83	0
1	K	177/199 (88%)	0.43	7 (3%) 42 56	30, 45, 70, 99	0
2	B	14/14 (100%)	-0.46	0 100 100	46, 52, 63, 64	0
2	G	14/14 (100%)	-0.05	0 100 100	46, 70, 85, 85	0
2	L	14/14 (100%)	-0.29	0 100 100	47, 52, 69, 70	0
3	C	11/11 (100%)	-0.32	0 100 100	38, 45, 61, 65	0
3	H	11/11 (100%)	-0.17	0 100 100	37, 46, 59, 61	0
3	M	11/11 (100%)	-0.24	0 100 100	35, 50, 60, 61	0
4	D	15/15 (100%)	-0.08	0 100 100	36, 47, 67, 67	0
4	I	15/15 (100%)	-0.30	0 100 100	34, 50, 69, 71	0
4	N	15/15 (100%)	-0.22	0 100 100	38, 47, 67, 68	0
5	E	10/10 (100%)	-0.47	0 100 100	44, 62, 65, 66	0
5	J	10/10 (100%)	0.80	0 100 100	58, 80, 89, 90	0
5	O	10/10 (100%)	-0.25	0 100 100	47, 62, 71, 74	0
All	All	701/747 (93%)	0.27	16 (2%) 64 76	29, 47, 72, 99	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	187	PRO	5.0
1	F	195	HIS	4.4
1	K	182	LEU	4.2
1	K	75	ASP	4.1
1	K	76	THR	3.3
1	F	192	LEU	3.3
1	F	194	HIS	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	K	180	SER	2.9
1	K	181	HIS	2.6
1	K	74	GLY	2.5
1	A	186	PRO	2.5
1	A	75	ASP	2.4
1	K	51	HIS	2.2
1	F	150	THR	2.2
1	A	152[A]	HIS	2.1
1	F	193	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
7	ACT	A	1191	4/4	0.91	0.23	2.75	54,58,59,60	0
7	ACT	F	1198	4/4	0.88	0.22	2.07	55,59,60,61	0
7	ACT	A	1190	4/4	0.84	0.19	1.31	53,62,64,68	0
6	MG	F	1196	1/1	0.94	0.13	-1.51	40,40,40,40	0
6	MG	A	1189	1/1	0.97	0.09	-2.54	42,42,42,42	0
6	MG	F	1197	1/1	0.86	0.09	-3.81	50,50,50,50	0
6	MG	A	1188	1/1	0.97	0.06	-3.98	46,46,46,46	0
6	MG	K	1184	1/1	0.97	0.05	-5.53	45,45,45,45	0
6	MG	K	1183	1/1	0.97	0.04	-9.68	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.