



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 29, 2016 – 04:17 AM EDT

PDB ID : 4D7K
Title : Crystal structure of N,N-8-amino-8-demethyl-D-riboflavin dimethyltransferase (RosA) from *Streptomyces davawensis*
Authors : Uhl, M.K.; Gruber, K.
Deposited on : 2014-11-25
Resolution : 2.22 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027457
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4405 (2.24-2.20)
Clashscore	102246	5146 (2.24-2.20)
Ramachandran outliers	100387	5065 (2.24-2.20)
Sidechain outliers	100360	5066 (2.24-2.20)
RSRZ outliers	91569	4414 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	<div> <div>88%</div> <div>7% 5%</div> </div>
1	B	353	<div> <div>86%</div> <div>8% 5%</div> </div>
1	C	353	<div> <div>81%</div> <div>13% 5%</div> </div>
1	D	353	<div> <div>8%</div> <div>88%</div> <div>6% 5%</div> </div>
1	E	353	<div> <div>10%</div> <div>83%</div> <div>12% 5%</div> </div>
1	F	353	<div> <div>87%</div> <div>8% 5%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 15787 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SAM-DEPENDENT METHYLTRANSFERASES.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	0	1	0
			2590	1640	461	481	8			
1	B	335	Total	C	N	O	S	0	0	0
			2582	1635	458	481	8			
1	C	335	Total	C	N	O	S	0	0	0
			2582	1635	458	481	8			
1	D	336	Total	C	N	O	S	0	0	0
			2591	1640	459	484	8			
1	E	335	Total	C	N	O	S	0	0	0
			2582	1635	458	481	8			
1	F	337	Total	C	N	O	S	0	0	0
			2591	1640	460	483	8			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	348	HIS	-	EXPRESSION TAG	UNP K4RFM2
A	349	HIS	-	EXPRESSION TAG	UNP K4RFM2
A	350	HIS	-	EXPRESSION TAG	UNP K4RFM2
A	351	HIS	-	EXPRESSION TAG	UNP K4RFM2
A	352	HIS	-	EXPRESSION TAG	UNP K4RFM2
A	353	HIS	-	EXPRESSION TAG	UNP K4RFM2
B	348	HIS	-	EXPRESSION TAG	UNP K4RFM2
B	349	HIS	-	EXPRESSION TAG	UNP K4RFM2
B	350	HIS	-	EXPRESSION TAG	UNP K4RFM2
B	351	HIS	-	EXPRESSION TAG	UNP K4RFM2
B	352	HIS	-	EXPRESSION TAG	UNP K4RFM2
B	353	HIS	-	EXPRESSION TAG	UNP K4RFM2
C	348	HIS	-	EXPRESSION TAG	UNP K4RFM2
C	349	HIS	-	EXPRESSION TAG	UNP K4RFM2
C	350	HIS	-	EXPRESSION TAG	UNP K4RFM2
C	351	HIS	-	EXPRESSION TAG	UNP K4RFM2
C	352	HIS	-	EXPRESSION TAG	UNP K4RFM2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	353	HIS	-	EXPRESSION TAG	UNP K4RFM2
D	348	HIS	-	EXPRESSION TAG	UNP K4RFM2
D	349	HIS	-	EXPRESSION TAG	UNP K4RFM2
D	350	HIS	-	EXPRESSION TAG	UNP K4RFM2
D	351	HIS	-	EXPRESSION TAG	UNP K4RFM2
D	352	HIS	-	EXPRESSION TAG	UNP K4RFM2
D	353	HIS	-	EXPRESSION TAG	UNP K4RFM2
E	348	HIS	-	EXPRESSION TAG	UNP K4RFM2
E	349	HIS	-	EXPRESSION TAG	UNP K4RFM2
E	350	HIS	-	EXPRESSION TAG	UNP K4RFM2
E	351	HIS	-	EXPRESSION TAG	UNP K4RFM2
E	352	HIS	-	EXPRESSION TAG	UNP K4RFM2
E	353	HIS	-	EXPRESSION TAG	UNP K4RFM2
F	348	HIS	-	EXPRESSION TAG	UNP K4RFM2
F	349	HIS	-	EXPRESSION TAG	UNP K4RFM2
F	350	HIS	-	EXPRESSION TAG	UNP K4RFM2
F	351	HIS	-	EXPRESSION TAG	UNP K4RFM2
F	352	HIS	-	EXPRESSION TAG	UNP K4RFM2
F	353	HIS	-	EXPRESSION TAG	UNP K4RFM2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	78	Total O 78 78	0	0
2	B	64	Total O 64 64	0	0
2	C	52	Total O 52 52	0	0
2	D	19	Total O 19 19	0	0
2	E	27	Total O 27 27	0	0
2	F	29	Total O 29 29	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: SAM-DEPENDENT METHYLTRANSFERASES




• Molecule 1: SAM-DEPENDENT METHYLTRANSFERASES




• Molecule 1: SAM-DEPENDENT METHYLTRANSFERASES





• Molecule 1: SAM-DEPENDENT METHYLTRANSFERASES







● Molecule 1: SAM-DEPENDENT METHYLTRANSFERASES



● Molecule 1: SAM-DEPENDENT METHYLTRANSFERASES



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	82.72Å 82.76Å 96.45Å 95.08° 98.67° 114.53°	Depositor
Resolution (Å)	34.77 – 2.22 34.77 – 2.22	Depositor EDS
% Data completeness (in resolution range)	77.4 (34.77-2.22) 73.4 (34.77-2.22)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 2.22Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.195 , 0.248 0.195 , 0.249	Depositor DCC
R_{free} test set	4212 reflections (5.02%)	DCC
Wilson B-factor (Å ²)	41.1	Xtriage
Anisotropy	0.183	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 50.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.023 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15787	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/2646	0.40	0/3592
1	B	0.24	0/2635	0.40	0/3578
1	C	0.23	0/2635	0.39	0/3578
1	D	0.21	0/2644	0.37	0/3590
1	E	0.21	0/2635	0.37	0/3578
1	F	0.22	0/2644	0.39	0/3590
All	All	0.23	0/15839	0.39	0/21506

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2590	0	2557	17	0
1	B	2582	0	2544	20	0
1	C	2582	0	2544	31	0
1	D	2591	0	2550	16	0
1	E	2582	0	2544	25	0
1	F	2591	0	2552	22	0
2	A	78	0	0	1	0
2	B	64	0	0	1	0
2	C	52	0	0	3	0
2	D	19	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	27	0	0	1	0
2	F	29	0	0	3	0
All	All	15787	0	15291	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:99:ARG:NH1	2:F:2008:HOH:O	2.24	0.70
1:A:119:ASP:OD2	1:A:134:ARG:NH1	2.26	0.68
1:D:233:ARG:HD3	1:D:240:VAL:HG12	1.76	0.67
1:C:127:GLY:HA2	1:D:62:ARG:HD2	1.75	0.66
1:C:62:ARG:NH2	1:D:303:LEU:O	2.27	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/353 (95%)	326 (98%)	8 (2%)	0	100	100
1	B	333/353 (94%)	326 (98%)	6 (2%)	1 (0%)	46	50
1	C	333/353 (94%)	324 (97%)	7 (2%)	2 (1%)	30	29
1	D	334/353 (95%)	324 (97%)	10 (3%)	0	100	100
1	E	333/353 (94%)	322 (97%)	11 (3%)	0	100	100
1	F	335/353 (95%)	324 (97%)	11 (3%)	0	100	100
All	All	2002/2118 (94%)	1946 (97%)	53 (3%)	3 (0%)	56	64

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	81	GLU
1	C	80	PRO
1	B	80	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/278 (95%)	262 (100%)	1 (0%)	93	97
1	B	262/278 (94%)	259 (99%)	3 (1%)	80	89
1	C	262/278 (94%)	262 (100%)	0	100	100
1	D	263/278 (95%)	260 (99%)	3 (1%)	80	89
1	E	262/278 (94%)	261 (100%)	1 (0%)	93	97
1	F	262/278 (94%)	262 (100%)	0	100	100
All	All	1574/1668 (94%)	1566 (100%)	8 (0%)	92	96

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	317	ASP
1	E	119	ASP
1	D	275	ASP
1	B	95	SER
1	D	209	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	335/353 (94%)	-0.20	2 (0%) 90 90	27, 51, 88, 136	0
1	B	335/353 (94%)	-0.14	7 (2%) 67 65	29, 51, 88, 133	0
1	C	335/353 (94%)	-0.09	4 (1%) 81 80	32, 58, 99, 166	0
1	D	336/353 (95%)	0.32	28 (8%) 14 13	40, 71, 115, 170	0
1	E	335/353 (94%)	0.54	37 (11%) 7 7	41, 80, 141, 165	0
1	F	337/353 (95%)	-0.11	4 (1%) 81 80	33, 59, 94, 144	0
All	All	2013/2118 (95%)	0.05	82 (4%) 41 39	27, 60, 115, 170	0

The worst 5 of 82 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	264	ARG	8.9
1	D	240	VAL	7.8
1	E	260	ALA	6.3
1	C	256	ASP	5.8
1	E	224	GLN	5.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.