



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 06:58 PM GMT

PDB ID : 1DCT
Title : DNA (CYTOSINE-5) METHYLASE FROM HAEIII COVALENTLY
BOUND TO DNA
Authors : Reinisch, K.M.; Chen, L.; Verdine, G.L.; Lipscomb, W.N.
Deposited on : 1995-05-17
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	F	18	
1	G	18	
2	M	18	
2	N	18	
3	A	324	
3	B	324	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	C49	F	10	X	-	-	-
1	C49	G	10	X	-	-	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6908 atoms, of which 268 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*AP*CP*CP*AP*GP*CP*AP*GP*GP*(C49)P*CP*AP*CP*CP*AP*GP*TP*G)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	F	18	Total	C	F	N	O	P	0	0	0
			367	174	1	73	102	17			
1	G	16	Total	C	F	N	O	P	0	0	0
			327	155	1	65	91	15			

- Molecule 2 is a DNA chain called DNA (5'-D(*TP*CP*AP*CP*TP*GP*GP*TP*GP*GP*(C5M)P*CP*TP*GP*CP*TP*GP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	M	18	Total	C	N	O	P	0	0	0
			368	176	65	110	17			
2	N	17	Total	C	N	O	P	0	0	0
			346	166	60	104	16			

- Molecule 3 is a protein called PROTEIN (MODIFICATION METHYLASE HAEIII).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	A	324	Total	C	H	N	O	S	0	0	0
			2749	1680	134	455	468	12			
3	B	324	Total	C	H	N	O	S	0	0	0
			2749	1680	134	455	468	12			

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		
4	A	1	Total	Ca	0	0
			1	1		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: DNA (5'-D(*AP*CP*CP*AP*GP*CP*AP*GP*GP*(C49)P*CP*AP*CP*CP*AP*GP*TP*G)-3')

Chain F: 




- Molecule 1: DNA (5'-D(*AP*CP*CP*AP*GP*CP*AP*GP*GP*(C49)P*CP*AP*CP*CP*AP*GP*TP*G)-3')

Chain G: 




- Molecule 2: DNA (5'-D(*TP*CP*AP*CP*TP*GP*GP*TP*GP*GP*(C5M)P*CP*TP*GP*CP*TP*GP*G)-3')

Chain M: 



- Molecule 2: DNA (5'-D(*TP*CP*AP*CP*TP*GP*GP*TP*GP*GP*(C5M)P*CP*TP*GP*CP*TP*GP*G)-3')


Chain N: 



- Molecule 3: PROTEIN (MODIFICATION METHYLASE HAEIII)


Chain A: 





• Molecule 3: PROTEIN (MODIFICATION METHYLASE HAEIII)

Chain B: 48% 44% 7%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.57 Å 108.04 Å 155.79 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.80	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-2.80)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.226 , 0.326	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6908	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, C49, 5CM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	F	1.54	2/388 (0.5%)	2.60	46/594 (7.7%)
1	G	1.57	5/343 (1.5%)	2.38	30/525 (5.7%)
2	M	1.73	5/388 (1.3%)	2.43	42/596 (7.0%)
2	N	1.75	6/363 (1.7%)	2.44	38/557 (6.8%)
3	A	0.52	0/2681	0.74	2/3615 (0.1%)
3	B	0.51	0/2681	0.76	3/3615 (0.1%)
All	All	0.89	18/6844 (0.3%)	1.37	161/9502 (1.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	1	2
1	G	1	0
2	M	0	1
3	A	0	1
3	B	0	1
All	All	2	5

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	N	8	DT	C3'-O3'	-8.62	1.32	1.44
2	M	8	DT	C3'-O3'	-8.24	1.33	1.44
1	G	14	DC	C4'-O4'	-6.53	1.38	1.45
1	F	9	DG	C3'-O3'	-6.34	1.35	1.44
2	N	10	DG	C4'-O4'	-6.30	1.38	1.45

The worst 5 of 161 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	2	DC	O4'-C4'-C3'	10.12	112.07	106.00
1	F	13	DC	C1'-O4'-C4'	-9.23	100.87	110.10
1	F	13	DC	O4'-C4'-C3'	9.15	111.49	106.00
1	F	4	DA	N1-C2-N3	-8.98	124.81	129.30
1	G	4	DA	N1-C2-N3	-8.92	124.84	129.30

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	F	10	C49	C1'
1	G	10	C49	C1'

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	A	147	TYR	Sidechain
3	B	38	TYR	Sidechain
1	F	1	DA	Sidechain
1	F	17	DT	Sidechain
2	M	17	DG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	367	0	200	18	0
1	G	327	0	178	10	0
2	M	368	0	207	14	0
2	N	346	0	196	5	0
3	A	2615	134	2606	153	0
3	B	2615	134	2606	160	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
All	All	6640	268	5993	343	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 343 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:4:DA:H2"	1:F:5:DG:C8	2.04	0.93
3:A:202:ASN:HB3	3:A:205:LYS:HB3	1.51	0.92
3:B:240:ALA:HB1	3:B:302:LYS:HA	1.52	0.90
3:B:212:GLU:HG3	3:B:297:LEU:HD21	1.55	0.87
3:B:153:ARG:HG3	3:B:237:THR:HG23	1.60	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	322/324 (99%)	273 (85%)	39 (12%)	10 (3%)	5	17
3	B	322/324 (99%)	283 (88%)	26 (8%)	13 (4%)	4	12
All	All	644/648 (99%)	556 (86%)	65 (10%)	23 (4%)	4	14

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	9	GLY
3	A	80	LEU
3	A	241	SER
3	B	9	GLY
3	B	186	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	281/281 (100%)	262 (93%)	19 (7%)	20	49
3	B	281/281 (100%)	260 (92%)	21 (8%)	17	43
All	All	562/562 (100%)	522 (93%)	40 (7%)	18	46

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	273	ARG
3	B	30	TYR
3	B	274	ARG
3	B	8	SER
3	B	56	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
3	A	262	ASN
3	A	284	GLN
3	B	211	HIS
3	A	244	GLN
3	B	210	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	C49	F	10	1,3	13,22,24	1.16	2 (15%)	17,33,38	4.24	6 (35%)
1	C49	G	10	1,3	13,22,24	1.14	1 (7%)	17,33,38	4.05	7 (41%)
2	5CM	M	11	1,2	13,21,22	0.85	1 (7%)	17,30,33	0.69	0
2	5CM	N	11	1,2	13,21,22	0.90	1 (7%)	17,30,33	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	C49	F	10	1,3	1/1/8/10	0/7/40/46	0/2/2/2
1	C49	G	10	1,3	1/1/8/10	0/7/40/46	0/2/2/2
2	5CM	M	11	1,2	-	0/3/21/22	0/2/2/2
2	5CM	N	11	1,2	-	0/3/21/22	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	N	11	5CM	C6-C5	-2.44	1.33	1.40
2	M	11	5CM	C6-C5	-2.43	1.33	1.40
1	F	10	C49	O3'-C3'	-2.08	1.38	1.43
1	F	10	C49	O4'-C1'	2.18	1.47	1.42
1	G	10	C49	O4'-C1'	2.58	1.48	1.42

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	10	C49	C4'-O4'-C1'	-4.84	97.21	109.46
1	F	10	C49	C4'-O4'-C1'	-4.79	97.34	109.46
1	F	10	C49	C2'-C3'-C4'	-2.94	96.68	102.77
1	G	10	C49	C6-N1-C1'	-2.02	115.10	119.62
1	G	10	C49	O4'-C1'-C2'	3.12	112.51	106.28

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	F	10	C49	C1'
1	G	10	C49	C1'

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	F	10	C49	4	0
1	G	10	C49	4	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.