



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:31 AM GMT

PDB ID : 2DDR
Title : Crystal structure of sphingomyelinase from Bacillus cereus with calcium ion
Authors : Ago, H.; Oda, M.; Takahashi, M.; Tsuge, H.; Ochi, S.; Katunuma, N.; Miyano, M.; Sakurai, J.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2006-02-02
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1199 (1.40-1.40)
Clashscore	102246	1295 (1.40-1.40)
Ramachandran outliers	100387	1259 (1.40-1.40)
Sidechain outliers	100360	1258 (1.40-1.40)
RSRZ outliers	91569	1198 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	<div> <div>3%</div> <div>81% 13% 6%</div> </div>
1	B	306	<div> <div>3%</div> <div>83% 11% 6%</div> </div>
1	C	306	<div> <div>7%</div> <div>85% 12% .</div> </div>
1	D	306	<div> <div>3%</div> <div>84% 12% . .</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10629 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sphingomyelin phosphodiesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	288	Total	C	N	O	S	0	0	0
			2289	1447	382	454	6			
1	B	289	Total	C	N	O	S	0	0	0
			2297	1451	384	456	6			
1	C	298	Total	C	N	O	S	0	0	0
			2356	1485	393	470	8			
1	D	298	Total	C	N	O	S	0	0	0
			2356	1485	393	470	8			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Ca	0	0
			2	2		
2	A	2	Total	Ca	0	0
			2	2		
2	D	2	Total	Ca	0	0
			2	2		
2	C	2	Total	Ca	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	338	Total	O	0	0
			338	338		
3	B	385	Total	O	0	0
			385	385		
3	C	287	Total	O	0	0
			287	287		
3	D	313	Total	O	0	0
			313	313		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spingomyelin phosphodiesterase




- Molecule 1: Spingomyelin phosphodiesterase



- Molecule 1: Spingomyelin phosphodiesterase



- Molecule 1: Spingomyelin phosphodiesterase



N187	E188	M196	I201	M214	A249	E250	H261	I267	E268	N269	K270	V271	L272	W279	H284	F285	Q286	A301	T302	I303	S304	M305	LYS
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----

● ● ●

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	65.43 Å 72.69 Å 77.86 Å 112.19° 89.97° 116.76°	Depositor
Resolution (Å)	19.71 – 1.40 40.54 – 1.40	Depositor EDS
% Data completeness (in resolution range)	93.3 (19.71-1.40) 77.8 (40.54-1.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 1.40 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.213 , 0.234 0.212 , 0.211	Depositor DCC
R_{free} test set	10725 reflections (5.02%)	DCC
Wilson B-factor (Å ²)	13.7	Xtriage
Anisotropy	0.403	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 29.9	EDS
Estimated twinning fraction	0.479 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 213798 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10629	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 81.36 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.6156e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/2343	0.56	0/3187
1	B	0.29	0/2351	0.58	0/3198
1	C	0.28	0/2413	0.57	0/3284
1	D	0.29	0/2413	0.58	0/3284
All	All	0.29	0/9520	0.57	0/12953

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2289	0	2198	24	0
1	B	2297	0	2204	22	0
1	C	2356	0	2258	28	0
1	D	2356	0	2258	29	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	338	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	385	0	0	2	0
3	C	287	0	0	0	0
3	D	313	0	0	3	0
All	All	10629	0	8918	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:LYS:HE2	1:A:142:ASP:HA	1.52	0.92
1:C:139:LYS:HE2	1:C:142:ASP:HA	1.57	0.86
1:D:270:LYS:HE3	1:D:272:LEU:HD21	1.59	0.85
1:D:161:LYS:H	1:D:161:LYS:HD2	1.43	0.81
1:B:139:LYS:HE3	1:B:142:ASP:HA	1.62	0.80
1:C:161:LYS:HD2	1:C:161:LYS:H	1.48	0.78
1:B:68:LYS:HD3	1:B:72:PRO:HA	1.67	0.75
1:C:280:THR:HG21	1:C:287:LYS:HE3	1.69	0.74
1:B:7:ASN:HD22	1:B:140:LYS:HZ1	1.36	0.72
1:B:7:ASN:HD22	1:B:140:LYS:NZ	1.87	0.72
1:D:139:LYS:HE2	1:D:142:ASP:HA	1.74	0.69
1:C:8:ASP:CG	1:C:9:THR:H	1.98	0.65
1:A:16:ASN:HA	1:A:52:ASN:HB2	1.80	0.64
1:A:186:ASN:HD22	1:A:187:ASN:N	1.97	0.63
1:C:16:ASN:HA	1:C:52:ASN:HB2	1.81	0.62
1:D:16:ASN:HD22	1:D:16:ASN:C	2.03	0.62
1:A:139:LYS:HE2	1:A:142:ASP:CA	2.28	0.61
1:C:161:LYS:NZ	1:C:161:LYS:HB3	2.17	0.60
1:C:139:LYS:HD3	1:C:144:PHE:CE2	2.37	0.59
1:D:16:ASN:HA	1:D:52:ASN:HB2	1.84	0.59
1:B:16:ASN:HA	1:B:52:ASN:HB2	1.85	0.59
1:D:161:LYS:CD	1:D:161:LYS:H	2.15	0.58
1:B:163:SER:HB3	1:B:164:PRO:HD3	1.85	0.58
1:B:303:ILE:N	1:B:303:ILE:HD12	2.19	0.57
1:A:17:VAL:HG23	1:A:19:MET:HG3	1.85	0.57
1:C:161:LYS:HD2	1:C:161:LYS:N	2.19	0.57
1:B:186:ASN:HD22	1:B:187:ASN:N	2.03	0.57
1:C:139:LYS:HE2	1:C:142:ASP:CA	2.32	0.56
1:A:303:ILE:HD12	1:A:303:ILE:N	2.22	0.55
1:A:163:SER:HB3	1:A:164:PRO:HD3	1.89	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:68:LYS:HD2	1:D:72:PRO:HA	1.89	0.54
1:A:198:VAL:HG13	1:A:210:GLU:HG2	1.89	0.54
1:D:186:ASN:HD22	1:D:187:ASN:N	2.06	0.54
1:C:198:VAL:HG13	1:C:210:GLU:HG2	1.89	0.53
1:A:139:LYS:HD3	1:A:144:PHE:CE2	2.43	0.53
1:A:244:PHE:HB3	1:A:247:SER:OG	2.09	0.53
1:B:201:ILE:HD12	1:B:250:GLU:HA	1.91	0.53
1:A:284:TRP:O	1:A:286:GLN:HG2	2.09	0.53
1:C:119:PHE:CD1	1:C:132:GLY:HA2	2.45	0.52
1:C:161:LYS:HB3	1:C:161:LYS:HZ2	1.75	0.51
1:D:303:ILE:N	1:D:303:ILE:HD12	2.26	0.51
1:D:161:LYS:N	1:D:161:LYS:HD2	2.20	0.51
1:C:68:LYS:HD2	1:C:72:PRO:HA	1.93	0.51
1:B:122:GLY:HA2	1:B:167:VAL:HG11	1.93	0.51
1:D:196:MET:HG2	1:D:214:MET:SD	2.51	0.50
1:A:119:PHE:CD1	1:A:132:GLY:HA2	2.47	0.49
1:D:119:PHE:CD1	1:D:132:GLY:HA2	2.48	0.49
1:A:186:ASN:HD22	1:A:186:ASN:C	2.14	0.48
1:B:119:PHE:CD1	1:B:132:GLY:HA2	2.49	0.48
1:D:131:LYS:HE2	3:D:1358:HOH:O	2.12	0.48
1:C:244:PHE:HB3	1:C:247:SER:OG	2.14	0.48
1:D:8:ASP:OD1	1:D:9:THR:N	2.46	0.47
1:B:33:ARG:HD3	1:B:279:TRP:CZ2	2.49	0.47
1:D:201:ILE:HD12	1:D:250:GLU:HA	1.95	0.47
1:C:8:ASP:CG	1:C:9:THR:N	2.67	0.47
1:A:68:LYS:HD2	1:A:72:PRO:HA	1.96	0.47
1:A:54:VAL:HG22	1:A:102:GLY:O	2.15	0.47
1:D:167:VAL:O	1:D:171:GLN:HG3	2.15	0.47
1:A:167:VAL:O	1:A:171:GLN:HG3	2.14	0.47
1:C:161:LYS:CD	1:C:161:LYS:H	2.24	0.46
1:C:280:THR:CG2	1:C:287:LYS:HG3	2.46	0.46
1:A:131:LYS:HE2	3:A:1338:HOH:O	2.15	0.46
1:A:33:ARG:HD3	1:A:279:TRP:CZ2	2.50	0.46
1:C:119:PHE:CE1	1:C:132:GLY:HA2	2.51	0.46
1:D:119:PHE:CE1	1:D:132:GLY:HA2	2.51	0.46
1:D:201:ILE:HD12	1:D:249:ALA:O	2.16	0.46
1:C:303:ILE:HD12	1:C:303:ILE:N	2.31	0.45
1:D:139:LYS:HD3	1:D:144:PHE:CE2	2.52	0.45
1:B:167:VAL:O	1:B:171:GLN:HG3	2.17	0.45
1:B:268:GLU:O	1:B:301:ALA:HA	2.16	0.45
1:D:139:LYS:HE2	1:D:142:ASP:CA	2.42	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:119:PHE:CE1	1:B:132:GLY:HA2	2.52	0.45
1:B:131:LYS:HE3	3:B:1346:HOH:O	2.16	0.44
1:C:54:VAL:HG22	1:C:102:GLY:O	2.17	0.44
1:C:17:VAL:HG23	1:C:19:MET:HG3	1.98	0.44
1:A:268:GLU:O	1:A:301:ALA:HA	2.17	0.44
1:D:54:VAL:HG22	1:D:102:GLY:O	2.18	0.43
1:B:54:VAL:HG22	1:B:102:GLY:O	2.18	0.43
1:D:270:LYS:HE3	1:D:272:LEU:CD2	2.41	0.43
1:B:186:ASN:HD22	1:B:186:ASN:C	2.21	0.43
1:A:119:PHE:CE1	1:A:132:GLY:HA2	2.54	0.43
1:B:140:LYS:HZ2	1:B:141:ASN:HD22	1.67	0.42
1:D:23:ASN:ND2	3:D:1447:HOH:O	2.52	0.42
1:C:201:ILE:HG22	1:C:202:ASN:ND2	2.34	0.42
1:A:142:ASP:OD1	1:A:142:ASP:O	2.38	0.42
1:B:140:LYS:NZ	1:B:141:ASN:ND2	2.68	0.42
1:C:268:GLU:O	1:C:301:ALA:HA	2.19	0.42
1:D:188:GLU:O	1:D:261:HIS:HB3	2.19	0.41
1:C:167:VAL:O	1:C:171:GLN:HG3	2.19	0.41
1:A:185:PRO:HB2	1:A:188:GLU:HG3	2.02	0.41
1:C:142:ASP:OD1	1:C:142:ASP:O	2.37	0.41
1:D:173:LYS:HD3	3:D:1383:HOH:O	2.21	0.41
1:B:23:ASN:ND2	3:B:1506:HOH:O	2.54	0.41
1:C:33:ARG:HD3	1:C:279:TRP:CZ2	2.56	0.41
1:A:32:GLU:HG3	3:A:1586:HOH:O	2.19	0.41
1:D:186:ASN:C	1:D:186:ASN:HD22	2.23	0.41
1:B:109:TRP:CE2	1:B:140:LYS:HG3	2.56	0.41
1:D:267:ILE:HG13	1:D:303:ILE:HG13	2.03	0.41
1:D:33:ARG:HD3	1:D:279:TRP:CZ2	2.56	0.41
1:D:268:GLU:O	1:D:301:ALA:HA	2.20	0.41
1:C:161:LYS:HG2	1:C:162:THR:N	2.35	0.40
1:C:280:THR:HG21	1:C:287:LYS:HG3	2.03	0.40
1:A:7:ASN:HA	3:A:1632:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/306 (92%)	272 (96%)	10 (4%)	0	100	100
1	B	283/306 (92%)	272 (96%)	11 (4%)	0	100	100
1	C	296/306 (97%)	285 (96%)	11 (4%)	0	100	100
1	D	296/306 (97%)	283 (96%)	13 (4%)	0	100	100
All	All	1157/1224 (94%)	1112 (96%)	45 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/270 (94%)	248 (98%)	6 (2%)	57	19
1	B	255/270 (94%)	249 (98%)	6 (2%)	57	19
1	C	262/270 (97%)	255 (97%)	7 (3%)	52	15
1	D	262/270 (97%)	254 (97%)	8 (3%)	47	12
All	All	1033/1080 (96%)	1006 (97%)	27 (3%)	54	16

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	TRP
1	A	92	ASN
1	A	176	GLN
1	A	181	ASN
1	A	186	ASN
1	A	269	ASN
1	B	28	TRP
1	B	176	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	181	ASN
1	B	186	ASN
1	B	204	GLU
1	B	269	ASN
1	C	28	TRP
1	C	92	ASN
1	C	161	LYS
1	C	176	GLN
1	C	181	ASN
1	C	183	ASN
1	C	269	ASN
1	D	16	ASN
1	D	28	TRP
1	D	161	LYS
1	D	176	GLN
1	D	181	ASN
1	D	186	ASN
1	D	196	MET
1	D	269	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (46) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	ASN
1	A	74	GLN
1	A	92	ASN
1	A	176	GLN
1	A	181	ASN
1	A	183	ASN
1	A	186	ASN
1	A	219	ASN
1	A	243	ASN
1	A	261	HIS
1	A	273	GLN
1	B	7	ASN
1	B	23	ASN
1	B	74	GLN
1	B	92	ASN
1	B	127	ASN
1	B	141	ASN
1	B	176	GLN
1	B	181	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	183	ASN
1	B	186	ASN
1	B	243	ASN
1	B	278	GLN
1	C	23	ASN
1	C	74	GLN
1	C	92	ASN
1	C	176	GLN
1	C	181	ASN
1	C	183	ASN
1	C	202	ASN
1	C	207	ASN
1	C	243	ASN
1	C	273	GLN
1	C	278	GLN
1	D	16	ASN
1	D	23	ASN
1	D	74	GLN
1	D	92	ASN
1	D	141	ASN
1	D	176	GLN
1	D	181	ASN
1	D	183	ASN
1	D	186	ASN
1	D	207	ASN
1	D	243	ASN
1	D	273	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	288/306 (94%)	0.06	10 (3%) 48 44	7, 15, 31, 42	0
1	B	289/306 (94%)	-0.06	9 (3%) 52 49	7, 14, 26, 39	0
1	C	298/306 (97%)	0.16	20 (6%) 21 18	7, 15, 38, 59	0
1	D	298/306 (97%)	0.00	10 (3%) 49 45	7, 14, 31, 52	0
All	All	1173/1224 (95%)	0.04	49 (4%) 40 38	7, 15, 31, 59	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	285	PHE	7.5
1	A	285	PHE	7.2
1	B	285	PHE	6.0
1	D	285	PHE	5.7
1	C	161	LYS	4.3
1	B	127	ASN	4.1
1	C	284	TRP	4.0
1	C	142	ASP	3.9
1	A	286	GLN	3.8
1	D	142	ASP	3.7
1	C	207	ASN	3.6
1	A	208	ASP	3.5
1	A	142	ASP	3.2
1	D	284	TRP	3.2
1	D	161	LYS	3.1
1	A	7	ASN	3.0
1	C	202	ASN	3.0
1	B	286	GLN	2.8
1	D	141	ASN	2.8
1	A	284	TRP	2.8
1	A	207	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	286	GLN	2.6
1	D	286	GLN	2.6
1	B	142	ASP	2.6
1	C	125	PRO	2.6
1	C	208	ASP	2.5
1	C	244	PHE	2.5
1	B	205	ASN	2.5
1	C	158	MET	2.5
1	C	206	ASN	2.5
1	B	170	ASN	2.4
1	A	122	GLY	2.4
1	D	124	GLY	2.4
1	D	121	LYS	2.3
1	B	287	LYS	2.3
1	D	158	MET	2.3
1	A	141	ASN	2.3
1	D	170	ASN	2.2
1	C	128	LEU	2.2
1	C	126	ASP	2.2
1	B	186	ASN	2.2
1	C	170	ASN	2.2
1	B	219	ASN	2.2
1	C	162	THR	2.1
1	C	246	ASP	2.1
1	C	122	GLY	2.1
1	C	141	ASN	2.1
1	A	202	ASN	2.0
1	C	219	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	CA	A	1324	1/1	1.00	0.04	-3.52	10,10,10,10	0
2	CA	B	1326	1/1	1.00	0.04	-4.38	10,10,10,10	0
2	CA	C	1328	1/1	1.00	0.03	-5.17	10,10,10,10	0
2	CA	D	1330	1/1	1.00	0.03	-5.18	11,11,11,11	0
2	CA	B	1327	1/1	0.99	0.08	-	19,19,19,19	0
2	CA	C	1329	1/1	0.99	0.07	-	20,20,20,20	0
2	CA	D	1331	1/1	0.99	0.06	-	20,20,20,20	0
2	CA	A	1325	1/1	0.98	0.07	-	20,20,20,20	0

6.5 Other polymers

There are no such residues in this entry.