



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:34 AM GMT

PDB ID : 2DKJ  
Title : Crystal Structure of T.th.HB8 Serine Hydroxymethyltransferase  
Authors : Kai, K.; Goto, M.; Miyahara, I.; Hirotsu, K.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2006-04-11  
Resolution : 1.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 1.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1495 (1.20-1.08)
Clashscore	102246	1593 (1.20-1.08)
Ramachandran outliers	100387	1521 (1.20-1.08)
Sidechain outliers	100360	1518 (1.20-1.08)
RSRZ outliers	91569	1498 (1.20-1.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	407	 3% 86% 12% ..
1	B	407	 9% 84% 14% ..

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7077 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called serine hydroxymethyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	402	Total	C	N	O	S	0	0	0
			3077	1959	548	561	9			
1	B	402	Total	C	N	O	S	0	0	0
			3064	1948	542	565	9			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C<sub>8</sub>H<sub>10</sub>NO<sub>6</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
3	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	473	Total	O	0	0
			473	473		
4	B	423	Total	O	0	0
			423	423		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: serine hydroxymethyltransferase



- Molecule 1: serine hydroxymethyltransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.99Å 82.88Å 94.07Å 90.00° 91.67° 90.00°	Depositor
Resolution (Å)	42.00 – 1.15 42.81 – 1.15	Depositor EDS
% Data completeness (in resolution range)	95.2 (42.00-1.15) 95.8 (42.81-1.15)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.03 (at 1.15Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.180 , 0.190 0.180 , 0.190	Depositor DCC
$R_{free}$ test set	30095 reflections (9.99%)	DCC
Wilson B-factor (Å <sup>2</sup> )	8.3	Xtriage
Anisotropy	0.161	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 43.5	EDS
Estimated twinning fraction	0.016 for h,-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 301336 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7077	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/3143	0.60	0/4262
1	B	0.27	0/3130	0.58	0/4249
All	All	0.28	0/6273	0.59	0/8511

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3077	0	3090	48	0
1	B	3064	0	3045	57	0
2	A	5	0	0	1	0
2	B	5	0	0	1	0
3	A	15	0	7	2	0
3	B	15	0	7	1	0
4	A	473	0	0	5	0
4	B	423	0	0	8	0
All	All	7077	0	6149	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:194:LEU:H	1:B:218:HIS:HD2	1.11	0.97
1:A:194:LEU:H	1:A:218:HIS:HD2	1.09	0.91
1:B:194:LEU:H	1:B:218:HIS:CD2	2.00	0.80
1:B:92:HIS:H	1:B:256:GLN:HE22	1.31	0.77
1:B:262:HIS:H	1:B:262:HIS:CD2	2.03	0.76
1:A:92:HIS:H	1:A:256:GLN:HE22	1.33	0.76
1:B:291:LYS:HG3	4:B:1762:HOH:O	1.85	0.76
1:A:194:LEU:H	1:A:218:HIS:CD2	2.00	0.75
1:A:262:HIS:CD2	1:A:262:HIS:H	2.04	0.74
1:B:23:ARG:HA	1:B:406:MET:CE	2.19	0.71
1:A:144:ARG:HG3	1:A:151:ASP:HB2	1.75	0.69
1:B:200:HIS:HE1	2:B:1520:SO4:O4	1.76	0.68
1:A:246:ARG:HG3	4:A:2846:HOH:O	2.00	0.62
1:B:319:ARG:HB2	1:B:320:PRO:HD3	1.83	0.61
1:B:200:HIS:HD2	1:B:312:HIS:NE2	1.99	0.60
1:B:165:LYS:HD2	4:B:1815:HOH:O	2.02	0.60
1:B:344:ILE:O	1:B:347:ASP:HB2	2.02	0.58
1:B:280:LYS:HG3	4:B:1909:HOH:O	2.04	0.58
1:A:400:LEU:O	1:A:404:HIS:HD2	1.85	0.58
1:A:200:HIS:HD2	1:A:312:HIS:NE2	2.01	0.57
1:B:120:GLY:O	1:B:171:ALA:HB1	2.05	0.56
1:B:212:ASN:ND2	1:B:215:PRO:HD3	2.20	0.56
1:B:342:ASN:HD22	1:B:343:ALA:H	1.54	0.56
1:B:305:VAL:HG12	1:B:306:THR:HG23	1.88	0.56
1:B:226:LYS:NZ	3:B:510:PLP:O3	2.36	0.55
1:A:24:GLU:HB3	1:A:398:ARG:HG3	1.89	0.55
1:A:226:LYS:NZ	3:A:510:PLP:O3	2.39	0.55
1:A:225:HIS:CD2	1:A:232:ARG:HA	2.42	0.55
1:B:342:ASN:HD22	1:B:343:ALA:N	2.06	0.54
1:A:120:GLY:O	1:A:171:ALA:HB1	2.08	0.54
1:A:406:MET:HE2	4:A:2728:HOH:O	2.07	0.53
1:B:122:HIS:H	1:B:125:HIS:CD2	2.26	0.53
1:B:225:HIS:CD2	1:B:232:ARG:HA	2.43	0.53
1:B:324:THR:OG1	1:B:327:GLU:HG3	2.08	0.53
1:A:262:HIS:H	1:A:262:HIS:HD2	1.55	0.53
1:A:122:HIS:H	1:A:125:HIS:CD2	2.26	0.52
1:B:262:HIS:H	1:B:262:HIS:HD2	1.57	0.52
1:B:22:GLN:O	1:B:406:MET:HE1	2.10	0.51
1:A:92:HIS:H	1:A:256:GLN:NE2	2.05	0.51
1:B:23:ARG:HA	1:B:406:MET:HE2	1.92	0.51

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:134:LYS:HD3	4:B:1749:HOH:O	2.10	0.51
1:B:225:HIS:O	1:B:226:LYS:HB2	2.10	0.51
1:B:284:ARG:NH1	1:B:288:GLU:OE2	2.44	0.50
1:A:291:LYS:O	1:A:295:GLU:HG3	2.12	0.49
1:A:312:HIS:HD1	1:A:312:HIS:H	1.60	0.49
1:B:123:LEU:HD12	1:B:124:THR:HG23	1.93	0.49
1:B:225:HIS:HD2	1:B:233:GLY:H	1.61	0.49
1:A:225:HIS:O	1:A:226:LYS:HB2	2.12	0.49
1:B:125:HIS:HE1	1:B:197:ASP:OD2	1.96	0.49
1:A:181:LYS:O	1:A:185:GLU:HG3	2.12	0.49
1:B:9:GLU:O	1:B:13:GLU:HG3	2.12	0.49
1:B:313:LEU:C	1:B:313:LEU:HD12	2.34	0.48
1:B:92:HIS:N	1:B:256:GLN:HE22	2.07	0.48
1:B:312:HIS:HD1	1:B:312:HIS:H	1.61	0.48
1:A:262:HIS:HE1	4:B:1527:HOH:O	1.96	0.48
1:B:92:HIS:H	1:B:256:GLN:NE2	2.04	0.48
1:B:319:ARG:HG3	1:B:355:SER:HB3	1.96	0.48
1:B:123:LEU:HD12	1:B:124:THR:N	2.28	0.48
1:B:40:ARG:HD2	4:B:1649:HOH:O	2.14	0.48
1:A:139:VAL:HG11	1:A:162:HIS:CG	2.49	0.48
1:A:326:LYS:O	1:A:330:GLU:HG3	2.14	0.47
1:A:225:HIS:CD2	1:A:225:HIS:H	2.31	0.47
1:B:225:HIS:CD2	1:B:233:GLY:H	2.32	0.47
1:B:24:GLU:HB3	1:B:398:ARG:HG3	1.95	0.47
1:A:225:HIS:HE1	1:B:51:TYR:OH	1.98	0.46
1:B:262:HIS:N	1:B:262:HIS:CD2	2.78	0.46
1:B:225:HIS:H	1:B:225:HIS:CD2	2.34	0.46
1:A:262:HIS:N	1:A:262:HIS:CD2	2.78	0.46
1:B:31:SER:HB2	1:B:226:LYS:HE2	1.98	0.46
1:B:129:VAL:C	1:B:134:LYS:HE2	2.37	0.45
1:A:144:ARG:CG	1:A:151:ASP:HB2	2.45	0.45
1:B:30:ALA:HA	1:B:360:GLY:HA3	1.97	0.45
1:A:361:THR:N	1:A:362:PRO:CD	2.80	0.45
1:B:361:THR:N	1:B:362:PRO:CD	2.79	0.45
1:B:400:LEU:O	1:B:404:HIS:HD2	2.00	0.45
1:A:91:PRO:HA	1:A:256:GLN:HE22	1.82	0.44
1:B:326:LYS:O	1:B:330:GLU:HG3	2.18	0.44
1:A:51:TYR:OH	1:B:225:HIS:HE1	2.00	0.44
1:B:80:ALA:HB2	4:B:1916:HOH:O	2.17	0.44
1:A:30:ALA:HA	1:A:360:GLY:HA3	1.99	0.44
4:A:2524:HOH:O	1:B:262:HIS:HE1	1.99	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:92:HIS:N	1:A:256:GLN:HE22	2.09	0.44
1:A:305:VAL:HG12	1:A:306:THR:HG23	2.00	0.43
1:A:214:LEU:HD23	4:A:2658:HOH:O	2.18	0.43
1:A:197:ASP:OD2	3:A:510:PLP:N1	2.51	0.43
1:A:172:SER:HB3	2:A:2520:SO4:O1	2.19	0.43
1:A:334:ALA:HB3	1:A:389:PRO:HB3	1.99	0.43
1:B:91:PRO:HA	1:B:256:GLN:HE22	1.82	0.43
1:A:147:THR:O	1:A:148:GLU:HB2	2.19	0.43
1:A:274:ALA:HA	1:A:279:PHE:CG	2.54	0.43
1:A:295:GLU:HG3	4:A:2572:HOH:O	2.19	0.42
1:A:139:VAL:HG11	1:A:162:HIS:CB	2.50	0.42
1:A:313:LEU:C	1:A:313:LEU:HD12	2.39	0.42
1:B:406:MET:HG3	1:B:407:PRO:HD2	2.02	0.41
1:A:125:HIS:HE1	1:A:197:ASP:OD2	2.03	0.41
1:A:335:VAL:HG11	1:A:393:LEU:HD23	2.02	0.41
1:A:139:VAL:HG11	1:A:162:HIS:HB3	2.02	0.41
1:B:361:THR:N	1:B:362:PRO:HD3	2.36	0.41
1:B:15:ILE:O	1:B:19:GLU:HG3	2.20	0.41
1:B:406:MET:HG2	4:B:1674:HOH:O	2.19	0.41
1:A:113:MET:HA	1:A:139:VAL:O	2.21	0.41
1:A:225:HIS:CD2	1:A:233:GLY:H	2.40	0.40
1:A:361:THR:N	1:A:362:PRO:HD3	2.35	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/407 (98%)	392 (98%)	7 (2%)	1 (0%)	46	17
1	B	400/407 (98%)	390 (98%)	9 (2%)	1 (0%)	46	17
All	All	800/814 (98%)	782 (98%)	16 (2%)	2 (0%)	46	17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	226	LYS
1	B	226	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/325 (96%)	310 (99%)	3 (1%)	82	52
1	B	310/325 (95%)	307 (99%)	3 (1%)	82	52
All	All	623/650 (96%)	617 (99%)	6 (1%)	82	52

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	TYR
1	A	200	HIS
1	A	262	HIS
1	B	51	TYR
1	B	200	HIS
1	B	342	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	GLN
1	A	125	HIS
1	A	200	HIS
1	A	218	HIS
1	A	225	HIS
1	A	256	GLN
1	A	262	HIS
1	A	404	HIS
1	B	96	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	125	HIS
1	B	200	HIS
1	B	218	HIS
1	B	225	HIS
1	B	256	GLN
1	B	262	HIS
1	B	342	ASN
1	B	404	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	2520	-	4,4,4	3.16	2 (50%)	6,6,6	0.93	0
3	PLP	A	510	1	15,15,16	1.80	2 (13%)	21,22,23	2.68	5 (23%)
2	SO4	B	1520	-	4,4,4	3.05	2 (50%)	6,6,6	0.94	0
3	PLP	B	510	1	15,15,16	1.77	1 (6%)	21,22,23	2.35	6 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	2520	-	-	0/0/0/0	0/0/0/0
3	PLP	A	510	1	-	0/6/6/8	0/1/1/1
2	SO4	B	1520	-	-	0/0/0/0	0/0/0/0
3	PLP	B	510	1	-	0/6/6/8	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	510	PLP	C3-C2	-5.66	1.36	1.40
3	A	510	PLP	C3-C2	-5.58	1.36	1.40
2	A	2520	SO4	O3-S	-4.50	1.31	1.47
2	B	1520	SO4	O3-S	-4.34	1.31	1.47
3	A	510	PLP	C2A-C2	2.29	1.54	1.50
2	B	1520	SO4	O1-S	4.09	1.61	1.47
2	A	2520	SO4	O1-S	4.34	1.62	1.47

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	510	PLP	C4A-C4-C3	-2.17	116.42	120.36
3	B	510	PLP	O2P-P-O4P	-2.16	100.33	106.56
3	A	510	PLP	O2P-P-O4P	-2.16	100.34	106.56
3	B	510	PLP	C5A-C5-C6	-2.04	115.43	119.28
3	B	510	PLP	C4A-C4-C3	-2.03	116.69	120.36
3	B	510	PLP	C4A-C4-C5	3.01	124.02	120.88
3	B	510	PLP	C5A-C5-C4	3.15	125.83	121.65
3	A	510	PLP	C5A-C5-C4	3.16	125.83	121.65
3	A	510	PLP	C4A-C4-C5	3.36	124.38	120.88
3	B	510	PLP	O4P-C5A-C5	8.45	122.96	108.99
3	A	510	PLP	O4P-C5A-C5	10.14	125.75	108.99

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2520	SO4	1	0
3	A	510	PLP	2	0
2	B	1520	SO4	1	0
3	B	510	PLP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	402/407 (98%)	0.46	11 (2%) 58 56	5, 9, 16, 22	0
1	B	402/407 (98%)	0.71	38 (9%) 10 12	5, 9, 23, 30	0
All	All	804/814 (98%)	0.58	49 (6%) 25 24	5, 9, 19, 30	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	405	PRO	6.7
1	B	323	LEU	6.1
1	B	348	PRO	5.5
1	B	344	ILE	5.5
1	B	407	PRO	5.4
1	A	407	PRO	5.2
1	B	391	GLU	5.1
1	B	389	PRO	5.0
1	B	319	ARG	4.5
1	B	322	GLY	4.5
1	B	386	LEU	4.2
1	B	406	MET	4.0
1	A	406	MET	3.7
1	B	395	GLU	3.4
1	B	327	GLU	3.3
1	B	387	GLU	3.3
1	B	353	VAL	3.3
1	B	388	GLY	3.3
1	B	324	THR	3.1
1	B	351	PRO	3.0
1	B	347	ASP	3.0
1	B	350	PRO	3.0
1	B	326	LYS	2.9
1	A	405	PRO	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	390	SER	2.6
1	B	146	ASP	2.6
1	B	331	ARG	2.6
1	B	118	ALA	2.5
1	A	119	ALA	2.5
1	B	392	ALA	2.5
1	A	139	VAL	2.4
1	B	299	ARG	2.4
1	B	394	ARG	2.4
1	B	334	ALA	2.4
1	B	51	TYR	2.4
1	B	354	THR	2.4
1	A	118	ALA	2.4
1	B	330	GLU	2.3
1	A	216	TYR	2.3
1	B	61	TYR	2.3
1	B	320	PRO	2.3
1	B	60	TYR	2.2
1	A	387	GLU	2.2
1	A	117	LEU	2.1
1	B	241	PRO	2.1
1	A	251	ILE	2.1
1	A	350	PRO	2.1
1	B	321	LYS	2.0
1	B	346	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors



of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	2520	5/5	0.95	0.15	1.62	15,16,17,17	0
3	PLP	A	510	15/16	0.96	0.13	1.52	6,13,16,17	0
3	PLP	B	510	15/16	0.97	0.12	1.21	5,10,12,13	0
2	SO4	B	1520	5/5	0.96	0.09	0.09	10,12,13,14	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.