



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:49 PM GMT

PDB ID : 4DKO  
Title : Crystal structure of clade A/E 93TH057 HIV-1 gp120 core in complex with TS-II-224  
Authors : Kwon, Y.D.; LaLonde, J.M.; Jones, D.M.; Sun, A.W.; Courter, J.R.; Soeta, T.; Kobayashi, T.; Princiotta, A.M.; Wu, X.; Mascola, J.; Schon, A.; Freire, E.; Sodroski, J.; Madani, N.; Smith III, A.B.; Kwong, P.D.  
Deposited on : 2012-02-03  
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865

**i**

## X-RAY DIFFRACTION

A.



Similar resolution  
(#Entries, resolution range(Å))

Quality of chain

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	506	-	-	-	X
2	NAG	C	511	-	-	-	X
4	OLM	A	513	-	-	-	X
4	OLM	C	513	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6090 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HIV-1 gp120 core.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	339	Total	C	N	O	S	0	0	0
			2654	1666	460	507	21			
1	C	339	Total	C	N	O	S	0	0	0
			2654	1666	460	507	21			

- Molecule 2 is TS-II-224 (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C<sub>8</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
3	C	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 4 is N-(4-CHLORO-3-FLUOROPHENYL)-N'-(1,2,2,6,6-PENTAMETHYLPIPERIDIN-4-YL)ETHANEDIAMIDE (three-letter code: OLM) (formula:  $C_{18}H_{25}ClF_2N_3O_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Cl	F	N	O	
			25	18	1	1	3	2	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	C	1	Total	C	Cl	F	N	O	0	0
			25	18	1	1	3	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	243	Total	O	0	0
			243	243		
5	C	151	Total	O	0	0
			151	151		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HIV-1 gp120 core



- Molecule 1: HIV-1 gp120 core





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.72Å 68.90Å 94.52Å 90.00° 91.23° 90.00°	Depositor
Resolution (Å)	41.94 – 1.98 47.17 – 1.98	Depositor EDS
% Data completeness (in resolution range)	93.1 (41.94-1.98) 87.9 (47.17-1.98)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.89 (at 1.98Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, $R_{free}$	0.203 , 0.239 0.202 , 0.233	Depositor DCC
$R_{free}$ test set	2697 reflections (5.57%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.0	Xtriage
Anisotropy	0.393	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 51.4	EDS
Estimated twinning fraction	0.026 for h,-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 53999 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6090	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 0LM, EPE, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.21	0/2709	0.40	0/3678
1	C	0.21	0/2709	0.39	0/3678
All	All	0.21	0/5418	0.39	0/7356

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2654	0	2589	17	0
1	C	2654	0	2589	17	0
2	A	154	0	143	1	0
2	C	154	0	143	1	0
3	A	15	0	17	0	0
3	C	15	0	17	0	0
4	A	25	0	25	1	0
4	C	25	0	25	1	0
5	A	243	0	0	8	0
5	C	151	0	0	9	0
All	All	6090	0	5548	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:391:PHE:O	5:A:672:HOH:O	1.85	0.92
1:C:59:LYS:NZ	5:C:739:HOH:O	2.04	0.91
1:A:471:GLY:O	5:A:698:HOH:O	1.96	0.82
1:C:490:GLN:O	5:C:726:HOH:O	2.08	0.71
1:C:411:ASN:N	5:C:680:HOH:O	2.24	0.70
1:A:396:ILE:O	5:A:829:HOH:O	2.12	0.66
1:C:231:LYS:HB3	1:C:268:GLU:HG3	1.78	0.65
1:A:121:LYS:HB2	1:A:201:ILE:HB	1.80	0.63
1:A:480:ARG:NH2	5:A:683:HOH:O	2.36	0.58
1:A:461:ASN:ND2	5:A:799:HOH:O	2.37	0.57
1:A:106:GLU:OE1	5:A:607:HOH:O	2.18	0.55
1:C:106:GLU:OE1	5:C:648:HOH:O	2.19	0.53
1:C:411:ASN:ND2	5:C:740:HOH:O	2.41	0.52
5:A:625:HOH:O	2:C:511:NAG:H81	2.08	0.51
1:C:426:MET:SD	1:C:430:THR:OG1	2.68	0.50
1:A:224:VAL:HG11	1:A:244:SER:HB2	1.95	0.49
1:A:426:MET:SD	1:A:430:THR:OG1	2.71	0.48
1:A:474:ASN:HA	4:A:513:OLM:H6	1.95	0.48
1:C:121:LYS:HE2	1:C:123:THR:HG21	1.94	0.48
1:C:101:VAL:HG21	1:C:480:ARG:HG3	1.95	0.48
1:C:44:VAL:HG12	5:C:741:HOH:O	2.14	0.47
1:C:327:ARG:NH2	1:C:422:GLN:OE1	2.36	0.46
4:C:513:OLM:H24	4:C:513:OLM:H3	1.72	0.46
1:C:198:GLY:N	5:C:734:HOH:O	2.28	0.45
1:A:361:PHE:O	1:A:393:ASN:ND2	2.44	0.45
1:C:288:LEU:HB3	5:C:724:HOH:O	2.17	0.44
1:A:273:ARG:HB2	1:A:285:ILE:HB	1.99	0.43
1:A:116:LEU:HD11	1:A:210:PHE:CE2	2.54	0.42
1:A:432:GLN:NE2	5:A:749:HOH:O	2.36	0.42
1:C:224:VAL:HG11	1:C:244:SER:HB2	2.02	0.41
1:A:420:ILE:HG21	1:A:438:PRO:HG3	2.03	0.41
1:C:268:GLU:HB2	5:C:729:HOH:O	2.20	0.41
1:A:207:LYS:HD3	1:A:439:ILE:HG23	2.03	0.41
1:C:411:ASN:N	1:C:411:ASN:HD22	2.19	0.41
2:A:511:NAG:H2	2:A:511:NAG:H83	1.92	0.41
1:A:361:PHE:HB3	1:A:391:PHE:HB3	2.02	0.40
1:C:257:THR:O	1:C:259:LEU:N	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/353 (94%)	323 (97%)	10 (3%)	0	100	100
1	C	333/353 (94%)	322 (97%)	11 (3%)	0	100	100
All	All	666/706 (94%)	645 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/311 (97%)	303 (100%)	0	100	100
1	C	303/311 (97%)	300 (99%)	3 (1%)	82	84
All	All	606/622 (97%)	603 (100%)	3 (0%)	92	93

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	44	VAL
1	C	411	ASN
1	C	465	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no

such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

26 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	501	1	14,14,15	0.52	0	15,19,21	0.62	0
2	NAG	A	502	1	14,14,15	0.50	0	15,19,21	0.61	0
2	NAG	A	503	1	14,14,15	0.48	0	15,19,21	0.80	0
2	NAG	A	504	1	14,14,15	0.48	0	15,19,21	0.67	0
2	NAG	A	505	1	14,14,15	0.48	0	15,19,21	0.81	0
2	NAG	A	506	1	14,14,15	0.47	0	15,19,21	0.94	0
2	NAG	A	507	1	14,14,15	0.49	0	15,19,21	0.72	0
2	NAG	A	508	1	14,14,15	0.53	0	15,19,21	0.67	0
2	NAG	A	509	1	14,14,15	0.50	0	15,19,21	0.66	0
2	NAG	A	510	1	14,14,15	0.52	0	15,19,21	0.65	0
2	NAG	A	511	1	14,14,15	0.47	0	15,19,21	0.72	0
3	EPE	A	512	-	14,15,15	0.34	0	18,20,20	1.84	7 (38%)
4	OLM	A	513	-	24,26,26	2.31	7 (29%)	37,40,40	2.67	8 (21%)
2	NAG	C	501	1	14,14,15	0.50	0	15,19,21	0.75	0
2	NAG	C	502	1	14,14,15	0.50	0	15,19,21	0.75	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	C	503	1	14,14,15	0.43	0	15,19,21	1.00	1 (6%)
2	NAG	C	504	1	14,14,15	0.51	0	15,19,21	0.68	0
2	NAG	C	505	1	14,14,15	0.49	0	15,19,21	1.11	1 (6%)
2	NAG	C	506	1	14,14,15	0.49	0	15,19,21	0.68	0
2	NAG	C	507	1	14,14,15	0.49	0	15,19,21	0.70	0
2	NAG	C	508	1	14,14,15	0.51	0	15,19,21	0.71	0
2	NAG	C	509	1	14,14,15	0.48	0	15,19,21	0.82	0
2	NAG	C	510	1	14,14,15	0.47	0	15,19,21	0.75	0
2	NAG	C	511	1	14,14,15	0.47	0	15,19,21	0.79	0
3	EPE	C	512	-	14,15,15	0.35	0	18,20,20	1.86	6 (33%)
4	OLM	C	513	-	24,26,26	2.26	7 (29%)	37,40,40	3.08	7 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	501	1	-	0/6/23/26	0/1/1/1
2	NAG	A	502	1	-	0/6/23/26	0/1/1/1
2	NAG	A	503	1	-	0/6/23/26	0/1/1/1
2	NAG	A	504	1	-	0/6/23/26	0/1/1/1
2	NAG	A	505	1	-	0/6/23/26	0/1/1/1
2	NAG	A	506	1	-	0/6/23/26	0/1/1/1
2	NAG	A	507	1	-	0/6/23/26	0/1/1/1
2	NAG	A	508	1	-	0/6/23/26	0/1/1/1
2	NAG	A	509	1	-	0/6/23/26	0/1/1/1
2	NAG	A	510	1	-	0/6/23/26	0/1/1/1
2	NAG	A	511	1	-	0/6/23/26	0/1/1/1
3	EPE	A	512	-	-	0/9/19/19	0/1/1/1
4	OLM	A	513	-	-	0/12/34/34	0/2/2/2
2	NAG	C	501	1	-	0/6/23/26	0/1/1/1
2	NAG	C	502	1	-	0/6/23/26	0/1/1/1
2	NAG	C	503	1	-	0/6/23/26	0/1/1/1
2	NAG	C	504	1	-	0/6/23/26	0/1/1/1
2	NAG	C	505	1	-	0/6/23/26	0/1/1/1
2	NAG	C	506	1	-	0/6/23/26	0/1/1/1
2	NAG	C	507	1	-	0/6/23/26	0/1/1/1
2	NAG	C	508	1	-	0/6/23/26	0/1/1/1
2	NAG	C	509	1	-	0/6/23/26	0/1/1/1
2	NAG	C	510	1	-	0/6/23/26	0/1/1/1
2	NAG	C	511	1	-	0/6/23/26	0/1/1/1

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EPE	C	512	-	-	0/9/19/19	0/1/1/1
4	OLM	C	513	-	-	0/12/34/34	0/2/2/2

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	513	OLM	C14-C13	-7.12	1.44	1.52
4	C	513	OLM	C14-C13	-6.84	1.44	1.52
4	C	513	OLM	C18-C13	-3.33	1.48	1.52
4	A	513	OLM	C18-C13	-3.33	1.48	1.52
4	A	513	OLM	C08-C10	-2.56	1.49	1.53
4	C	513	OLM	C08-C10	-2.52	1.49	1.53
4	A	513	OLM	C18-C17	-2.35	1.48	1.53
4	C	513	OLM	C18-C17	-2.33	1.49	1.53
4	C	513	OLM	C13-N11	2.16	1.50	1.46
4	A	513	OLM	C13-N11	2.25	1.51	1.46
4	A	513	OLM	C08-N07	3.49	1.43	1.35
4	C	513	OLM	C08-N07	3.50	1.43	1.35
4	C	513	OLM	C10-N11	4.30	1.43	1.34
4	A	513	OLM	C10-N11	4.45	1.43	1.34

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	512	EPE	C5-C6-N1	-2.61	105.95	110.63
3	C	512	EPE	C5-C6-N1	-2.60	105.97	110.63
3	A	512	EPE	O2S-S-C10	2.08	108.68	106.91
2	C	503	NAG	C1-O5-C5	2.09	114.90	112.25
3	C	512	EPE	C2-C3-N4	2.16	114.50	110.63
3	C	512	EPE	C7-N4-C5	2.19	116.89	111.27
3	A	512	EPE	C7-N4-C5	2.21	116.93	111.27
3	A	512	EPE	C2-C3-N4	2.22	114.59	110.63
3	A	512	EPE	C7-N4-C3	2.24	117.01	111.27
3	C	512	EPE	C7-N4-C3	2.34	117.27	111.27
4	A	513	OLM	C17-C18-C13	2.43	116.01	113.85
4	C	513	OLM	C17-C18-C13	2.59	116.16	113.85
4	A	513	OLM	C18-C13-N11	2.69	113.54	110.23
3	C	512	EPE	O2S-S-C10	2.78	109.28	106.91
4	A	513	OLM	C15-C14-C13	2.86	116.40	113.85
3	A	512	EPE	O1S-S-C10	3.00	109.47	106.91
2	C	505	NAG	C1-O5-C5	3.53	116.72	112.25
4	C	513	OLM	C10-C08-N07	3.85	118.58	112.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	513	OLM	C10-C08-N07	4.11	119.01	112.30
3	A	512	EPE	C5-N4-C3	4.55	118.75	108.90
4	A	513	OLM	C14-C13-C18	4.64	112.86	108.72
3	C	512	EPE	C5-N4-C3	4.70	119.08	108.90
4	C	513	OLM	C15-C14-C13	5.28	118.55	113.85
4	C	513	OLM	C18-C17-N16	6.07	114.31	108.37
4	A	513	OLM	C18-C17-N16	6.58	114.82	108.37
4	A	513	OLM	C08-C10-N11	7.13	119.56	113.34
4	C	513	OLM	C14-C13-C18	7.42	115.34	108.72
4	C	513	OLM	C08-C10-N11	8.37	120.64	113.34
4	A	513	OLM	C14-C15-N16	8.81	117.00	108.37
4	C	513	OLM	C14-C15-N16	10.17	118.33	108.37

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	511	NAG	1	0
4	A	513	OLM	1	0
2	C	511	NAG	1	0
4	C	513	OLM	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	339/353 (96%)	0.28	10 (2%) 55 58	17, 30, 63, 95	0
1	C	339/353 (96%)	0.89	43 (12%) 5 6	23, 45, 89, 123	0
All	All	678/706 (96%)	0.59	53 (7%) 16 19	17, 37, 79, 123	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	460	ALA	7.9
1	C	461	ASN	6.6
1	C	464	SER	6.0
1	A	492	GLU	5.9
1	A	124	GLY	5.8
1	C	492	GLU	5.7
1	C	472	GLY	5.1
1	C	396	ILE	5.0
1	C	465	ASN	4.9
1	C	44	VAL	4.8
1	C	462	ASN	4.7
1	C	358	THR	4.5
1	C	463	THR	4.4
1	C	278	THR	4.2
1	A	472	GLY	4.1
1	C	89	VAL	4.0
1	A	473	GLY	3.8
1	C	354	ASN	3.8
1	C	491	ILE	3.8
1	C	355	ASN	3.7
1	C	466	GLU	3.5
1	A	301	ASN	3.4
1	C	473	GLY	3.3
1	C	45	TRP	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	324	GLY	3.0
1	C	233	PHE	3.0
1	C	394	THR	2.9
1	C	301	ASN	2.9
1	C	359	ILE	2.9
1	C	277	LEU	2.8
1	C	458	GLY	2.8
1	C	353	PHE	2.7
1	C	224	VAL	2.7
1	C	82	GLN	2.6
1	C	430	THR	2.6
1	C	350	LYS	2.5
1	C	352	HIS	2.5
1	C	390	LEU	2.5
1	C	236	THR	2.4
1	A	324	GLY	2.4
1	A	44	VAL	2.4
1	C	392	ASN	2.3
1	C	271	ILE	2.3
1	C	365	SER	2.2
1	C	87	GLU	2.2
1	C	235	GLY	2.2
1	C	61	HIS	2.2
1	A	45	TRP	2.2
1	C	116	LEU	2.1
1	A	411	ASN	2.1
1	C	88	ASN	2.1
1	C	459	GLY	2.1
1	A	116	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	506	14/15	0.90	0.22	5.39	32,56,68,72	0
4	OLM	C	513	25/25	0.78	0.32	4.84	49,105,133,133	0
2	NAG	C	511	14/15	0.81	0.20	4.49	43,64,74,77	0
4	OLM	A	513	25/25	0.80	0.31	3.92	46,83,125,128	0
2	NAG	A	501	14/15	0.90	0.15	1.95	35,50,58,65	0
2	NAG	C	505	14/15	0.78	0.18	1.85	42,56,64,65	0
2	NAG	C	506	14/15	0.88	0.14	1.37	39,52,64,72	0
2	NAG	A	507	14/15	0.85	0.15	0.82	44,53,65,69	0
2	NAG	A	508	14/15	0.91	0.13	0.57	25,42,67,77	0
3	EPE	A	512	15/15	0.98	0.13	0.40	20,25,33,34	0
3	EPE	C	512	15/15	0.97	0.15	0.36	22,31,44,45	0
2	NAG	C	501	14/15	0.89	0.20	0.34	54,64,76,77	0
2	NAG	A	503	14/15	0.95	0.12	-0.11	16,24,37,38	0
2	NAG	A	505	14/15	0.93	0.10	-1.40	29,42,53,65	0
2	NAG	C	503	14/15	0.97	0.11	-1.52	16,23,27,29	0
2	NAG	C	508	14/15	0.64	0.51	-	100,109,117,118	0
2	NAG	C	507	14/15	0.72	0.24	-	57,72,79,81	0
2	NAG	C	509	14/15	0.89	0.15	-	47,58,65,66	0
2	NAG	C	502	14/15	0.56	0.36	-	83,101,107,107	0
2	NAG	A	510	14/15	0.67	0.35	-	72,86,94,94	0
2	NAG	A	502	14/15	0.46	0.42	-	79,88,96,99	0
2	NAG	A	511	14/15	0.55	0.32	-	79,88,96,98	0
2	NAG	C	504	14/15	0.68	0.29	-	66,84,88,89	0
2	NAG	A	504	14/15	0.87	0.15	-	32,41,49,54	0
2	NAG	A	509	14/15	0.94	0.10	-	31,44,53,64	0
2	NAG	C	510	14/15	0.74	0.34	-	71,80,91,98	0

## 6.5 Other polymers

There are no such residues in this entry.