



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:37 AM GMT

PDB ID : 2DPE
Title : Crystal structure of a secretory 40KDA glycoprotein from sheep at 2.0A resolution
Authors : Srivastava, D.B.; Ethayathulla, A.S.; Kumar, J.; Singh, N.; Sharma, S.; Das, U.; Srinivasan, A.; Singh, T.P.
Deposited on : 2006-05-11
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1799 (2.08-2.04)
Clashscore	102246	1910 (2.08-2.04)
Ramachandran outliers	100387	1893 (2.08-2.04)
Sidechain outliers	100360	1893 (2.08-2.04)
RSRZ outliers	91569	1802 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	<div> <div>4%</div> <div>80%</div> <div>17%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	363	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MAN	A	366	X	-	-	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3168 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chitinase-3-like protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	361	Total	C	N	O	S	0	0	0
			2869	1832	499	529	9			

- Molecule 2 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	5	Total	C	N	O	0	0
			63	34	2	27		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	236	Total	O	0	0
			236	236		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chitinase-3-like protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.71Å 66.44Å 107.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.94 – 2.07 56.52 – 2.07	Depositor EDS
% Data completeness (in resolution range)	99.7 (19.94-2.07) 99.8 (56.52-2.07)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 2.07Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.185 , 0.203 0.183 , 0.209	Depositor DCC
R_{free} test set	551 reflections (2.01%)	DCC
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 64.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 28016 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3168	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	1/2945 (0.0%)	0.77	5/3995 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	A	1	0
All	All	1	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	191	TRP	N-CA	7.04	1.60	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	ALA	O-C-N	-9.54	107.43	122.70
1	A	320	LEU	CA-CB-CG	8.51	134.88	115.30
1	A	189	GLY	N-CA-C	-8.50	91.84	113.10
1	A	190	ALA	CA-C-N	6.56	131.64	117.20
1	A	191	TRP	N-CA-C	5.75	126.52	111.00

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	366	MAN	C1

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	190	ALA	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2869	0	2794	57	0
2	A	63	0	53	1	0
3	A	236	0	0	20	0
All	All	3168	0	2847	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:166:ARG:HA	3:A:502:HOH:O	1.50	1.11
1:A:204:GLY:HA2	1:A:292:ASP:HB3	1.48	0.95
1:A:57:ASP:O	1:A:61:THR:HG23	1.71	0.89
1:A:188:HIS:CD2	1:A:189:GLY:H	1.92	0.87
1:A:96:ARG:HB2	3:A:486:HOH:O	1.78	0.83
1:A:347:LEU:HD12	3:A:598:HOH:O	1.84	0.76
1:A:78:TRP:HH2	3:A:452:HOH:O	1.69	0.75
1:A:28:PHE:HE1	3:A:576:HOH:O	1.73	0.72
1:A:204:GLY:CA	1:A:292:ASP:HB3	2.21	0.69
1:A:281:PHE:O	1:A:300:LYS:HE3	1.93	0.69
1:A:239:THR:HG22	1:A:335:LEU:HB2	1.77	0.67
1:A:95:ARG:NH2	1:A:131:THR:HG21	2.10	0.66
1:A:239:THR:HG21	1:A:332:ALA:O	1.97	0.65
1:A:188:HIS:HD2	1:A:189:GLY:H	1.41	0.64
1:A:190:ALA:O	1:A:191:TRP:HB2	1.98	0.64
1:A:362:VAL:HG12	1:A:362:VAL:OXT	1.98	0.63
1:A:313:VAL:HG13	1:A:355:VAL:CG2	2.29	0.62
1:A:107:ARG:HD3	1:A:143:GLU:OE2	2.00	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:317:ALA:HA	1:A:320:LEU:HD13	1.81	0.61
1:A:95:ARG:CZ	1:A:131:THR:HG21	2.30	0.60
1:A:15:GLU:OE1	1:A:268:LYS:NZ	2.35	0.60
1:A:148:THR:HB	3:A:491:HOH:O	2.02	0.59
1:A:210:SER:O	1:A:213:PHE:N	2.36	0.58
1:A:210:SER:C	1:A:213:PHE:H	2.06	0.58
1:A:287:THR:HB	3:A:560:HOH:O	2.03	0.58
1:A:96:ARG:CB	3:A:486:HOH:O	2.45	0.58
1:A:188:HIS:CD2	1:A:189:GLY:N	2.68	0.58
1:A:262:VAL:H	1:A:303:GLN:HE22	1.52	0.58
1:A:96:ARG:CA	3:A:486:HOH:O	2.53	0.56
1:A:148:THR:HG23	3:A:499:HOH:O	2.06	0.56
1:A:313:VAL:HG13	1:A:355:VAL:HG22	1.86	0.56
1:A:166:ARG:CA	3:A:502:HOH:O	2.25	0.55
1:A:166:ARG:CB	3:A:502:HOH:O	2.55	0.54
1:A:239:THR:CG2	1:A:335:LEU:HB2	2.37	0.53
1:A:95:ARG:CZ	1:A:131:THR:CG2	2.88	0.52
1:A:314:LYS:HE2	3:A:596:HOH:O	2.10	0.51
1:A:126:LYS:NZ	3:A:568:HOH:O	2.45	0.50
1:A:198:HIS:CD2	1:A:198:HIS:H	2.31	0.49
1:A:291:ARG:HG3	3:A:530:HOH:O	2.12	0.49
1:A:289:ARG:NH2	1:A:309:ASP:OD2	2.44	0.48
1:A:205:ASN:HD22	1:A:205:ASN:C	2.16	0.48
1:A:78:TRP:CB	3:A:580:HOH:O	2.61	0.48
1:A:210:SER:C	1:A:213:PHE:N	2.66	0.48
1:A:362:VAL:CG1	1:A:362:VAL:OXT	2.62	0.48
1:A:78:TRP:CH2	3:A:452:HOH:O	2.55	0.48
1:A:239:THR:HG23	1:A:239:THR:O	2.15	0.46
1:A:95:ARG:NH2	1:A:131:THR:CG2	2.77	0.46
1:A:28:PHE:CE1	3:A:576:HOH:O	2.54	0.45
1:A:102:VAL:HB	1:A:103:PRO:HD3	1.97	0.45
1:A:188:HIS:HD2	1:A:189:GLY:N	2.10	0.45
1:A:262:VAL:H	1:A:303:GLN:NE2	2.14	0.44
1:A:348:THR:HG21	3:A:574:HOH:O	2.18	0.44
1:A:237:ILE:CG2	1:A:313:VAL:HG22	2.49	0.43
1:A:272:ILE:HG22	1:A:273:LEU:N	2.35	0.41
2:A:365:BMA:H61	2:A:366:MAN:H2	1.40	0.41
1:A:204:GLY:HA2	1:A:292:ASP:CB	2.35	0.41
1:A:321:LYS:NZ	3:A:602:HOH:O	2.54	0.41
1:A:344:GLY:O	1:A:345:GLN:C	2.60	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	359/361 (99%)	344 (96%)	9 (2%)	6 (2%)	11 3

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	207	ASP
1	A	190	ALA
1	A	205	ASN
1	A	346	ASN
1	A	193	GLN
1	A	212	ARG

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	302/302 (100%)	272 (90%)	30 (10%)	10 4

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LEU
1	A	69	LEU
1	A	83	GLU
1	A	85	PHE
1	A	86	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	93	GLN
1	A	106	LEU
1	A	107	ARG
1	A	129	LEU
1	A	149	GLU
1	A	175	ARG
1	A	193	GLN
1	A	195	VAL
1	A	205	ASN
1	A	210	SER
1	A	212	ARG
1	A	239	THR
1	A	252	ASP
1	A	258	SER
1	A	265	ARG
1	A	282	LEU
1	A	299	THR
1	A	311	GLU
1	A	313	VAL
1	A	320	LEU
1	A	345	GLN
1	A	347	LEU
1	A	348	THR
1	A	351	LEU
1	A	359	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	ASN
1	A	109	HIS
1	A	188	HIS
1	A	193	GLN
1	A	198	HIS
1	A	205	ASN
1	A	294	GLN
1	A	303	GLN
1	A	315	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

5 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	363	1,2	14,14,15	0.72	0	15,19,21	0.97	1 (6%)
2	NAG	A	364	2	14,14,15	0.57	0	15,19,21	1.79	3 (20%)
2	BMA	A	365	2	11,11,12	2.21	3 (27%)	14,15,17	3.50	6 (42%)
2	MAN	A	366	2	12,12,12	0.73	0	17,17,17	1.99	3 (17%)
2	MAN	A	367	2	12,12,12	0.97	1 (8%)	17,17,17	1.97	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	363	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	364	2	-	0/6/23/26	0/1/1/1
2	BMA	A	365	2	-	0/2/19/22	0/1/1/1
2	MAN	A	366	2	1/1/5/5	0/2/22/22	0/1/1/1
2	MAN	A	367	2	-	0/2/22/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	365	BMA	O3-C3	-5.34	1.30	1.43
2	A	365	BMA	C4-C5	-4.06	1.44	1.53
2	A	367	MAN	O5-C5	-2.31	1.38	1.44
2	A	365	BMA	C4-C3	-2.05	1.47	1.52

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	365	BMA	O3-C3-C4	-6.21	96.35	110.34
2	A	365	BMA	O3-C3-C2	-6.17	98.86	110.00
2	A	365	BMA	C3-C4-C5	-5.31	100.94	110.20
2	A	365	BMA	C1-C2-C3	-4.44	104.28	109.54
2	A	366	MAN	O1-C1-O5	-3.11	101.74	110.25
2	A	366	MAN	C4-C3-C2	-2.89	105.40	110.79
2	A	363	NAG	C2-N2-C7	-2.80	119.44	123.04
2	A	367	MAN	O1-C1-O5	-2.76	102.70	110.25
2	A	364	NAG	C3-C2-N2	-2.13	105.45	110.56
2	A	367	MAN	C3-C4-C5	-2.05	106.61	110.20
2	A	367	MAN	O2-C2-C1	2.04	114.31	109.82
2	A	367	MAN	C1-C2-C3	2.57	114.25	110.43
2	A	365	BMA	O5-C5-C6	3.21	114.29	107.35
2	A	364	NAG	C4-C3-C2	3.52	116.70	111.23
2	A	364	NAG	C3-C4-C5	4.85	118.65	110.20
2	A	365	BMA	C2-C3-C4	5.20	119.86	111.04
2	A	367	MAN	O5-C1-C2	5.69	118.88	109.80
2	A	366	MAN	O5-C1-C2	5.70	118.89	109.80

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	366	MAN	C1

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	365	BMA	1	0
2	A	366	MAN	1	0

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	361/361 (100%)	0.01	15 (4%)	40 44	22, 35, 64, 99	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	208	ALA	8.4
1	A	191	TRP	8.1
1	A	189	GLY	7.7
1	A	207	ASP	7.1
1	A	209	SER	6.2
1	A	362	VAL	5.5
1	A	190	ALA	5.4
1	A	210	SER	3.9
1	A	204	GLY	3.4
1	A	192	ARG	3.0
1	A	212	ARG	2.9
1	A	253	VAL	2.3
1	A	265	ARG	2.2
1	A	78	TRP	2.1
1	A	131	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAG	A	363	14/15	0.87	0.36	8.85	35,37,39,40	14
2	MAN	A	367	12/12	0.55	0.45	-	40,41,44,45	11
2	MAN	A	366	12/12	0.52	0.34	-	40,50,62,74	11
2	NAG	A	364	14/15	0.67	0.27	-	37,40,41,41	14
2	BMA	A	365	11/12	0.65	0.30	-	36,39,41,42	11

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.