



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:53 PM GMT

PDB ID : 4DRS
Title : Crystal structure of Cryptosporidium parvum pyruvate kinase
Authors : Cook, W.J.; Chattopadhyay, D.
Deposited on : 2012-02-17
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	526	
1	B	526	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	604	-	-	X	-
4	ACT	B	605	-	-	-	X
4	ACT	B	607	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7505 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	485	Total	C	N	O	S	0	0	0
			3644	2279	629	704	32			
1	B	485	Total	C	N	O	S	0	0	0
			3644	2279	629	704	32			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	74	Total O 74 74	0	0
5	B	75	Total O 75 75	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pyruvate kinase



• Molecule 1: Pyruvate kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	129.90 Å 136.94 Å 77.23 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.71 – 2.50 49.71 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.8 (49.71-2.50) 97.8 (49.71-2.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.39 (at 2.51 Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.213 , 0.249 0.213 , 0.249	Depositor DCC
R_{free} test set	2392 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	47.3	Xtriage
Anisotropy	0.133	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 47350 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7505	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/3692	0.48	0/4991
1	B	0.35	0/3692	0.48	0/4991
All	All	0.35	0/7384	0.48	0/9982

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3644	0	3733	36	0
1	B	3644	0	3733	41	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
3	A	12	0	16	1	0
3	B	12	0	16	5	0
4	A	12	0	9	1	0
4	B	12	0	9	0	0
5	A	74	0	0	0	0
5	B	75	0	0	0	0
All	All	7505	0	7516	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:229:ASN:HD21	3:B:604:GOL:H31	1.52	0.73
1:B:339:ARG:HG2	1:B:340:PRO:HD2	1.71	0.72
1:B:229:ASN:ND2	3:B:604:GOL:H31	2.06	0.69
1:B:229:ASN:HD21	3:B:604:GOL:C3	2.05	0.69
1:A:175:THR:HB	1:A:187:CYS:HB3	1.79	0.65
1:A:168:ILE:HD12	1:A:173:LEU:HD23	1.78	0.65
1:B:331:LEU:HD12	1:B:361:LEU:HD21	1.78	0.64
1:B:168:ILE:HG12	1:B:201:MET:HG3	1.80	0.64
1:B:335:ILE:HG23	1:B:368:GLY:HA2	1.81	0.62
1:A:339:ARG:HG2	1:A:340:PRO:HD2	1.81	0.62
1:A:32:ILE:HG12	1:B:305:LYS:HD3	1.81	0.62
1:B:175:THR:HB	1:B:187:CYS:HB3	1.81	0.61
1:A:331:LEU:HB2	1:A:364:GLU:HG2	1.82	0.60
1:A:431:ILE:HB	1:A:507:HIS:HB3	1.84	0.60
1:A:193:VAL:HG12	1:A:194:THR:N	2.16	0.59
1:A:335:ILE:HG23	1:A:368:GLY:HA2	1.84	0.59
1:B:168:ILE:HB	1:B:173:LEU:HB3	1.87	0.57
1:B:327:ALA:HB2	1:B:360:MET:HB3	1.86	0.57
1:A:298:GLY:HA2	1:A:306:ILE:HD11	1.87	0.57
1:A:384:GLN:HG3	1:B:308:VAL:HG21	1.88	0.56
1:A:120:LEU:HD22	1:A:126:ILE:HD12	1.87	0.56
1:A:331:LEU:HD12	1:A:361:LEU:HD21	1.88	0.55
1:A:168:ILE:HG12	1:A:201:MET:HG3	1.89	0.55
1:A:168:ILE:HB	1:A:173:LEU:HB3	1.91	0.53
1:B:361:LEU:HG	1:B:378:MET:CE	2.40	0.52
1:A:305:LYS:HD3	1:B:32:ILE:HG12	1.91	0.52
1:B:168:ILE:HD12	1:B:173:LEU:HD23	1.91	0.52
1:B:193:VAL:HG12	1:B:194:THR:N	2.25	0.52
1:A:361:LEU:HG	1:A:378:MET:CE	2.40	0.52
1:A:80:GLY:H	4:A:606:ACT:H3	1.75	0.52
1:A:64:GLY:O	1:A:68:LYS:HG2	2.09	0.52
1:A:193:VAL:HG12	1:A:194:THR:H	1.74	0.51
1:B:461:ARG:NH2	3:B:604:GOL:H11	2.26	0.51
1:A:342:ARG:CZ	1:B:294:ARG:HG3	2.40	0.50
1:B:120:LEU:HD22	1:B:126:ILE:HD12	1.93	0.50
1:A:167:LEU:HB2	1:A:202:ASN:HB2	1.94	0.49
1:B:331:LEU:HB2	1:B:364:GLU:HG2	1.95	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:327:ALA:CB	1:B:360:MET:HB3	2.43	0.48
1:B:452:ALA:HB3	1:B:471:THR:HG22	1.95	0.48
1:A:294:ARG:HG3	1:B:342:ARG:CZ	2.44	0.48
1:A:156:LEU:N	1:A:157:PRO:HD2	2.29	0.47
1:A:49:ILE:HG12	1:A:72:VAL:HB	1.95	0.47
1:A:118:GLY:HA3	1:A:150:SER:HB3	1.96	0.47
1:A:29:MET:HA	1:A:32:ILE:HD12	1.97	0.46
1:B:329:GLN:HG3	1:B:362:SER:HB2	1.97	0.46
1:A:214:GLY:O	1:A:218:ARG:HG3	2.16	0.46
1:A:74:ARG:HH22	3:A:603:GOL:H2	1.81	0.46
1:A:327:ALA:HB2	1:A:360:MET:HB3	1.97	0.45
1:A:181:GLY:HA3	1:A:184:PHE:CE1	2.51	0.45
1:B:431:ILE:HB	1:B:507:HIS:HB3	1.98	0.45
1:A:431:ILE:HG23	1:A:477:ILE:HG13	2.00	0.44
1:A:193:VAL:CG1	1:A:194:THR:N	2.80	0.44
1:B:156:LEU:N	1:B:157:PRO:HD2	2.33	0.43
1:B:361:LEU:HG	1:B:378:MET:HE2	2.00	0.43
1:B:365:THR:HA	1:B:371:PRO:HB3	2.01	0.43
1:B:431:ILE:HG23	1:B:477:ILE:HG13	2.00	0.43
1:B:302:PRO:HA	1:B:303:PRO:HD3	1.87	0.43
1:B:470:LYS:HE2	1:B:494:GLU:OE1	2.19	0.43
1:A:193:VAL:CG1	1:A:194:THR:H	2.32	0.42
1:B:29:MET:HA	1:B:32:ILE:HD12	2.02	0.42
1:B:118:GLY:HA3	1:B:150:SER:HB3	2.02	0.41
1:A:272:GLU:HB3	1:A:295:GLY:HA3	2.03	0.41
1:B:419:SER:O	1:B:423:VAL:HG22	2.20	0.41
1:B:339:ARG:HG2	1:B:340:PRO:CD	2.46	0.41
1:B:270:LYS:HZ1	3:B:603:GOL:H11	1.84	0.41
1:B:181:GLY:HA3	1:B:184:PHE:CE1	2.55	0.41
1:B:218:ARG:HH11	1:B:218:ARG:HB3	1.84	0.41
1:B:218:ARG:NH1	1:B:218:ARG:HB3	2.35	0.41
1:A:333:SER:O	1:A:337:SER:HB3	2.21	0.41
1:B:272:GLU:HB3	1:B:295:GLY:HA3	2.01	0.41
1:A:302:PRO:HA	1:A:303:PRO:HD3	1.83	0.41
1:B:272:GLU:HG2	1:B:293:ALA:HB3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	479/526 (91%)	463 (97%)	14 (3%)	2 (0%)	39	61
1	B	479/526 (91%)	460 (96%)	17 (4%)	2 (0%)	39	61
All	All	958/1052 (91%)	923 (96%)	31 (3%)	4 (0%)	39	61

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	121	GLU
1	A	123	GLY
1	B	121	GLU
1	B	123	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	414/449 (92%)	413 (100%)	1 (0%)	95	99
1	B	414/449 (92%)	413 (100%)	1 (0%)	95	99
All	All	828/898 (92%)	826 (100%)	2 (0%)	95	99

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	419	SER
1	B	215	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	518	ASN
1	B	229	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	601	-	4,4,4	0.25	0	6,6,6	0.17	0
2	SO4	A	602	-	4,4,4	0.40	0	6,6,6	0.22	0
3	GOL	A	603	-	5,5,5	0.28	0	5,5,5	0.29	0
3	GOL	A	604	-	5,5,5	0.26	0	5,5,5	0.36	0
4	ACT	A	605	-	1,3,3	1.44	0	0,3,3	0.00	-
4	ACT	A	606	-	1,3,3	1.20	0	0,3,3	0.00	-
4	ACT	A	607	-	1,3,3	1.25	0	0,3,3	0.00	-
2	SO4	B	601	-	4,4,4	0.32	0	6,6,6	0.09	0
2	SO4	B	602	-	4,4,4	0.40	0	6,6,6	0.18	0
3	GOL	B	603	-	5,5,5	0.29	0	5,5,5	0.34	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	604	-	5,5,5	0.23	0	5,5,5	0.33	0
4	ACT	B	605	-	1,3,3	1.24	0	0,3,3	0.00	-
4	ACT	B	606	-	1,3,3	1.24	0	0,3,3	0.00	-
4	ACT	B	607	-	1,3,3	1.48	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	601	-	-	0/0/0/0	0/0/0/0
2	SO4	A	602	-	-	0/0/0/0	0/0/0/0
3	GOL	A	603	-	-	0/4/4/4	0/0/0/0
3	GOL	A	604	-	-	0/4/4/4	0/0/0/0
4	ACT	A	605	-	-	0/0/0/0	0/0/0/0
4	ACT	A	606	-	-	0/0/0/0	0/0/0/0
4	ACT	A	607	-	-	0/0/0/0	0/0/0/0
2	SO4	B	601	-	-	0/0/0/0	0/0/0/0
2	SO4	B	602	-	-	0/0/0/0	0/0/0/0
3	GOL	B	603	-	-	0/4/4/4	0/0/0/0
3	GOL	B	604	-	-	0/4/4/4	0/0/0/0
4	ACT	B	605	-	-	0/0/0/0	0/0/0/0
4	ACT	B	606	-	-	0/0/0/0	0/0/0/0
4	ACT	B	607	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	603	GOL	1	0
4	A	606	ACT	1	0
3	B	603	GOL	1	0
3	B	604	GOL	4	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	485/526 (92%)	0.45	37 (7%)	17 18	32, 51, 111, 142	0
1	B	485/526 (92%)	0.64	63 (12%)	5 4	33, 50, 139, 176	0
All	All	970/1052 (92%)	0.54	100 (10%)	9 9	32, 51, 125, 176	0

All (100) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	119	MET	11.0
1	B	128	LEU	8.5
1	A	126	ILE	8.3
1	B	195	ILE	7.1
1	B	194	THR	7.0
1	B	134	LEU	6.9
1	B	125	PRO	6.9
1	A	173	LEU	6.6
1	B	196	GLY	6.1
1	A	195	ILE	6.0
1	A	119	MET	5.5
1	B	130	ALA	5.3
1	B	127	GLU	5.2
1	A	128	LEU	5.2
1	A	127	GLU	4.9
1	A	121	GLU	4.8
1	B	120	LEU	4.8
1	B	121	GLU	4.8
1	B	126	ILE	4.8
1	A	147	GLU	4.7
1	A	143	LEU	4.6
1	A	149	ILE	4.6
1	A	196	GLY	4.5
1	A	365	THR	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	133	THR	4.4
1	B	208	VAL	4.4
1	B	140	TYR	4.3
1	A	134	LEU	4.3
1	A	189	VAL	4.3
1	B	143	LEU	4.2
1	B	191	ASN	4.2
1	B	147	GLU	4.1
1	B	146	SER	4.0
1	B	118	GLY	4.0
1	B	172	SER	3.9
1	A	118	GLY	3.9
1	B	151	CYS	3.8
1	B	122	GLY	3.8
1	B	173	LEU	3.8
1	B	150	SER	3.8
1	B	198	ARG	3.8
1	A	170	ASP	3.7
1	A	198	ARG	3.7
1	B	149	ILE	3.7
1	B	129	LYS	3.7
1	A	193	VAL	3.6
1	A	175	THR	3.6
1	B	169	ALA	3.6
1	B	117	THR	3.5
1	A	330	MET	3.5
1	B	139	ASP	3.5
1	B	42	THR	3.4
1	A	146	SER	3.3
1	B	193	VAL	3.2
1	B	339	ARG	3.1
1	A	194	THR	3.1
1	A	327	ALA	3.1
1	B	369	ALA	3.1
1	A	334	MET	3.0
1	B	124	LYS	3.0
1	B	190	LEU	3.0
1	B	185	ILE	3.0
1	B	327	ALA	3.0
1	A	336	LYS	3.0
1	B	179	GLU	2.9
1	B	192	SER	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	201	MET	2.9
1	B	142	MET	2.9
1	A	120	LEU	2.9
1	A	125	PRO	2.8
1	A	148	CYS	2.8
1	B	170	ASP	2.7
1	B	123	GLY	2.6
1	B	200	ASN	2.5
1	B	197	GLU	2.5
1	B	212	ILE	2.5
1	B	113	PRO	2.4
1	A	141	SER	2.4
1	A	122	GLY	2.4
1	A	123	GLY	2.4
1	B	365	THR	2.4
1	B	187	CYS	2.4
1	B	178	LEU	2.4
1	A	124	LYS	2.4
1	B	57	ASN	2.4
1	B	29	MET	2.3
1	A	190	LEU	2.3
1	B	370	PHE	2.3
1	A	363	GLY	2.2
1	A	133	THR	2.2
1	B	207	LYS	2.1
1	B	334	MET	2.1
1	B	199	LYS	2.1
1	A	188	LYS	2.1
1	B	148	CYS	2.1
1	B	168	ILE	2.1
1	B	341	THR	2.1
1	B	131	GLY	2.0
1	B	209	HIS	2.0
1	A	172	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	ACT	B	607	4/4	0.85	0.26	6.74	69,69,70,71	0
4	ACT	B	605	4/4	0.91	0.24	2.94	65,67,67,68	0
3	GOL	A	603	6/6	0.87	0.17	1.03	62,63,65,66	0
3	GOL	B	603	6/6	0.83	0.16	0.17	72,73,75,77	0
4	ACT	A	605	4/4	0.88	0.17	-0.32	64,66,67,68	0
2	SO4	B	601	5/5	0.98	0.12	-0.89	59,59,61,61	0
2	SO4	A	602	5/5	0.96	0.13	-1.24	59,63,66,66	0
2	SO4	A	601	5/5	0.98	0.12	-1.46	54,54,56,57	0
2	SO4	B	602	5/5	0.91	0.13	-1.66	78,82,84,85	0
4	ACT	A	607	4/4	0.87	0.14	-	65,65,67,67	0
3	GOL	B	604	6/6	0.90	0.22	-	55,55,58,58	0
4	ACT	B	606	4/4	0.94	0.10	-	71,73,73,74	0
4	ACT	A	606	4/4	0.96	0.13	-	68,68,70,70	0
3	GOL	A	604	6/6	0.90	0.21	-	47,49,51,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.