



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:56 PM GMT

PDB ID : 4DVF  
Title : Crystal structure of BACE1 with its inhibitor  
Authors : Xu, Y.C.; Chen, W.Y.; Li, L.; Chen, T.T.  
Deposited on : 2012-02-23  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> <span style="position: absolute; top: -5px; left: 0;">%</span> </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <span>79%</span> <span>7% • 14%</span> </div> </div>
1	B	433	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> <span style="position: absolute; top: -5px; left: 0;">6%</span> </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <span>75%</span> <span>11% • 14%</span> </div> </div>
2	C	7	<div> <div style="width: 100%; height: 10px; background-color: green; position: relative;"> <span style="position: absolute; top: -5px; left: 0;">57%</span> </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <span>29%</span> <span>14%</span> </div> </div>
2	D	7	<div> <div style="width: 100%; height: 10px; background-color: green; position: relative;"> <span style="position: absolute; top: -5px; left: 0;">86%</span> </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <span></span> <span>14%</span> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6500 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-secretase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	374	Total	C	N	O	S	0	0	0
			2913	1872	486	541	14			
1	B	373	Total	C	N	O	S	0	1	0
			2911	1869	481	547	14			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-39	MET	-	EXPRESSION TAG	UNP P56817
A	-38	GLY	-	EXPRESSION TAG	UNP P56817
A	-37	SER	-	EXPRESSION TAG	UNP P56817
A	-36	SER	-	EXPRESSION TAG	UNP P56817
A	-35	HIS	-	EXPRESSION TAG	UNP P56817
A	-34	HIS	-	EXPRESSION TAG	UNP P56817
A	-33	HIS	-	EXPRESSION TAG	UNP P56817
A	-32	HIS	-	EXPRESSION TAG	UNP P56817
A	-31	HIS	-	EXPRESSION TAG	UNP P56817
A	-30	HIS	-	EXPRESSION TAG	UNP P56817
A	-29	SER	-	EXPRESSION TAG	UNP P56817
A	-28	ALA	-	EXPRESSION TAG	UNP P56817
A	-27	GLY	-	EXPRESSION TAG	UNP P56817
A	-26	GLU	-	EXPRESSION TAG	UNP P56817
A	-25	ASN	-	EXPRESSION TAG	UNP P56817
A	-24	LEU	-	EXPRESSION TAG	UNP P56817
A	-23	TYR	-	EXPRESSION TAG	UNP P56817
A	-22	PHE	-	EXPRESSION TAG	UNP P56817
A	-21	GLN	-	EXPRESSION TAG	UNP P56817
A	-20	GLY	-	EXPRESSION TAG	UNP P56817
A	-19	THR	-	EXPRESSION TAG	UNP P56817
A	75	ALA	LYS	ENGINEERED MUTATION	UNP P56817
A	77	ALA	GLU	ENGINEERED MUTATION	UNP P56817
B	-39	MET	-	EXPRESSION TAG	UNP P56817
B	-38	GLY	-	EXPRESSION TAG	UNP P56817

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-37	SER	-	EXPRESSION TAG	UNP P56817
B	-36	SER	-	EXPRESSION TAG	UNP P56817
B	-35	HIS	-	EXPRESSION TAG	UNP P56817
B	-34	HIS	-	EXPRESSION TAG	UNP P56817
B	-33	HIS	-	EXPRESSION TAG	UNP P56817
B	-32	HIS	-	EXPRESSION TAG	UNP P56817
B	-31	HIS	-	EXPRESSION TAG	UNP P56817
B	-30	HIS	-	EXPRESSION TAG	UNP P56817
B	-29	SER	-	EXPRESSION TAG	UNP P56817
B	-28	ALA	-	EXPRESSION TAG	UNP P56817
B	-27	GLY	-	EXPRESSION TAG	UNP P56817
B	-26	GLU	-	EXPRESSION TAG	UNP P56817
B	-25	ASN	-	EXPRESSION TAG	UNP P56817
B	-24	LEU	-	EXPRESSION TAG	UNP P56817
B	-23	TYR	-	EXPRESSION TAG	UNP P56817
B	-22	PHE	-	EXPRESSION TAG	UNP P56817
B	-21	GLN	-	EXPRESSION TAG	UNP P56817
B	-20	GLY	-	EXPRESSION TAG	UNP P56817
B	-19	THR	-	EXPRESSION TAG	UNP P56817
B	75	ALA	LYS	ENGINEERED MUTATION	UNP P56817
B	77	ALA	GLU	ENGINEERED MUTATION	UNP P56817

- Molecule 2 is a protein called METHYL (2S)-1-[(2R,5S,8S,12S,13S)-2,13-DIBENZYL-12-HYDROXY-3,5-DIMETHYL-8-(2-METHYLPROPYL)-15-(3-[(METHYLSULFONYL)AMINO]-5-[[[(1R)-1-PHENYLETHYL]CARBAMOYL}PHENYL)-4,7,10,15-TETRAOXO-3,6,9,14-TETRAAZAPENTADECAN-1-OYL]PYRROLIDINE-2-CARBOXYLATE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	7	Total	C	N	O	S	0	0	0
			72	53	7	11	1			
2	D	7	Total	C	N	O	S	0	0	0
			72	53	7	11	1			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	291	Total	O	0	0
			291	291		
3	B	234	Total	O	0	0
			234	234		
3	C	3	Total	O	0	0
			3	3		



Continued on next page...

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	4	Total	O	0	0
			4	4		



- Molecule 1: Beta-secretase 1



THYL|CARBAMOYL}PHENYL)-4,7,10,15-TETRAOXO-3,6,9,14-TETRAAZAPENTADECAN  
-1-OYL|PYRROLIDINE-2-CARBOXYLATE

Chain D:  86% 14%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.57Å 85.82Å 176.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.48 – 1.80 48.94 – 1.80	Depositor EDS
% Data completeness (in resolution range)	91.2 (35.48-1.80) 96.6 (48.94-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.67 (at 1.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, $R_{free}$	0.207 , 0.238 0.211 , 0.241	Depositor DCC
$R_{free}$ test set	4116 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	25.2	Xtriage
Anisotropy	0.341	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 49.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 81739 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6500	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: QSC, PSA, USC, ZAE, PLJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/2987	0.55	0/4061
1	B	0.35	0/2988	0.52	0/4067
2	C	1.57	0/12	1.30	0/15
2	D	1.50	0/12	1.35	0/15
All	All	0.37	0/5999	0.54	0/8158

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	2
2	D	0	2
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	3	PSA	Mainchain,Peptide
2	D	3	PSA	Mainchain,Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2913	0	2827	20	0
1	B	2911	0	2803	31	0
2	C	72	0	67	2	0
2	D	72	0	67	2	0
3	A	291	0	0	5	0
3	B	234	0	0	3	0
3	C	3	0	0	0	0
3	D	4	0	0	0	0
All	All	6500	0	5764	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:271:GLN:H	1:B:271:GLN:CD	1.85	0.80
1:A:278:ASN:HB2	3:A:671:HOH:O	1.84	0.77
1:B:239:LYS:HE2	3:B:661:HOH:O	1.98	0.64
1:B:298[B]:ILE:HG13	1:B:341:PHE:CZ	2.33	0.63
1:A:11:GLY:HA2	1:A:307:ARG:HH21	1.64	0.61
1:B:386:ILE:O	1:B:386:ILE:HG22	2.00	0.60
1:A:65:LYS:HE3	1:A:80:LEU:HD12	1.85	0.59
1:A:9:LYS:HE2	1:A:114:ASN:HB2	1.86	0.57
1:B:125:GLU:OE2	1:B:195:ARG:NH1	2.38	0.56
1:B:14:TYR:CE2	1:B:170:VAL:HG13	2.43	0.54
1:B:149:LEU:C	1:B:149:LEU:HD23	2.27	0.54
1:A:371:GLU:HG2	3:A:790:HOH:O	2.06	0.54
1:A:224:LYS:HE2	3:A:773:HOH:O	2.10	0.52
1:A:61:ARG:HD3	3:A:598:HOH:O	2.09	0.52
1:A:298:ILE:HB	1:A:341:PHE:CZ	2.45	0.51
1:B:151:SER:OG	1:B:175:ILE:HB	2.11	0.50
1:A:333:MET:HE3	1:A:337:ILE:HG21	1.93	0.50
1:B:205:ARG:HB3	1:B:286:TYR:HB2	1.94	0.49
1:B:10:SER:OG	1:B:169:SER:HB3	2.11	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:11:GLY:HA2	1:A:307:ARG:NH2	2.28	0.48
1:B:149:LEU:HD22	1:B:178:GLY:CA	2.43	0.48
1:A:155:CYS:O	1:A:170:VAL:HG22	2.13	0.48
1:B:222:TYR:HA	1:B:223:ASP:HA	1.69	0.47
1:B:149:LEU:HD23	1:B:150:PHE:N	2.29	0.47
1:A:73:GLN:O	2:C:3:PSA:HE1	2.15	0.47
1:A:15:TYR:CD1	1:A:28:ASN:HB3	2.50	0.46
1:B:73:GLN:O	2:D:3:PSA:HE1	2.16	0.46
1:A:11:GLY:CA	1:A:307:ARG:HH21	2.29	0.46
1:A:364:GLU:HG2	3:A:735:HOH:O	2.15	0.46
1:A:2:MET:HG2	1:A:90:GLY:HA2	1.97	0.45
1:A:222:TYR:HA	1:A:223:ASP:HA	1.75	0.45
1:B:45:HIS:CG	1:B:46:PRO:HD2	2.51	0.45
1:A:385:ASN:C	1:A:386:ILE:HG12	2.37	0.44
1:B:242:GLU:O	1:B:246:LYS:HG2	2.18	0.44
1:B:271:GLN:H	1:B:271:GLN:NE2	2.14	0.43
1:B:49:HIS:CD2	3:B:703:HOH:O	2.72	0.43
1:B:126:ILE:HG23	1:B:197:TRP:HB2	2.00	0.42
1:B:301:LEU:HB3	1:B:302:PRO:HD2	2.00	0.42
1:A:245:VAL:HG12	1:A:249:LYS:HD2	2.02	0.42
1:B:244:ALA:O	1:B:248:ILE:HG13	2.19	0.42
1:B:2:MET:HG2	1:B:90:GLY:HA2	2.01	0.42
1:B:288:MET:HE2	1:B:379:MET:HB3	2.02	0.42
1:B:337:ILE:O	1:B:341:PHE:HD1	2.03	0.42
1:B:246:LYS:HD3	1:B:246:LYS:HA	1.78	0.41
1:A:42:ALA:CB	1:A:101:ALA:HB1	2.50	0.41
1:B:30:LEU:CD2	2:D:3:PSA:HD2	2.49	0.41
1:B:11:GLY:HA2	1:B:307:ARG:NH2	2.35	0.41
1:B:11:GLY:HA2	1:B:307:ARG:HH21	1.84	0.41
1:B:47:PHE:CE2	1:B:111:ASN:HB2	2.55	0.41
2:C:5:ALA:HA	2:C:6:ZAE:H11	1.92	0.41
1:B:252:SER:HA	3:B:623:HOH:O	2.21	0.41
1:B:359:CYS:O	1:B:359:CYS:SG	2.79	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	368/433 (85%)	361 (98%)	7 (2%)	0	100	100
1	B	368/433 (85%)	357 (97%)	10 (3%)	1 (0%)	46	29
2	C	2/7 (29%)	2 (100%)	0	0	100	100
2	D	2/7 (29%)	2 (100%)	0	0	100	100
All	All	740/880 (84%)	722 (98%)	17 (2%)	1 (0%)	56	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	10	SER

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/368 (84%)	305 (98%)	6 (2%)	65	52
1	B	311/368 (84%)	303 (97%)	8 (3%)	54	37
2	C	1/1 (100%)	1 (100%)	0	100	100
2	D	1/1 (100%)	1 (100%)	0	100	100
All	All	624/738 (85%)	610 (98%)	14 (2%)	60	45

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LYS
1	A	61	ARG
1	A	169	SER
1	A	170	VAL
1	A	197	TRP
1	A	271	GLN
1	B	104	GLU
1	B	138	ASP
1	B	149	LEU
1	B	197	TRP
1	B	211	GLN
1	B	279	ILE
1	B	362	HIS
1	B	380	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	360	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PSA	C	3	2	14,14,15	3.24	6 (42%)	15,17,19	1.34	2 (13%)
2	ZAE	C	6	2	10,12,13	3.94	6 (60%)	10,14,16	0.98	1 (10%)
2	PLJ	C	7	2	9,9,9	1.25	1 (11%)	10,11,11	1.55	2 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PSA	D	3	2	14,14,15	3.31	7 (50%)	15,17,19	1.58	2 (13%)
2	ZAE	D	6	-	10,12,13	4.01	6 (60%)	10,14,16	1.31	2 (20%)
2	PLJ	D	7	-	9,9,9	1.21	1 (11%)	10,11,11	1.37	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PSA	C	3	2	-	0/11/11/12	0/1/1/1
2	ZAE	C	6	2	-	0/4/8/10	0/1/1/1
2	PLJ	C	7	2	-	0/6/13/13	0/1/1/1
2	PSA	D	3	2	-	0/11/11/12	0/1/1/1
2	ZAE	D	6	-	-	0/4/8/10	0/1/1/1
2	PLJ	D	7	-	-	0/6/13/13	0/1/1/1

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	PSA	OH-CH	-2.08	1.38	1.43
2	C	7	PLJ	OXT-C	2.36	1.39	1.33
2	D	7	PLJ	OXT-C	2.45	1.39	1.33
2	D	3	PSA	CZ-CE1	4.04	1.48	1.38
2	C	3	PSA	CZ-CE2	4.09	1.48	1.38
2	D	6	ZAE	CZ-CE2	4.26	1.48	1.38
2	C	6	ZAE	CZ-CE1	4.28	1.49	1.38
2	C	3	PSA	CZ-CE1	4.28	1.49	1.38
2	D	3	PSA	CZ-CE2	4.29	1.49	1.38
2	C	6	ZAE	CZ-CE2	4.36	1.49	1.38
2	D	6	ZAE	CZ-CE1	4.45	1.49	1.38
2	C	3	PSA	CE2-CD2	4.67	1.48	1.38
2	C	3	PSA	CD2-CG	4.75	1.48	1.38
2	D	3	PSA	CD2-CG	4.87	1.49	1.38
2	C	6	ZAE	CD1-CG	4.88	1.49	1.38
2	D	3	PSA	CE1-CD1	5.20	1.49	1.38
2	C	3	PSA	CE1-CD1	5.25	1.49	1.38
2	D	3	PSA	CE2-CD2	5.26	1.49	1.38
2	D	6	ZAE	CD1-CG	5.27	1.50	1.38
2	C	6	ZAE	CD2-CG	5.34	1.50	1.38
2	C	6	ZAE	CE1-CD1	5.41	1.50	1.38
2	D	6	ZAE	CD2-CG	5.45	1.50	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	PSA	CD1-CG	5.46	1.50	1.38
2	C	3	PSA	CD1-CG	5.47	1.50	1.38
2	D	6	ZAE	CE2-CD2	5.52	1.50	1.38
2	D	6	ZAE	CE1-CD1	5.70	1.50	1.38
2	C	6	ZAE	CE2-CD2	5.76	1.50	1.38

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	6	ZAE	CG-CB-CA	-3.12	108.89	114.26
2	C	6	ZAE	CG-CB-CA	-2.42	110.09	114.26
2	C	7	PLJ	OXT-C-O	-2.40	118.84	123.79
2	D	6	ZAE	O-C-CA	-2.15	119.77	125.44
2	D	7	PLJ	OXT-C-O	-2.12	119.42	123.79
2	C	3	PSA	OH-CH-CA	2.71	113.50	109.49
2	D	7	PLJ	OXT-C-CA	2.73	118.61	111.52
2	C	7	PLJ	OXT-C-CA	2.80	118.79	111.52
2	D	3	PSA	OH-CH-CM	2.91	116.10	109.34
2	C	3	PSA	OH-CH-CM	3.32	117.06	109.34
2	D	3	PSA	OH-CH-CA	4.52	116.19	109.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	3	PSA	1	0
2	C	6	ZAE	1	0
2	D	3	PSA	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	374/433 (86%)	-0.24	4 (1%) 82 80	17, 27, 43, 63	0
1	B	373/433 (86%)	0.19	24 (6%) 23 18	19, 30, 56, 91	0
2	C	2/7 (28%)	-0.16	0 100 100	20, 20, 20, 21	0
2	D	2/7 (28%)	-0.37	0 100 100	26, 26, 26, 27	0
All	All	751/880 (85%)	-0.03	28 (3%) 45 39	17, 29, 50, 91	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	386	ILE	6.5
1	B	310	GLU	5.9
1	B	169	SER	5.7
1	B	315	SER	5.3
1	B	272	ALA	5.0
1	B	362	HIS	4.7
1	B	316	GLN	4.6
1	B	157	ALA	4.6
1	B	271	GLN	4.1
1	B	267	LEU	3.2
1	B	10	SER	3.2
1	B	317	ASP	3.1
1	B	273	GLY	3.0
1	B	319	CYS	2.7
1	B	254	THR	2.6
1	A	92	ASN	2.5
1	A	316	GLN	2.5
1	B	8	GLY	2.4
1	B	255	GLU	2.3
1	B	361	VAL	2.3
1	B	11	GLY	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	259	ASP	2.2
1	B	309	VAL	2.2
1	A	157	ALA	2.1
1	B	385	ASN	2.1
1	B	257	PHE	2.1
1	B	318	ASP	2.1
1	A	49	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	PLJ	C	7	9/9	0.94	0.14	-	20,25,28,29	0
2	PSA	D	3	14/15	0.94	0.13	-	21,23,25,25	0
2	PLJ	D	7	9/9	0.93	0.09	-	29,31,37,37	0
2	ZAE	C	6	12/13	0.96	0.10	-	21,23,25,27	0
2	ZAE	D	6	12/13	0.91	0.09	-	25,31,35,35	0
2	PSA	C	3	14/15	0.96	0.10	-	17,19,21,22	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.