



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 07:04 PM GMT

PDB ID : 1DW1
Title : STRUCTURE OF THE CYANIDE COMPLEX OF SHP, AN OXYGEN BINDING CYTOCHROME C
Authors : Leys, D.; Backers, K.; Meyer, T.E.; Hagen, W.R.; Cusanovich, M.A.; Van Beeumen, J.J.
Deposited on : 2000-01-24
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	112	<div> <div>4%</div> <div> <div></div> <div>73%</div> <div>23%</div> <div>.</div> </div> </div>
1	B	112	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>13%</div> <div>..</div> </div> </div>
1	C	112	<div> <div></div> <div> <div></div> <div>80%</div> <div>17%</div> <div>..</div> </div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 2847 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYTOCHROME C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	112	Total	C	N	O	S	0	1	0
			801	488	149	160	4			
1	B	112	Total	C	N	O	S	0	0	0
			794	483	143	164	4			
1	C	112	Total	C	N	O	S	0	1	0
			801	486	150	161	4			

- Molecule 2 is CYANIDE ION (three-letter code: CYN) (formula: CN).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			2	1	1		
2	B	1	Total	C	N	0	0
			2	1	1		
2	C	1	Total	C	N	0	0
			2	1	1		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	1
			45	34	1	4	6		
3	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	101	Total	O	0	0
			101	101		
4	B	95	Total	O	0	0
			95	95		
4	C	118	Total	O	0	0
			118	118		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CYTOCHROME C



• Molecule 1: CYTOCHROME C



• Molecule 1: CYTOCHROME C



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	83.34Å 103.50Å 113.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 28.70 – 1.79	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-1.90) 93.9 (28.70-1.79)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 1.79Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.171 , 0.198 0.161 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	20.7	Xtriage
Anisotropy	0.433	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 45700 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2847	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, CYN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.26	15/823 (1.8%)	1.98	25/1120 (2.2%)
1	B	1.19	10/809 (1.2%)	1.78	19/1105 (1.7%)
1	C	1.08	9/823 (1.1%)	1.53	13/1122 (1.2%)
All	All	1.18	34/2455 (1.4%)	1.77	57/3347 (1.7%)

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	34	THR	CA-CB	11.62	1.83	1.53
1	B	61	LYS	CA-C	9.93	1.78	1.52
1	B	58	ARG	CA-CB	9.40	1.74	1.53
1	B	61	LYS	C-O	-9.26	1.05	1.23
1	A	23	GLU	CA-CB	8.80	1.73	1.53
1	B	15	ALA	C-O	7.55	1.37	1.23
1	B	58	ARG	CA-C	-7.55	1.33	1.52
1	C	79	ALA	CA-CB	7.49	1.68	1.52
1	C	70	ALA	C-O	7.28	1.37	1.23
1	C	18	ALA	CA-CB	7.24	1.67	1.52
1	C	16	ALA	C-O	6.80	1.36	1.23
1	A	111	ALA	C-O	6.72	1.36	1.23
1	A	107	ALA	CA-CB	6.71	1.66	1.52
1	C	20	ALA	C-O	6.56	1.35	1.23
1	A	27	ALA	C-O	6.47	1.35	1.23
1	A	111	ALA	CA-C	-6.38	1.36	1.52
1	B	58	ARG	C-O	6.33	1.35	1.23
1	C	64	ALA	CA-CB	6.20	1.65	1.52
1	C	6	ALA	CA-CB	6.14	1.65	1.52
1	A	79	ALA	CA-CB	6.04	1.65	1.52
1	B	10	ALA	CA-CB	6.00	1.65	1.52
1	A	70	ALA	CA-C	-5.99	1.37	1.52
1	A	64	ALA	CA-CB	5.94	1.65	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	54	ALA	C-O	5.84	1.34	1.23
1	A	49	ALA	CA-CB	5.73	1.64	1.52
1	B	6	ALA	CA-CB	5.63	1.64	1.52
1	A	49	ALA	CA-C	-5.54	1.38	1.52
1	A	67	ALA	CA-CB	5.46	1.64	1.52
1	C	70	ALA	CA-CB	5.43	1.63	1.52
1	A	49	ALA	C-O	5.42	1.33	1.23
1	B	111	ALA	CA-CB	5.40	1.63	1.52
1	C	22	ALA	CA-CB	5.24	1.63	1.52
1	B	67	ALA	C-O	5.15	1.33	1.23
1	A	34	THR	C-O	5.04	1.32	1.23

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	2	ASP	N-CA-CB	25.21	155.98	110.60
1	A	80[A]	ARG	CD-NE-CZ	19.51	150.91	123.60
1	A	80[B]	ARG	CD-NE-CZ	19.51	150.91	123.60
1	A	87	ARG	NE-CZ-NH1	17.62	129.11	120.30
1	A	87	ARG	NE-CZ-NH2	-13.96	113.32	120.30
1	A	80[A]	ARG	NE-CZ-NH1	12.60	126.60	120.30
1	A	80[B]	ARG	NE-CZ-NH1	12.60	126.60	120.30
1	B	74	ARG	NE-CZ-NH2	11.83	126.22	120.30
1	A	87	ARG	CD-NE-CZ	10.95	138.93	123.60
1	C	80[A]	ARG	CD-NE-CZ	10.17	137.84	123.60
1	C	80[B]	ARG	CD-NE-CZ	10.17	137.84	123.60
1	C	53	ARG	NE-CZ-NH1	9.60	125.10	120.30
1	B	61	LYS	CA-C-N	-8.61	98.27	117.20
1	A	80[A]	ARG	NE-CZ-NH2	-8.10	116.25	120.30
1	A	80[B]	ARG	NE-CZ-NH2	-8.10	116.25	120.30
1	A	58	ARG	NE-CZ-NH1	7.98	124.29	120.30
1	B	67	ALA	CB-CA-C	-7.87	98.30	110.10
1	A	111	ALA	CA-C-N	7.64	134.02	117.20
1	B	61	LYS	O-C-N	7.22	134.25	122.70
1	C	77	ASP	CB-CG-OD1	7.22	124.80	118.30
1	C	53	ARG	NE-CZ-NH2	-7.06	116.77	120.30
1	C	44	THR	N-CA-CB	-7.05	96.91	110.30
1	B	15	ALA	O-C-N	-6.83	111.78	122.70
1	C	70	ALA	O-C-N	-6.82	111.79	122.70
1	B	58	ARG	CA-C-N	6.73	132.01	117.20
1	B	2	ASP	CB-CG-OD2	-6.62	112.34	118.30
1	A	58	ARG	NE-CZ-NH2	-6.55	117.03	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	18	ALA	N-CA-CB	-6.55	100.94	110.10
1	C	104	ASP	CB-CG-OD2	6.54	124.18	118.30
1	B	53	ARG	NE-CZ-NH2	-6.45	117.08	120.30
1	A	34	THR	CA-C-N	6.33	128.85	116.20
1	A	70	ALA	CA-C-N	6.16	130.75	117.20
1	A	34	THR	CB-CA-C	6.12	128.12	111.60
1	C	49	ALA	CB-CA-C	-6.04	101.05	110.10
1	B	62	GLU	CB-CG-CD	5.99	130.38	114.20
1	B	58	ARG	O-C-N	-5.89	113.27	122.70
1	B	2	ASP	CB-CA-C	-5.80	98.79	110.40
1	B	62	GLU	CG-CD-OE1	5.69	129.69	118.30
1	B	50	ASP	CB-CG-OD2	5.51	123.26	118.30
1	B	62	GLU	OE1-CD-OE2	-5.51	116.69	123.30
1	B	62	GLU	CA-C-O	-5.47	108.61	120.10
1	A	2	ASP	CB-CG-OD2	5.44	123.20	118.30
1	A	24	ARG	CD-NE-CZ	5.40	131.16	123.60
1	A	50	ASP	CB-CG-OD1	-5.40	113.44	118.30
1	A	74	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	A	49	ALA	CA-C-N	5.33	128.92	117.20
1	C	53	ARG	CD-NE-CZ	5.33	131.06	123.60
1	A	34	THR	N-CA-CB	-5.31	100.21	110.30
1	B	96	ASP	CB-CG-OD1	5.28	123.05	118.30
1	C	20	ALA	N-CA-CB	-5.24	102.76	110.10
1	A	111	ALA	N-CA-CB	-5.22	102.80	110.10
1	A	2	ASP	CB-CG-OD1	-5.16	113.66	118.30
1	A	72	PRO	N-CA-C	5.15	125.50	112.10
1	A	111	ALA	O-C-N	-5.14	114.47	122.70
1	B	62	GLU	N-CA-C	5.11	124.79	111.00
1	C	50	ASP	CB-CG-OD2	5.08	122.87	118.30
1	C	50	ASP	CB-CG-OD1	-5.05	113.76	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	801	0	758	14	0
1	B	794	0	740	12	0
1	C	801	0	752	7	0
2	A	2	0	0	1	0
2	B	2	0	0	1	0
2	C	2	0	0	0	0
3	A	45	0	0	0	0
3	B	43	0	30	1	0
3	C	43	0	30	4	0
4	A	101	0	0	4	0
4	B	95	0	0	2	0
4	C	118	0	0	0	0
All	All	2847	0	2310	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:58:ARG:CA	1:B:58:ARG:CB	1.74	1.64
1:A:34:THR:CA	1:A:34:THR:CB	1.83	1.55
1:B:61:LYS:CA	1:B:61:LYS:C	1.78	1.48
1:C:46:CYS:SG	3:C:113:HEM:HAC	1.58	1.42
1:B:62:GLU:HA	4:B:197:HOH:O	1.53	1.07
2:B:114:CYN:C	2:B:114:CYN:N	0.81	0.96
1:B:61:LYS:CA	1:B:62:GLU:N	2.37	0.87
1:A:78:SER:HB2	4:A:129:HOH:O	1.74	0.86
1:B:61:LYS:C	1:B:61:LYS:CB	2.52	0.79
1:C:80[A]:ARG:NH2	3:C:113:HEM:O2D	2.16	0.78
1:B:58:ARG:CB	1:B:58:ARG:C	2.50	0.77
1:B:62:GLU:CA	4:B:197:HOH:O	2.22	0.77
1:A:34:THR:CB	1:A:34:THR:N	2.55	0.70
1:C:46:CYS:SG	3:C:113:HEM:C3C	2.87	0.66
1:A:59:THR:HB	4:A:122:HOH:O	2.02	0.58
1:A:65:PRO:O	1:A:71:THR:HG23	2.05	0.56
1:A:73:ASP:OD1	1:A:73:ASP:N	2.41	0.54
1:A:90:ASN:ND2	4:A:215:HOH:O	2.42	0.53
1:C:56:GLN:HA	1:C:63:ILE:HG13	1.91	0.52
1:B:58:ARG:CB	1:B:58:ARG:N	2.66	0.52
1:A:73:ASP:HB3	4:A:211:HOH:O	2.11	0.50
1:A:88:ASN:HD22	2:A:114:CYN:C	2.25	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:74:ARG:HD3	3:C:113:HEM:O1A	2.17	0.44
1:C:42:SER:OG	1:C:44:THR:HB	2.19	0.43
1:B:80:ARG:NH2	3:B:113:HEM:O2D	2.51	0.43
1:B:61:LYS:N	1:B:61:LYS:C	2.62	0.43
1:A:26:ARG:HG3	1:A:108:TRP:CE2	2.54	0.42
1:A:65:PRO:O	1:A:71:THR:CG2	2.68	0.41
1:A:8:LEU:HD21	1:A:78:SER:HB3	2.03	0.41
1:B:98:THR:HB	1:B:99:PRO:HD2	2.02	0.41
1:B:61:LYS:CA	1:B:62:GLU:H	2.30	0.41
1:C:64:ALA:HA	1:C:65:PRO:HD3	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	111/112 (99%)	111 (100%)	0	0	100	100
1	B	110/112 (98%)	108 (98%)	1 (1%)	1 (1%)	21	9
1	C	111/112 (99%)	109 (98%)	2 (2%)	0	100	100
All	All	332/336 (99%)	328 (99%)	3 (1%)	1 (0%)	46	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	62	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	77/81 (95%)	74 (96%)	3 (4%)	39	27
1	B	77/81 (95%)	76 (99%)	1 (1%)	76	73
1	C	77/81 (95%)	75 (97%)	2 (3%)	54	45
All	All	231/243 (95%)	225 (97%)	6 (3%)	54	45

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	59	THR
1	A	71	THR
1	A	72	PRO
1	B	8	LEU
1	C	44	THR
1	C	90	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	88	ASN
1	A	90	ASN
1	B	88	ASN
1	B	90	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HEM	A	113[A]	-	30,50,50	2.51	9 (30%)	24,82,82	3.14	10 (41%)
3	HEM	A	113[B]	-	30,50,50	2.51	9 (30%)	24,82,82	3.14	10 (41%)
2	CYN	A	114	3	0,1,1	0.00	-	0,0,0	0.00	-
3	HEM	B	113	1,2	30,50,50	2.93	10 (33%)	24,82,82	3.26	12 (50%)
2	CYN	B	114	3	0,1,1	0.00	-	0,0,0	0.00	-
3	HEM	C	113	1,2	30,50,50	2.61	9 (30%)	24,82,82	3.61	10 (41%)
2	CYN	C	114	3	0,1,1	0.00	-	0,0,0	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	113[A]	-	-	0/10/54/54	0/0/8/8
3	HEM	A	113[B]	-	-	0/10/54/54	0/0/8/8
2	CYN	A	114	3	-	0/0/0/0	0/0/0/0
3	HEM	B	113	1,2	-	0/10/54/54	0/0/8/8
2	CYN	B	114	3	-	0/0/0/0	0/0/0/0
3	HEM	C	113	1,2	-	0/10/54/54	0/0/8/8
2	CYN	C	114	3	-	0/0/0/0	0/0/0/0

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	113	HEM	C3B-C4B	-9.78	1.43	1.51
3	C	113	HEM	C3B-C4B	-7.68	1.45	1.51
3	B	113	HEM	C2D-C3D	-7.11	1.33	1.54
3	A	113[B]	HEM	C3B-C4B	-7.10	1.45	1.51
3	A	113[A]	HEM	C3B-C4B	-7.10	1.45	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	113	HEM	C2D-C3D	-7.02	1.33	1.54
3	A	113[B]	HEM	C2D-C3D	-6.73	1.34	1.54
3	A	113[A]	HEM	C2D-C3D	-6.73	1.34	1.54
3	B	113	HEM	C3D-C4D	-5.75	1.44	1.51
3	A	113[B]	HEM	C3D-C4D	-5.14	1.45	1.51
3	A	113[A]	HEM	C3D-C4D	-5.14	1.45	1.51
3	C	113	HEM	C3D-C4D	-4.91	1.45	1.51
3	B	113	HEM	C2C-C1C	-3.78	1.45	1.52
3	A	113[B]	HEM	C2C-C1C	-3.52	1.45	1.52
3	A	113[A]	HEM	C2C-C1C	-3.52	1.45	1.52
3	C	113	HEM	C2C-C1C	-2.97	1.46	1.52
3	B	113	HEM	C2B-C1B	-2.35	1.44	1.51
3	B	113	HEM	C4C-NC	2.01	1.38	1.36
3	A	113[B]	HEM	CBB-CAB	2.04	1.41	1.29
3	A	113[A]	HEM	CBB-CAB	2.04	1.41	1.29
3	B	113	HEM	CBB-CAB	2.26	1.42	1.29
3	A	113[B]	HEM	C1C-NC	2.28	1.38	1.36
3	A	113[A]	HEM	C1C-NC	2.28	1.38	1.36
3	A	113[B]	HEM	C3B-CAB	2.30	1.55	1.51
3	A	113[A]	HEM	C3B-CAB	2.30	1.55	1.51
3	C	113	HEM	C3B-CAB	2.39	1.55	1.51
3	B	113	HEM	CAA-C2A	2.55	1.56	1.52
3	C	113	HEM	CBB-CAB	2.59	1.44	1.29
3	C	113	HEM	CAA-C2A	2.65	1.56	1.52
3	B	113	HEM	C3B-CAB	2.77	1.56	1.51
3	A	113[B]	HEM	C3C-CAC	2.87	1.56	1.51
3	A	113[A]	HEM	C3C-CAC	2.87	1.56	1.51
3	A	113[B]	HEM	CAA-C2A	2.96	1.57	1.52
3	A	113[A]	HEM	CAA-C2A	2.96	1.57	1.52
3	B	113	HEM	C3C-CAC	3.05	1.57	1.51
3	C	113	HEM	C4C-NC	3.26	1.40	1.36
3	C	113	HEM	C3C-CAC	3.46	1.57	1.51

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	113	HEM	C3B-CAB-CBB	-8.91	110.79	124.46
3	B	113	HEM	C3B-CAB-CBB	-7.09	113.58	124.46
3	A	113[B]	HEM	C3B-CAB-CBB	-7.08	113.60	124.46
3	A	113[A]	HEM	C3B-CAB-CBB	-7.08	113.60	124.46
3	C	113	HEM	C3C-CAC-CBC	-6.40	114.64	124.46
3	B	113	HEM	CAA-C2A-C1A	-5.42	121.13	127.01

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	113	HEM	CAA-C2A-C1A	-5.19	121.37	127.01
3	A	113[B]	HEM	C3C-CAC-CBC	-5.03	116.73	124.46
3	A	113[A]	HEM	C3C-CAC-CBC	-5.03	116.73	124.46
3	C	113	HEM	CMA-C3A-C4A	-3.48	122.61	128.36
3	A	113[B]	HEM	CMA-C3A-C4A	-3.29	122.92	128.36
3	A	113[A]	HEM	CMA-C3A-C4A	-3.29	122.92	128.36
3	B	113	HEM	CMA-C3A-C4A	-3.18	123.10	128.36
3	B	113	HEM	C3C-CAC-CBC	-3.09	119.71	124.46
3	A	113[B]	HEM	CMA-C3A-C2A	2.13	129.69	125.24
3	A	113[A]	HEM	CMA-C3A-C2A	2.13	129.69	125.24
3	B	113	HEM	CMA-C3A-C2A	2.31	130.07	125.24
3	B	113	HEM	CAD-C3D-C4D	2.44	121.06	112.47
3	B	113	HEM	CAD-CBD-CGD	2.90	124.84	113.02
3	C	113	HEM	CAD-C3D-C4D	3.00	123.06	112.47
3	A	113[B]	HEM	CMD-C2D-C3D	3.32	129.02	114.35
3	A	113[A]	HEM	CMD-C2D-C3D	3.32	129.02	114.35
3	B	113	HEM	CMD-C2D-C3D	3.36	129.23	114.35
3	C	113	HEM	CMD-C2D-C3D	3.53	129.96	114.35
3	A	113[B]	HEM	CAD-C3D-C4D	3.55	124.98	112.47
3	A	113[A]	HEM	CAD-C3D-C4D	3.55	124.98	112.47
3	A	113[B]	HEM	C2D-C3D-C4D	4.13	108.50	101.50
3	A	113[A]	HEM	C2D-C3D-C4D	4.13	108.50	101.50
3	C	113	HEM	C2D-C3D-C4D	4.30	108.79	101.50
3	B	113	HEM	C2D-C3D-C4D	4.35	108.88	101.50
3	A	113[B]	HEM	CAD-C3D-C2D	4.62	126.51	113.22
3	A	113[A]	HEM	CAD-C3D-C2D	4.62	126.51	113.22
3	B	113	HEM	CMB-C2B-C3B	4.96	128.92	116.53
3	C	113	HEM	CAD-C3D-C2D	5.18	128.10	113.22
3	A	113[B]	HEM	CMC-C2C-C3C	5.34	129.86	116.53
3	A	113[A]	HEM	CMC-C2C-C3C	5.34	129.86	116.53
3	C	113	HEM	CMB-C2B-C3B	5.57	130.43	116.53
3	B	113	HEM	CAD-C3D-C2D	5.85	130.03	113.22
3	A	113[B]	HEM	CMB-C2B-C3B	5.96	131.41	116.53
3	A	113[A]	HEM	CMB-C2B-C3B	5.96	131.41	116.53
3	B	113	HEM	CMC-C2C-C3C	5.99	131.48	116.53
3	C	113	HEM	CMC-C2C-C3C	6.34	132.35	116.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	114	CYN	1	0
3	B	113	HEM	1	0
2	B	114	CYN	1	0
3	C	113	HEM	4	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	112/112 (100%)	-0.49	4 (3%) 46 50	16, 23, 37, 55	0
1	B	112/112 (100%)	-0.48	2 (1%) 71 74	17, 23, 41, 51	0
1	C	112/112 (100%)	-0.36	0 100 100	15, 23, 38, 47	0
All	All	336/336 (100%)	-0.44	6 (1%) 71 74	15, 23, 39, 55	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	62	GLU	6.0
1	B	1	GLY	3.7
1	A	1	GLY	3.4
1	A	2	ASP	2.9
1	A	72	PRO	2.7
1	A	38	PRO	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	HEM	A	113[B]	43/43	0.99	0.08	0.35	15,18,27,32	2
3	HEM	A	113[A]	43/43	0.99	0.08	0.35	15,18,27,32	2
3	HEM	B	113	43/43	0.98	0.08	0.31	16,19,34,41	0
3	HEM	C	113	43/43	0.97	0.08	-0.45	16,20,32,41	0
2	CYN	A	114	2/2	0.99	0.06	-	19,19,19,23	0
2	CYN	B	114	2/2	0.99	0.08	-	22,22,22,31	0
2	CYN	C	114	2/2	0.97	0.07	-	23,23,23,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.