



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:15 AM GMT

PDB ID : 3DX9  
Title : Crystal Structure of the DM1 TCR at 2.75Å  
Authors : Archbold, J.K.; Macdonald, W.A.; Gras, S.; Rossjohn, J.  
Deposited on : 2008-07-24  
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3340 (2.80-2.72)
Clashscore	102246	3829 (2.80-2.72)
Ramachandran outliers	100387	3767 (2.80-2.72)
Sidechain outliers	100360	3770 (2.80-2.72)
RSRZ outliers	91569	3352 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	198	<div> <div>21%</div> <div>72%</div> <div>22%</div> <div>6%</div> </div>
1	C	198	<div> <div>21%</div> <div>73%</div> <div>22%</div> <div>6%</div> </div>
2	B	244	<div> <div>7%</div> <div>80%</div> <div>14%</div> <div>5%</div> </div>
2	D	244	<div> <div>8%</div> <div>79%</div> <div>17%</div> <div>.</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7257 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DM1 T cell receptor alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	198	Total	C	N	O	S	0	1	0
			1557	975	262	310	10			
1	C	198	Total	C	N	O	S	0	1	0
			1556	973	262	311	10			

- Molecule 2 is a protein called DM1 T cell receptor beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	244	Total	C	N	O	S	0	1	0
			1992	1249	358	379	6			
2	D	244	Total	C	N	O	S	0	1	0
			1993	1249	358	380	6			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	21	Total	O	0	0
			21	21		
3	B	53	Total	O	0	0
			53	53		
3	C	31	Total	O	0	0
			31	31		
3	D	54	Total	O	0	0
			54	54		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DM1 T cell receptor alpha chain




- Molecule 1: DM1 T cell receptor alpha chain



- Molecule 2: DM1 T cell receptor beta chain



- Molecule 2: DM1 T cell receptor beta chain





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	41.74Å 70.73Å 154.41Å 90.00° 94.69° 90.00°	Depositor
Resolution (Å)	154.30 – 2.75 52.07 – 2.75	Depositor EDS
% Data completeness (in resolution range)	96.2 (154.30-2.75) 96.2 (52.07-2.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.65 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.2	Depositor
R, $R_{free}$	0.204 , 0.263 0.228 , 0.272	Depositor DCC
$R_{free}$ test set	1159 reflections (5.39%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.5	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 57.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 22660 reflections (0.004%)	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	7257	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.32 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.0090e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.59	1/1596 (0.1%)	0.75	2/2170 (0.1%)
1	C	0.57	1/1595 (0.1%)	0.76	3/2171 (0.1%)
2	B	0.62	2/2049 (0.1%)	1.18	4/2782 (0.1%)
2	D	0.55	0/2050	0.74	3/2782 (0.1%)
All	All	0.58	4/7290 (0.1%)	0.89	12/9905 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	9
1	C	0	9
2	B	1	2
2	D	0	1
All	All	1	21

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	61	GLN	C-N	9.75	1.56	1.34
1	A	113	VAL	CB-CG1	-7.69	1.36	1.52
1	C	113	VAL	CB-CG2	-7.69	1.36	1.52
2	B	184	CYS	CB-SG	-5.77	1.72	1.81

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	61	GLN	O-C-N	-41.52	56.28	122.70
2	B	61	GLN	CA-C-N	25.34	172.94	117.20
2	B	194	PRO	N-CA-C	6.37	128.67	112.10
1	A	30	GLY	N-CA-C	-6.35	97.23	113.10

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	D	194	PRO	N-CA-C	6.26	128.38	112.10

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	66	LEU	CA

5 of 21 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	116	GLY	Peptide
1	A	119	THR	Peptide
1	A	29	SER	Peptide
1	A	55	HIS	Peptide
1	A	58	LYS	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1557	0	1492	36	1
1	C	1556	0	1485	32	0
2	B	1992	0	1886	38	0
2	D	1993	0	1886	33	1
3	A	21	0	0	1	0
3	B	53	0	0	0	0
3	C	31	0	0	1	0
3	D	54	0	0	0	0
All	All	7257	0	6749	129	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 129 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:193:GLN:C	2:B:195:ALA:HB3	1.66	1.15
2:D:193:GLN:C	2:D:195:ALA:HB3	1.67	1.13
2:B:129:ASP:OD2	2:B:131:LYS:HG2	1.52	1.09
2:B:194:PRO:N	2:B:195:ALA:HB3	1.77	1.00
2:D:194:PRO:N	2:D:195:ALA:HB3	1.75	0.99

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:143:ASP:O	2:D:237:THR:OG1[1_455]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/198 (100%)	176 (89%)	16 (8%)	5 (2%)	7	21
1	C	197/198 (100%)	180 (91%)	13 (7%)	4 (2%)	9	27
2	B	243/244 (100%)	226 (93%)	12 (5%)	5 (2%)	9	25
2	D	243/244 (100%)	228 (94%)	12 (5%)	3 (1%)	16	43
All	All	880/884 (100%)	810 (92%)	53 (6%)	17 (2%)	11	28

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	59	ASN
1	A	143	ASP
1	A	144[A]	LYS
1	A	144[B]	LYS
2	B	66	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/177 (101%)	166 (93%)	12 (7%)	20	46
1	C	178/177 (101%)	168 (94%)	10 (6%)	26	56
2	B	217/216 (100%)	200 (92%)	17 (8%)	16	38
2	D	217/216 (100%)	202 (93%)	15 (7%)	19	45
All	All	790/786 (100%)	736 (93%)	54 (7%)	21	46

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	196	LEU
1	C	69	ASN
2	D	188	GLN
2	B	206	ARG
2	B	218	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	219	ASN
1	C	20	ASN
1	C	122	GLN
1	A	203	ASN
2	B	30	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	198/198 (100%)	1.36	41 (20%) <b>1</b> <b>1</b>	40, 42, 43, 45	10 (5%)
1	C	198/198 (100%)	1.26	42 (21%) <b>1</b> <b>1</b>	23, 42, 43, 44	14 (7%)
2	B	244/244 (100%)	0.63	16 (6%) <b>22</b> <b>16</b>	22, 42, 43, 45	18 (7%)
2	D	244/244 (100%)	0.70	19 (7%) <b>16</b> <b>11</b>	37, 42, 43, 45	12 (4%)
All	All	884/884 (100%)	0.95	118 (13%) <b>4</b> <b>3</b>	22, 42, 43, 45	54 (6%)

The worst 5 of 118 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	110	TYR	7.8
1	C	108	GLY	7.1
1	C	110	TYR	6.8
1	A	108	GLY	6.6
1	A	111	GLN	5.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.