



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 21, 2016 – 11:40 AM EDT

PDB ID : 5DX6  
Title : Acetolactate Synthase from *Klebsiella pneumoniae* soaked with beta-fluoropyruvate  
Authors : Latta, A.J.; McLeish, M.J.; Andrews, F.H.  
Deposited on : 2015-09-23  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027939  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027939




# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1609 (1.76-1.76)
Clashscore	102246	1730 (1.76-1.76)
Ramachandran outliers	100387	1711 (1.76-1.76)
Sidechain outliers	100360	1711 (1.76-1.76)
RSRZ outliers	91569	1610 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	579	<div> <div>4%</div> <div>88%</div> <div>5% • 7%</div> </div>
1	B	579	<div> <div>%</div> <div>91%</div> <div>5% •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	65S	B	604	-	-	-	X



## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 9100 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetolactate synthase, catabolic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	541	Total	C	N	O	S	0	2	0
			4064	2575	711	760	18			
1	B	557	Total	C	N	O	S	0	5	0
			4184	2651	726	789	18			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	MET	-	initiating methionine	UNP P27696
B	-18	GLY	-	expression tag	UNP P27696
B	-17	SER	-	expression tag	UNP P27696
B	-16	SER	-	expression tag	UNP P27696
B	-15	HIS	-	expression tag	UNP P27696
B	-14	HIS	-	expression tag	UNP P27696
B	-13	HIS	-	expression tag	UNP P27696
B	-12	HIS	-	expression tag	UNP P27696
B	-11	HIS	-	expression tag	UNP P27696
B	-10	HIS	-	expression tag	UNP P27696
B	-9	SER	-	expression tag	UNP P27696
B	-8	SER	-	expression tag	UNP P27696
B	-7	GLY	-	expression tag	UNP P27696
B	-6	LEU	-	expression tag	UNP P27696
B	-5	VAL	-	expression tag	UNP P27696
B	-4	PRO	-	expression tag	UNP P27696
B	-3	ARG	-	expression tag	UNP P27696
B	-2	GLY	-	expression tag	UNP P27696
B	-1	SER	-	expression tag	UNP P27696
B	0	HIS	-	expression tag	UNP P27696
B	407	SER	THR	conflict	UNP P27696
A	-19	MET	-	initiating methionine	UNP P27696
A	-18	GLY	-	expression tag	UNP P27696
A	-17	SER	-	expression tag	UNP P27696
A	-16	SER	-	expression tag	UNP P27696


*Continued on next page...*



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-15	HIS	-	expression tag	UNP P27696
A	-14	HIS	-	expression tag	UNP P27696
A	-13	HIS	-	expression tag	UNP P27696
A	-12	HIS	-	expression tag	UNP P27696
A	-11	HIS	-	expression tag	UNP P27696
A	-10	HIS	-	expression tag	UNP P27696
A	-9	SER	-	expression tag	UNP P27696
A	-8	SER	-	expression tag	UNP P27696
A	-7	GLY	-	expression tag	UNP P27696
A	-6	LEU	-	expression tag	UNP P27696
A	-5	VAL	-	expression tag	UNP P27696
A	-4	PRO	-	expression tag	UNP P27696
A	-3	ARG	-	expression tag	UNP P27696
A	-2	GLY	-	expression tag	UNP P27696
A	-1	SER	-	expression tag	UNP P27696
A	0	HIS	-	expression tag	UNP P27696
A	407	SER	THR	conflict	UNP P27696

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Mg 2 2	0	0
3	A	2	Total Mg 2 2	0	0


- Molecule 4 is 3-[(4-amino-2-methylpyrimidin-5-yl)methyl]-2-[(1R)-2-fluoro-1-hydroxyethyl]-5-(2-{[(S)-hydroxy(phosphonooxy)phosphoryl]oxy}ethyl)-4-methyl-1,3-thiazol-3-ium (three-letter code: 5GY) (formula: C<sub>14</sub>H<sub>22</sub>FN<sub>4</sub>O<sub>8</sub>P<sub>2</sub>S).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	P	S	0	0
			30	14	1	4	8	2	1		


- Molecule 5 is THIAMINE DIPHOSPHATE (three-letter code: TPP) (formula:  $\text{C}_{12}\text{H}_{19}\text{N}_4\text{O}_7\text{P}_2\text{S}$ ).





Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		

- Molecule 6 is 3-fluoro-2-oxopropanoic acid (three-letter code: 65S) (formula:  $\text{C}_3\text{H}_3\text{FO}_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	F	O	0	0
			7	3	1	3		

- Molecule 7 is water.




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	332	Total 332	O 332	0	0
7	B	448	Total 448	O 448	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetolactate synthase, catabolic



- Molecule 1: Acetolactate synthase, catabolic





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.80Å 134.01Å 110.77Å 90.00° 95.50° 90.00°	Depositor
Resolution (Å)	42.57 – 1.75 42.57 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (42.57-1.75) 95.6 (42.57-1.75)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.93 (at 1.75Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.151 , 0.190 0.158 , 0.200	Depositor DCC
$R_{free}$ test set	1879 reflections (1.59%)	DCC
Wilson B-factor (Å <sup>2</sup> )	22.2	Xtriage
Anisotropy	0.243	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 48.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	9100	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5GY, MG, TPP, PO4, 65S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/4147	0.53	1/5641 (0.0%)
1	B	0.38	0/4279	0.54	0/5830
All	All	0.37	0/8426	0.53	1/11471 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	259	ARG	NE-CZ-NH2	-7.13	116.73	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4064	0	4056	34	0
1	B	4184	0	4140	19	0
2	A	5	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	30	0	0	4	0
5	B	26	0	16	4	0
6	B	7	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	332	0	0	28	2
7	B	448	0	0	11	1
All	All	9100	0	8212	60	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:232:GLU:OE1	7:A:701:HOH:O	1.81	0.97
1:A:542:ASP:OD2	7:A:702:HOH:O	1.99	0.80
1:B:220:GLN:OE1	7:B:701:HOH:O	1.99	0.79
1:A:545:ASP:OD1	7:A:703:HOH:O	2.01	0.78
1:A:28:GLN:NE2	7:A:711:HOH:O	2.16	0.77
1:A:488:TYR:OH	7:A:704:HOH:O	2.04	0.75
1:B:439:ARG:NH1	7:B:705:HOH:O	2.18	0.75
1:B:254:SER:OG	7:B:702:HOH:O	2.04	0.74
1:A:355:GLN:OE1	7:A:706:HOH:O	2.07	0.72
1:A:512:GLY:O	7:A:705:HOH:O	2.07	0.72
1:A:409:ARG:NH1	7:A:713:HOH:O	2.24	0.71
1:A:506:GLU:OE1	7:A:707:HOH:O	2.08	0.71
1:A:265:ASN:OD1	1:A:403:ARG:NH2	2.25	0.70
5:B:603:TPP:HN42	5:B:603:TPP:C2	2.06	0.69
1:A:527:ALA:O	7:A:708:HOH:O	2.09	0.69
1:A:368:GLN:NE2	7:A:716:HOH:O	2.27	0.67
1:A:499:MET:O	7:A:709:HOH:O	2.13	0.67
1:A:381:GLN:NE2	7:A:717:HOH:O	2.28	0.66
4:A:604:5GY:O19	7:A:710:HOH:O	2.14	0.65
1:B:381:GLN:O	1:B:409:ARG:NH1	2.30	0.63
1:B:381:GLN:OE1	1:B:407:SER:OG	2.18	0.61
5:B:603:TPP:H2	5:B:603:TPP:HN42	1.66	0.60
1:A:259:ARG:NH2	7:A:721:HOH:O	2.34	0.60
1:A:527:ALA:HB1	7:A:708:HOH:O	2.02	0.59
1:A:527:ALA:CA	7:A:708:HOH:O	2.50	0.59
1:B:310:GLU:OE2	7:B:704:HOH:O	2.17	0.58
1:B:141:PRO:HB3	1:B:176:VAL:HG12	1.87	0.55
4:A:604:5GY:C06	4:A:604:5GY:N01	2.70	0.54
4:A:604:5GY:P18	7:A:710:HOH:O	2.65	0.54
1:B:269:ASP:OD1	7:B:706:HOH:O	2.19	0.54
1:A:73:LYS:NZ	7:A:728:HOH:O	2.41	0.53

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:523:PRO:CA	7:A:720:HOH:O	2.59	0.50
1:A:486:LYS:NZ	7:A:731:HOH:O	2.45	0.49
1:B:411:ARG:N	7:B:715:HOH:O	2.42	0.49
5:B:603:TPP:H2	7:B:810:HOH:O	2.13	0.49
1:B:118:VAL:HG13	7:B:893:HOH:O	2.12	0.49
1:A:369:PHE:HA	1:A:370:ALA:C	2.33	0.49
1:A:229:ARG:NH1	1:A:329:ASN:OD1	2.46	0.49
1:A:512:GLY:N	7:A:705:HOH:O	2.25	0.48
1:B:352:ARG:NE	7:B:703:HOH:O	2.16	0.48
5:B:603:TPP:N4'	5:B:603:TPP:C2	2.75	0.47
1:A:547:PRO:O	7:A:712:HOH:O	2.20	0.47
1:A:409:ARG:NH1	7:A:735:HOH:O	2.47	0.47
1:A:472:TRP:HB3	1:A:541[A]:VAL:HG21	1.98	0.45
1:B:86:SER:HB3	1:B:127:MET:HE1	1.98	0.45
1:A:527:ALA:CB	7:A:708:HOH:O	2.63	0.45
1:B:367:ASN:O	7:B:707:HOH:O	2.21	0.44
1:B:384:VAL:CG1	1:B:410:ALA:HB2	2.48	0.44
1:A:403:ARG:HA	1:A:403:ARG:HD3	1.84	0.44
1:B:31:GLY:HA3	1:B:78:LEU:O	2.17	0.44
1:A:440:LYS:HE3	1:A:529:MET:HE2	1.98	0.44
1:A:259:ARG:HD2	1:A:259:ARG:N	2.33	0.44
1:A:527:ALA:C	7:A:708:HOH:O	2.53	0.43
1:B:482:ILE:HD11	7:B:936:HOH:O	2.18	0.43
1:A:420:GLN:O	4:A:604:5GY:N01	2.52	0.43
1:B:22:GLU:OE2	1:B:47[B]:SER:HB2	2.20	0.41
1:B:-5:VAL:N	1:B:-4:PRO:CD	2.84	0.41
1:A:338:ARG:NH1	7:A:738:HOH:O	2.50	0.41
1:A:523:PRO:CB	7:A:720:HOH:O	2.69	0.41
1:B:369:PHE:HA	1:B:370:ALA:C	2.41	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:743:HOH:O	7:B:804:HOH:O[2_756]	1.92	0.28
7:A:997:HOH:O	7:A:1010:HOH:O[2_756]	2.08	0.12



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	537/579 (93%)	523 (97%)	14 (3%)	0	100	100
1	B	558/579 (96%)	544 (98%)	14 (2%)	0	100	100
All	All	1095/1158 (95%)	1067 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	421/462 (91%)	415 (99%)	6 (1%)	74	58
1	B	434/462 (94%)	430 (99%)	4 (1%)	84	72
All	All	855/924 (92%)	845 (99%)	10 (1%)	78	62

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	311	GLU
1	A	365	GLN
1	A	368	GLN
1	A	403	ARG
1	A	447	ASP
1	A	529	MET
1	B	28	GLN
1	B	235	HIS

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	B	259	ARG
1	B	354	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	A	601	-	4,4,4	0.83	0	6,6,6	0.22	0
4	5GY	A	604	3	23,31,31	1.20	2 (8%)	28,46,46	2.46	10 (35%)
5	TPP	B	603	3	20,27,27	1.35	3 (15%)	27,40,40	1.95	8 (29%)
6	65S	B	604	-	2,6,6	0.88	0	1,7,7	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means



no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	601	-	-	0/0/0/0	0/0/0/0
4	5GY	A	604	3	-	0/16/23/23	0/2/2/2
5	TPP	B	603	3	-	0/16/17/17	0/2/2/2
6	65S	B	604	-	-	0/0/6/6	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	603	TPP	C5'-C4'	-3.00	1.37	1.42
5	B	603	TPP	PB-O2B	-2.01	1.47	1.54
5	B	603	TPP	C4'-N4'	2.54	1.40	1.34
4	A	604	5GY	C02-N01	2.75	1.41	1.34
4	A	604	5GY	C04-C03	3.18	1.57	1.51

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	604	5GY	C03-C04-N05	-7.15	100.81	113.37
4	A	604	5GY	C25-C24-C12	-3.36	121.79	128.91
5	B	603	TPP	CM4-C4-C5	-3.29	121.95	128.91
5	B	603	TPP	C5'-C7'-N3	-3.12	107.89	113.37
4	A	604	5GY	C04-N05-C24	-2.83	118.93	124.44
5	B	603	TPP	N1'-C2'-N3'	-2.80	120.24	125.50
4	A	604	5GY	N27-C28-N30	-2.71	120.41	125.50
4	A	604	5GY	O15-C14-C13	-2.55	98.72	109.24
4	A	604	5GY	C13-C12-C24	-2.30	124.77	127.34
5	B	603	TPP	C5'-C6'-N1'	-2.21	120.00	123.86
4	A	604	5GY	C12-C24-N05	2.04	112.70	107.94
5	B	603	TPP	C6'-C5'-C4'	2.07	118.58	115.65
5	B	603	TPP	C6'-N1'-C2'	2.34	120.37	115.92
4	A	604	5GY	C29-C28-N27	3.14	120.93	117.05
4	A	604	5GY	C26-N27-C28	3.32	122.23	115.92
5	B	603	TPP	CM2-C2'-N1'	3.33	121.16	117.05
5	B	603	TPP	CM4-C4-N3	5.69	129.41	122.43
4	A	604	5GY	C13-C12-S11	6.16	128.86	120.24

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 8 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	604	5GY	4	0
5	B	603	TPP	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	541/579 (93%)	0.17	23 (4%) 39 45	19, 39, 60, 85	0
1	B	557/579 (96%)	-0.23	8 (1%) 78 84	15, 27, 49, 82	0
All	All	1098/1158 (94%)	-0.03	31 (2%) 56 62	15, 32, 57, 85	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	366	LEU	8.4
1	A	364	ALA	6.6
1	B	-5	VAL	5.4
1	A	115	ALA	5.0
1	A	365	GLN	3.6
1	A	11	ALA	3.6
1	A	9	GLN	3.2
1	A	187	PRO	3.1
1	A	6	PRO	2.9
1	A	524	THR	2.9
1	B	0	HIS	2.9
1	A	7	VAL	2.6
1	B	355	GLN	2.6
1	A	367	ASN	2.5
1	B	336	ASP	2.4
1	A	488	TYR	2.4
1	A	517	SER	2.3
1	B	4	GLN	2.3
1	A	48	SER	2.3
1	A	520	ALA	2.2
1	A	532	ASP	2.2
1	A	265	ASN	2.2
1	A	370	ALA	2.2
1	A	518	ALA	2.2

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	312	ARG	2.2
1	A	176	VAL	2.2
1	A	489	GLN	2.2
1	B	337	HIS	2.2
1	A	519	GLU	2.2
1	B	253	PHE	2.1
1	A	45	LEU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	65S	B	604	7/7	0.83	0.21	4.71	60,65,67,70	0
3	MG	B	601	1/1	0.99	0.10	1.45	19,19,19,19	0
3	MG	B	602	1/1	0.99	0.09	1.19	31,31,31,31	0
4	5GY	A	604	30/30	0.97	0.12	0.62	30,38,51,53	6
5	TPP	B	603	26/26	0.99	0.09	-0.04	17,20,25,29	0
3	MG	A	602	1/1	0.99	0.09	-0.27	41,41,41,41	0
2	PO4	A	601	5/5	0.99	0.06	-0.94	33,33,34,35	0
3	MG	A	603	1/1	0.97	0.05	-1.79	41,41,41,41	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.