



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:41 AM GMT

PDB ID : 2DYU
Title : Helicobacter pylori formamidase AmiF contains a fine-tuned cysteine-glutamate-lysine catalytic triad
Authors : Wang, W.C.; Hung, C.L.
Deposited on : 2006-09-18
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1609 (1.76-1.76)
Clashscore	102246	1730 (1.76-1.76)
Ramachandran outliers	100387	1711 (1.76-1.76)
Sidechain outliers	100360	1711 (1.76-1.76)
RSRZ outliers	91569	1610 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	<div> <div>8%</div> <div>82%</div> <div>14%</div> <div>..</div> </div>
1	B	334	<div> <div>12%</div> <div>82%</div> <div>13%</div> <div>..</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5811 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Formamidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	322	Total	C	N	O	S	0	0	0
			2549	1641	427	469	12			
1	B	322	Total	C	N	O	S	0	0	0
			2549	1641	427	469	12			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	412	Total	O	0	0
			412	412		
2	B	301	Total	O	0	0
			301	301		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Formamidase



• Molecule 1: Formamidase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	147.22Å 147.22Å 72.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 1.75 25.27 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.0 (30.00-1.75) 98.8 (25.27-1.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.174 , 0.215 0.175 , 0.173	Depositor DCC
R_{free} test set	2955 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	17.7	Xtriage
Anisotropy	0.025	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 65.0	EDS
Estimated twinning fraction	0.015 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 58456 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5811	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/2623	0.53	0/3567
1	B	0.32	0/2623	0.51	0/3567
All	All	0.34	0/5246	0.52	0/7134

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2549	0	2480	71	0
1	B	2549	0	2480	75	0
2	A	412	0	0	8	0
2	B	301	0	0	8	0
All	All	5811	0	4960	115	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (115) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:209:TRP:HE1	1:B:170:MET:CE	1.51	1.23

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:166:CYS:SG	1:A:167:HIS:N	2.11	1.19
1:A:170:MET:SD	2:A:733:HOH:O	1.96	1.19
1:B:137:TRP:CD1	1:B:167:HIS:HB2	1.77	1.18
1:A:170:MET:CE	1:B:209:TRP:HE1	1.60	1.13
1:A:170:MET:HE1	1:B:209:TRP:HE1	1.11	1.08
1:B:170:MET:HE3	1:B:203:THR:HG21	1.36	1.08
1:A:209:TRP:HE1	1:B:170:MET:HE1	1.19	1.06
1:B:170:MET:CE	1:B:203:THR:HG21	1.87	1.03
1:B:170:MET:SD	2:B:628:HOH:O	2.17	1.01
1:B:192:TYR:HA	2:B:619:HOH:O	1.59	1.00
1:A:170:MET:CE	1:A:203:THR:HG21	1.95	0.96
1:A:170:MET:HE3	1:A:203:THR:HG21	1.48	0.96
1:B:137:TRP:HD1	1:B:167:HIS:CB	1.79	0.96
1:B:170:MET:HA	1:B:170:MET:HE3	1.50	0.94
1:A:170:MET:HE1	1:B:209:TRP:NE1	1.82	0.93
1:A:170:MET:HA	1:A:170:MET:HE3	1.49	0.92
1:A:19:ILE:HD11	1:A:56:ILE:HG23	1.49	0.92
1:B:137:TRP:HD1	1:B:167:HIS:HB2	1.26	0.91
1:B:137:TRP:CD1	1:B:167:HIS:CB	2.53	0.90
1:A:209:TRP:NE1	1:B:170:MET:CE	2.35	0.90
1:A:209:TRP:HE1	1:B:170:MET:HE2	1.34	0.88
1:A:209:TRP:NE1	1:B:170:MET:HE1	1.88	0.87
1:A:13:GLY:N	2:A:721:HOH:O	2.10	0.84
1:A:170:MET:CE	1:B:209:TRP:NE1	2.41	0.78
1:A:334:LYS:HD2	1:A:334:LYS:H	1.48	0.78
1:A:167:HIS:CE1	1:B:275:ASN:ND2	2.53	0.76
1:A:19:ILE:HD11	1:A:56:ILE:CG2	2.14	0.76
1:B:170:MET:HE1	1:B:203:THR:HG21	1.70	0.72
1:B:23:VAL:CG2	1:B:35:ASN:HD21	2.02	0.71
1:A:167:HIS:HE1	1:B:275:ASN:ND2	1.87	0.71
1:A:170:MET:HA	1:A:170:MET:CE	2.22	0.70
1:A:170:MET:HE1	1:A:203:THR:HG21	1.74	0.70
1:A:209:TRP:NE1	1:B:170:MET:HE2	2.05	0.69
1:A:232:GLU:OE1	2:A:728:HOH:O	2.09	0.69
1:B:136:PRO:O	2:B:623:HOH:O	2.10	0.68
1:B:23:VAL:HG23	1:B:35:ASN:HD21	1.60	0.67
1:B:136:PRO:O	1:B:138:ASN:N	2.29	0.66
1:B:170:MET:CE	1:B:170:MET:HA	2.25	0.65
1:A:170:MET:HE2	1:B:209:TRP:HE1	1.59	0.64
1:A:229:TYR:HA	2:A:723:HOH:O	1.98	0.63
1:B:137:TRP:NE1	1:B:167:HIS:HB2	2.14	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:291:GLU:HG3	1:A:293:ASP:H	1.66	0.59
1:B:292:HIS:HD2	2:B:342:HOH:O	1.84	0.59
1:B:170:MET:HE1	1:B:203:THR:CG2	2.33	0.58
1:B:170:MET:CA	1:B:170:MET:HE3	2.30	0.57
1:B:229:TYR:HA	2:B:627:HOH:O	2.05	0.56
1:A:283:GLY:HA3	1:B:145:PRO:HG3	1.86	0.56
1:A:167:HIS:CE1	1:B:275:ASN:HD21	2.22	0.56
1:A:135:PHE:O	1:A:167:HIS:HD2	1.89	0.55
1:A:275:ASN:ND2	1:B:167:HIS:CE1	2.76	0.54
1:A:170:MET:HE3	1:A:170:MET:CA	2.31	0.53
1:A:170:MET:CA	1:A:170:MET:CE	2.85	0.52
1:A:282:ARG:HG2	1:B:135:PHE:CE1	2.45	0.52
1:B:281:HIS:HD2	1:B:289:GLY:O	1.93	0.52
1:B:170:MET:CE	1:B:170:MET:CA	2.87	0.51
1:A:196:VAL:CG1	1:A:196:VAL:O	2.58	0.50
1:B:167:HIS:CD2	1:B:167:HIS:C	2.84	0.50
1:A:285:VAL:HA	1:B:327:THR:HG22	1.93	0.50
1:A:334:LYS:HD2	1:A:334:LYS:N	2.23	0.50
1:A:167:HIS:CE1	1:B:275:ASN:HD22	2.26	0.49
1:A:194:THR:HG23	1:A:195:GLN:HG2	1.94	0.49
1:B:170:MET:CE	1:B:203:THR:CG2	2.74	0.48
1:A:278:ASN:HD22	1:B:167:HIS:HE1	1.62	0.48
1:A:312:GLU:OE2	1:B:281:HIS:HE1	1.96	0.48
1:B:122:ASP:HB2	1:B:123:PRO:CD	2.43	0.48
1:B:196:VAL:HG23	2:B:514:HOH:O	2.12	0.48
1:B:309:LEU:HB2	1:B:312:GLU:HG3	1.96	0.48
1:A:85:THR:HG22	1:A:103:ILE:CD1	2.43	0.47
1:A:122:ASP:HB2	1:A:123:PRO:CD	2.44	0.47
1:B:193:SER:OG	1:B:193:SER:O	2.32	0.47
1:A:332:PHE:HA	1:A:334:LYS:HE3	1.97	0.47
1:A:247:HIS:O	1:A:252:GLU:OE1	2.33	0.46
1:A:192:TYR:HA	2:A:405:HOH:O	2.14	0.46
1:A:170:MET:HE1	1:B:209:TRP:CD1	2.48	0.46
1:A:287:LYS:HD3	1:A:291:GLU:HB2	1.98	0.46
1:A:272:LEU:HD21	1:B:140:ILE:HG22	1.98	0.45
1:B:194:THR:HG23	1:B:195:GLN:HG2	1.98	0.45
1:B:167:HIS:CD2	1:B:167:HIS:O	2.69	0.45
1:B:133:LYS:HD3	1:B:136:PRO:HA	1.99	0.45
1:B:247:HIS:O	1:B:252:GLU:OE1	2.34	0.44
1:A:196:VAL:HG13	1:A:196:VAL:O	2.16	0.44
1:A:282:ARG:HD2	1:A:294:ALA:HB2	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:209:TRP:CD1	1:B:170:MET:HE1	2.51	0.44
1:A:19:ILE:CD1	1:A:56:ILE:CG2	2.92	0.44
1:B:227:PHE:CZ	1:B:229:TYR:HB2	2.53	0.44
1:A:29:ARG:HD2	2:A:377:HOH:O	2.18	0.43
1:B:39:ILE:HD11	1:B:62:SER:O	2.18	0.43
1:A:280:GLY:HA3	1:B:307:TYR:CD2	2.53	0.43
1:B:194:THR:HG22	2:B:621:HOH:O	2.19	0.43
1:B:89:ALA:HA	1:B:121:ILE:HG21	2.00	0.43
1:A:82:GLY:O	1:A:86:GLU:HG2	2.18	0.43
1:B:167:HIS:HB3	2:B:393:HOH:O	2.18	0.43
1:A:227:PHE:CZ	1:A:229:TYR:HB2	2.54	0.43
1:B:19:ILE:CD1	1:B:19:ILE:N	2.81	0.43
1:A:210:HIS:HD2	2:A:342:HOH:O	2.01	0.43
1:A:248:ARG:C	2:A:723:HOH:O	2.56	0.43
1:A:89:ALA:HA	1:A:121:ILE:HG21	2.01	0.43
1:A:334:LYS:CD	1:A:334:LYS:H	2.23	0.42
1:A:170:MET:HE1	1:A:203:THR:CG2	2.44	0.42
1:A:278:ASN:ND2	1:B:167:HIS:HE1	2.17	0.42
1:A:281:HIS:HE1	1:B:312:GLU:OE2	2.02	0.42
1:B:166:CYS:HG	1:B:192:TYR:HE2	1.63	0.42
1:B:141:GLU:OE2	1:B:192:TYR:OH	2.32	0.42
1:A:272:LEU:HD22	1:B:137:TRP:CD2	2.55	0.41
1:A:170:MET:HE2	1:B:209:TRP:NE1	2.25	0.41
1:A:138:ASN:HD21	1:B:283:GLY:HA2	1.85	0.41
1:B:137:TRP:HD1	1:B:167:HIS:HB3	1.77	0.41
1:A:166:CYS:SG	1:A:167:HIS:CB	3.09	0.41
1:B:18:ALA:HB1	1:B:218:VAL:HB	2.02	0.41
1:B:19:ILE:HD11	1:B:56:ILE:HG23	2.03	0.40
1:A:138:ASN:HD21	1:B:283:GLY:CA	2.34	0.40
1:A:194:THR:C	1:A:196:VAL:H	2.24	0.40
1:A:18:ALA:HB1	1:A:218:VAL:HB	2.03	0.40
1:B:170:MET:HE2	1:B:170:MET:HB3	1.80	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/334 (96%)	305 (95%)	15 (5%)	0	100	100
1	B	320/334 (96%)	301 (94%)	17 (5%)	2 (1%)	30	12
All	All	640/668 (96%)	606 (95%)	32 (5%)	2 (0%)	46	25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	137	TRP
1	B	192	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/277 (97%)	264 (98%)	4 (2%)	72	55
1	B	268/277 (97%)	256 (96%)	12 (4%)	34	11
All	All	536/554 (97%)	520 (97%)	16 (3%)	48	22

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	TRP
1	A	166	CYS
1	A	298	TYR
1	A	334	LYS
1	B	19	ILE
1	B	66	LEU
1	B	71	TRP
1	B	111	ASN
1	B	137	TRP
1	B	167	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	174	LEU
1	B	192	TYR
1	B	193	SER
1	B	196	VAL
1	B	298	TYR
1	B	334	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	138	ASN
1	A	167	HIS
1	A	210	HIS
1	A	275	ASN
1	A	278	ASN
1	A	292	HIS
1	B	35	ASN
1	B	111	ASN
1	B	167	HIS
1	B	210	HIS
1	B	275	ASN
1	B	278	ASN
1	B	281	HIS
1	B	292	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	322/334 (96%)	0.30	27 (8%) 14 16	9, 16, 31, 58	0
1	B	322/334 (96%)	0.97	41 (12%) 5 7	10, 21, 62, 98	0
All	All	644/668 (96%)	0.63	68 (10%) 8 10	9, 18, 44, 98	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	323	TYR	15.4
1	B	332	PHE	14.7
1	B	137	TRP	14.7
1	B	326	PRO	13.0
1	B	320	GLY	12.5
1	B	322	ILE	12.3
1	A	285	VAL	12.3
1	B	196	VAL	12.3
1	B	325	TYR	12.0
1	B	328	THR	11.5
1	A	284	TYR	10.8
1	B	327	THR	10.8
1	B	333	GLY	10.1
1	B	324	GLY	9.9
1	A	286	ALA	9.8
1	B	192	TYR	8.7
1	A	290	GLY	8.7
1	A	288	PRO	8.6
1	B	330	GLY	7.7
1	B	195	GLN	7.7
1	A	292	HIS	7.6
1	B	140	ILE	7.6
1	A	289	GLY	7.4
1	B	329	GLY	7.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	331	ARG	7.4
1	B	334	LYS	7.3
1	B	319	ASP	7.1
1	B	317	ILE	7.1
1	A	283	GLY	7.1
1	B	321	SER	7.0
1	A	291	GLU	6.6
1	B	111	ASN	5.6
1	B	139	PRO	5.2
1	A	195	GLN	5.0
1	B	69	ALA	4.8
1	A	287	LYS	4.8
1	B	318	LYS	4.6
1	B	34	HIS	4.5
1	B	314	HIS	4.3
1	A	323	TYR	4.2
1	B	110	SER	4.2
1	A	294	ALA	4.2
1	B	194	THR	4.1
1	B	193	SER	3.9
1	A	293	ASP	3.6
1	A	192	TYR	3.6
1	B	316	LYS	3.5
1	B	109	ASP	3.3
1	B	247	HIS	3.1
1	A	322	ILE	3.0
1	B	30	LYS	3.0
1	A	247	HIS	2.7
1	A	279	LEU	2.7
1	B	167	HIS	2.7
1	A	295	GLY	2.7
1	A	196	VAL	2.6
1	A	314	HIS	2.6
1	A	248	ARG	2.6
1	B	112	LYS	2.6
1	A	313	ASP	2.3
1	B	225	ASN	2.3
1	B	37	GLU	2.3
1	B	306	LYS	2.3
1	A	223	TYR	2.2
1	A	272	LEU	2.2
1	B	223	TYR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	111	ASN	2.0
1	A	34	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.