



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:15 AM GMT

PDB ID : 3DYL  
Title : human phosphodiesterase 9 substrate complex (ES complex)  
Authors : Liu, S.; Mansour, M.N.; Dillman, K.; Perez, J.; Danley, D.; Menniti, F.  
Deposited on : 2008-07-28  
Resolution : 2.70 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	
1	B	329	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	FMT	B	1	-	-	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5635 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called human phosphodiesterase 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	328	Total	C	N	O	S	0	0	0
			2706	1732	449	494	31			
1	B	328	Total	C	N	O	S	0	0	0
			2706	1732	449	494	31			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	178	GLY	-	EXPRESSION TAG	UNP O76083
A	179	SER	-	EXPRESSION TAG	UNP O76083
A	180	HIS	-	EXPRESSION TAG	UNP O76083
A	181	MET	-	EXPRESSION TAG	UNP O76083
B	178	GLY	-	EXPRESSION TAG	UNP O76083
B	179	SER	-	EXPRESSION TAG	UNP O76083
B	180	HIS	-	EXPRESSION TAG	UNP O76083
B	181	MET	-	EXPRESSION TAG	UNP O76083


- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mn	0	0
			1	1		
2	A	1	Total	Mn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is CYCLIC GUANOSINE MONOPHOSPHATE (three-letter code: PCG) (formula:  $C_{10}H_{12}N_5O_7P$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 5 is 3-ISOBUTYL-1-METHYLNANTHINE (three-letter code: IBM) (formula:  $C_{10}H_{14}N_4O_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			16	10	4	2		

- Molecule 6 is FORMIC ACID (three-letter code: FMT) (formula:  $\text{CH}_2\text{O}_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			3	1	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	96	Total	O	0	0
			96	96		
7	B	81	Total	O	0	0
			81	81		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: human phosphodiesterase 9



- Molecule 1: human phosphodiesterase 9



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	103.79 Å   103.79 Å   269.80 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	96.67 – 2.70 49.66 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (96.67-2.70) 99.7 (49.66-2.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.05 (at 2.69 Å)	Xtriage
Refinement program	REFMAC 5.3.0008	Depositor
R, $R_{free}$	0.188   ,   0.219 0.192   ,   0.217	Depositor DCC
$R_{free}$ test set	4208 reflections (11.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	46.7	Xtriage
Anisotropy	0.504	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 46.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Outliers	1 of 41463 reflections (0.002%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5635	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MN, FMT, IBM, PCG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.16	10/2776 (0.4%)	0.98	9/3759 (0.2%)
1	B	1.11	8/2776 (0.3%)	0.95	7/3759 (0.2%)
All	All	1.13	18/5552 (0.3%)	0.96	16/7518 (0.2%)

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	326	CYS	CB-SG	-11.25	1.63	1.82
1	A	411	GLU	CG-CD	9.32	1.66	1.51
1	B	370	GLU	CG-CD	8.95	1.65	1.51
1	B	411	GLU	CG-CD	8.39	1.64	1.51
1	A	370	GLU	CG-CD	8.04	1.64	1.51
1	A	291	CYS	CB-SG	-7.77	1.69	1.82
1	B	354	GLN	CG-CD	7.23	1.67	1.51
1	A	241	CYS	CB-SG	-6.65	1.71	1.82
1	A	400	CYS	CB-SG	-6.61	1.71	1.82
1	B	370	GLU	CB-CG	6.55	1.64	1.52
1	A	411	GLU	CD-OE1	6.23	1.32	1.25
1	A	370	GLU	CB-CG	6.11	1.63	1.52
1	B	419	CYS	CB-SG	-6.05	1.72	1.82
1	A	419	CYS	CB-SG	-6.01	1.72	1.82
1	B	231	ASN	CB-CG	5.97	1.64	1.51
1	B	310	GLU	CD-OE2	5.28	1.31	1.25
1	A	501	LYS	CD-CE	5.03	1.63	1.51
1	B	257	CYS	CB-SG	-5.00	1.73	1.81

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	255	ARG	NE-CZ-NH1	8.19	124.40	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	495	ARG	NE-CZ-NH1	7.43	124.01	120.30
1	A	426	MET	CG-SD-CE	7.40	112.04	100.20
1	A	402	ASP	CB-CG-OD1	6.81	124.42	118.30
1	A	237	ARG	NE-CZ-NH2	-6.61	117.00	120.30
1	A	379	MET	CG-SD-CE	-6.45	89.87	100.20
1	A	498	ASP	CB-CG-OD1	6.26	123.94	118.30
1	B	379	MET	CG-SD-CE	-6.15	90.36	100.20
1	B	429	ASP	CB-CG-OD2	-5.89	112.99	118.30
1	A	429	ASP	CB-CG-OD2	-5.82	113.06	118.30
1	B	241	CYS	CA-CB-SG	-5.47	104.15	114.00
1	B	429	ASP	CB-CG-OD1	5.33	123.10	118.30
1	A	255	ARG	NE-CZ-NH2	-5.33	117.64	120.30
1	A	295	ASP	CB-CG-OD1	5.23	123.01	118.30
1	B	314	ARG	NE-CZ-NH1	5.08	122.84	120.30
1	B	495	ARG	NE-CZ-NH2	-5.01	117.80	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2706	0	2643	45	0
1	B	2706	0	2643	31	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	23	0	11	2	0
5	B	16	0	14	0	0
6	B	3	0	1	1	0
7	A	96	0	0	15	0
7	B	81	0	0	3	0
All	All	5635	0	5312	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 7.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:184:PRO:HB2	7:A:143:HOH:O	1.51	1.10
1:A:209:ASN:HB3	7:A:140:HOH:O	1.53	1.06
1:A:448:THR:HG22	1:A:451:THR:H	1.24	1.02
1:B:375:PHE:CZ	1:B:379:MET:HE1	1.94	1.01
1:A:448:THR:HG21	7:A:125:HOH:O	1.71	0.89
1:A:375:PHE:CZ	1:A:379:MET:HE1	2.12	0.84
1:A:210:GLU:HG3	7:A:155:HOH:O	1.78	0.82
1:B:375:PHE:CE1	1:B:379:MET:HE1	2.15	0.81
1:A:277:PHE:CE1	1:A:390:MET:HE3	2.17	0.80
1:A:375:PHE:CE1	1:A:379:MET:HE1	2.16	0.80
1:B:277:PHE:CE1	1:B:390:MET:CE	2.71	0.73
1:A:300:ASN:C	1:A:300:ASN:HD22	1.93	0.71
1:B:277:PHE:CE1	1:B:390:MET:HE3	2.26	0.70
1:A:277:PHE:CE1	1:A:390:MET:CE	2.74	0.70
4:A:900:PCG:PA	7:A:1000:HOH:O	2.49	0.70
1:A:178:GLY:HA2	7:A:160:HOH:O	1.91	0.70
1:A:474:MET:HE2	7:A:154:HOH:O	1.94	0.66
1:A:448:THR:HG22	1:A:451:THR:N	2.04	0.64
1:A:375:PHE:CZ	1:A:379:MET:CE	2.81	0.63
1:B:375:PHE:CZ	1:B:379:MET:CE	2.76	0.63
1:B:277:PHE:CE1	1:B:390:MET:HE1	2.33	0.62
1:A:403:ILE:O	1:A:461:LEU:HD21	2.02	0.60
1:A:414:GLU:HG3	1:A:449:LYS:HZ1	1.67	0.60
1:B:443:ASP:O	1:B:447:VAL:HG23	2.03	0.59
1:A:403:ILE:HD11	1:A:464:MET:CE	2.35	0.57
1:A:186:TYR:N	7:A:143:HOH:O	2.36	0.57
1:A:442:MET:HE1	7:A:165:HOH:O	2.05	0.57
1:B:375:PHE:CE1	1:B:379:MET:CE	2.87	0.57
1:A:403:ILE:HD11	1:A:464:MET:HE1	1.88	0.56
1:B:379:MET:O	7:B:108:HOH:O	2.18	0.56
1:B:497:ASP:HB3	1:B:501:LYS:NZ	2.21	0.55
1:B:403:ILE:HD11	1:B:464:MET:CE	2.37	0.55
1:A:400:CYS:O	1:A:404:SER:HB3	2.07	0.55
1:B:450:ALA:O	1:B:454:ILE:HG13	2.08	0.54
1:A:186:TYR:CD2	7:A:143:HOH:O	2.52	0.54
1:B:454:ILE:HD13	1:B:491:GLU:HG2	1.90	0.53
1:A:375:PHE:CE1	1:A:379:MET:CE	2.91	0.53
1:A:231:ASN:ND2	1:A:234:THR:OG1	2.40	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:209:ASN:ND2	7:A:155:HOH:O	2.42	0.52
1:B:375:PHE:CD2	1:B:396:ILE:HG13	2.43	0.52
1:B:403:ILE:HD11	1:B:464:MET:HE2	1.92	0.52
1:A:316:ASN:O	1:B:357:ILE:HD13	2.09	0.52
1:A:386:ASN:HD22	1:A:386:ASN:C	2.13	0.52
6:B:1:FMT:C	7:B:33:HOH:O	2.59	0.49
4:A:900:PCG:H8	7:A:131:HOH:O	2.10	0.49
1:B:458:LYS:HE3	1:B:491:GLU:OE1	2.13	0.48
1:A:300:ASN:ND2	1:A:303:TYR:H	2.10	0.48
1:A:300:ASN:O	1:A:304:GLN:HG2	2.13	0.47
1:B:457:ILE:HA	1:B:461:LEU:HB2	1.96	0.47
1:A:386:ASN:HD21	1:A:388:GLU:HB2	1.78	0.47
1:A:372:MET:O	1:A:376:LYS:HG3	2.14	0.47
1:B:403:ILE:HD12	1:B:461:LEU:HD23	1.97	0.47
1:A:277:PHE:CE1	1:A:390:MET:HE1	2.48	0.46
1:B:267:MET:HE3	1:B:479:MET:SD	2.55	0.46
1:A:476:GLU:HG3	1:A:480:LEU:HD23	1.98	0.46
1:A:203:VAL:C	7:A:123:HOH:O	2.55	0.45
1:A:448:THR:HG23	7:A:132:HOH:O	2.17	0.45
1:B:497:ASP:HB3	1:B:501:LYS:HZ2	1.81	0.44
1:A:186:TYR:CG	7:A:143:HOH:O	2.70	0.44
1:B:429:ASP:OD2	1:B:444:ARG:NH1	2.50	0.44
1:A:260:VAL:HG22	1:A:402:ASP:HA	1.99	0.44
1:B:271:CYS:HB3	1:B:472:PHE:CD2	2.52	0.44
1:A:252:HIS:ND1	1:A:423:GLU:OE2	2.34	0.44
1:B:400:CYS:O	1:B:404:SER:HB3	2.18	0.44
1:B:277:PHE:CD1	1:B:390:MET:CE	3.01	0.43
1:A:304:GLN:NE2	1:A:309:THR:HG21	2.34	0.43
1:A:404:SER:O	1:A:405:ASN:C	2.56	0.43
1:A:288:ALA:O	1:A:292:HIS:HB3	2.19	0.42
1:A:277:PHE:CD1	1:A:390:MET:CE	3.02	0.42
1:B:318:ILE:O	1:B:319:SER:C	2.58	0.42
1:A:300:ASN:ND2	1:A:302:THR:N	2.67	0.42
1:B:461:LEU:HD13	1:B:483:LEU:HD21	2.01	0.42
1:B:322:GLU:HA	1:B:325:HIS:CD2	2.54	0.42
1:B:440:PRO:HD2	7:B:134:HOH:O	2.20	0.42
1:B:319:SER:N	1:B:320:PRO:HD3	2.36	0.41
1:A:300:ASN:HD21	1:A:302:THR:HB	1.86	0.41
1:A:302:THR:HG22	1:A:306:ASN:ND2	2.36	0.41
1:B:379:MET:HE3	1:B:379:MET:HB2	1.79	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/329 (99%)	316 (97%)	8 (2%)	2 (1%)	30	59
1	B	326/329 (99%)	316 (97%)	9 (3%)	1 (0%)	46	75
All	All	652/658 (99%)	632 (97%)	17 (3%)	3 (0%)	34	63

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	444	ARG
1	A	184	PRO
1	B	184	PRO

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	304/305 (100%)	289 (95%)	15 (5%)	31	61
1	B	304/305 (100%)	300 (99%)	4 (1%)	76	92
All	All	608/610 (100%)	589 (97%)	19 (3%)	47	78

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	181	MET
1	A	257	CYS
1	A	267	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	300	ASN
1	A	326	CYS
1	A	386	ASN
1	A	444	ARG
1	A	448	THR
1	A	449	LYS
1	A	471	LEU
1	A	474	MET
1	A	480	LEU
1	A	496	ILE
1	A	500	MET
1	A	505	LYS
1	B	181	MET
1	B	257	CYS
1	B	474	MET
1	B	493	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	231	ASN
1	A	300	ASN
1	A	304	GLN
1	A	386	ASN
1	A	389	HIS
1	B	304	GLN
1	B	351	GLN
1	B	389	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PCG	A	900	3,2	20,26,26	5.12	1 (5%)	21,41,41	2.43	7 (33%)
6	FMT	B	1	2	0,2,2	0.00	-	0,1,1	0.00	-
5	IBM	B	900	-	9,17,17	1.18	1 (11%)	9,25,25	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PCG	A	900	3,2	-	0/0/31/31	0/4/4/4
6	FMT	B	1	2	-	0/0/0/0	0/0/0/0
5	IBM	B	900	-	-	0/4/4/4	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	900	IBM	C5-C4	2.09	1.45	1.40
4	A	900	PCG	PA-O3'	22.53	1.96	1.58

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	900	PCG	O5'-PA-O3'	-4.49	99.16	105.75
4	A	900	PCG	C2'-C1'-N9	-3.28	109.29	114.29
4	A	900	PCG	C5-C6-N1	-3.03	119.44	123.59
4	A	900	PCG	N3-C2-N1	-2.10	124.25	127.44
4	A	900	PCG	C6-N1-C2	2.93	120.00	115.94
4	A	900	PCG	O5'-C5'-C4'	4.16	116.80	105.78
4	A	900	PCG	O1A-PA-O2A	5.91	127.64	108.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	900	PCG	2	0
6	B	1	FMT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	328/329 (99%)	0.16	9 (2%) 58 58	39, 49, 62, 85	0
1	B	328/329 (99%)	0.11	6 (1%) 71 72	41, 49, 62, 85	0
All	All	656/658 (99%)	0.14	15 (2%) 64 64	39, 49, 62, 85	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	179	SER	6.2
1	A	180	HIS	5.7
1	A	178	GLY	5.0
1	B	505	LYS	4.6
1	B	178	GLY	3.9
1	B	179	SER	3.8
1	A	181	MET	2.9
1	A	432	LYS	2.9
1	B	180	HIS	2.8
1	A	186	TYR	2.8
1	A	182	THR	2.8
1	B	181	MET	2.5
1	B	495	ARG	2.2
1	A	185	LYS	2.2
1	A	183	TYR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	FMT	B	1	3/3	0.88	0.21	2.36	69,69,70,71	0
5	IBM	B	900	16/16	0.94	0.17	-0.49	56,58,62,63	0
4	PCG	A	900	23/23	0.97	0.13	-1.30	47,48,59,66	0
3	MG	B	902	1/1	0.95	0.09	-2.96	46,46,46,46	0
2	MN	B	901	1/1	0.99	0.09	-4.54	57,57,57,57	0
3	MG	A	902	1/1	0.94	0.08	-5.08	39,39,39,39	0
2	MN	A	901	1/1	0.99	0.05	-8.37	57,57,57,57	0

## 6.5 Other polymers

There are no such residues in this entry.