



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:58 PM GMT

PDB ID : 4E0U
Title : Crystal structure of CdpNPT in complex with thiolodiphosphate and (S)-benzodiazependione
Authors : Schuller, J.M.; Zocher, G.; Stehle, T.
Deposited on : 2012-03-05
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	428	 4% 85% 10% • 5%
1	B	428	 4% 87% 8% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	A	504	-	-	-	X
5	EDO	B	503	-	-	-	X
5	EDO	B	505	-	-	-	X
6	NA	A	505	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6623 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclic dipeptide N-prenyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	408	Total	C	N	O	S	0	0	0
			3242	2090	539	595	18			
1	B	408	Total	C	N	O	S	0	0	0
			3243	2091	540	594	18			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	13	HIS	-	EXPRESSION TAG	UNP D1D8L6
A	14	HIS	-	EXPRESSION TAG	UNP D1D8L6
A	15	HIS	-	EXPRESSION TAG	UNP D1D8L6
A	16	HIS	-	EXPRESSION TAG	UNP D1D8L6
A	17	HIS	-	EXPRESSION TAG	UNP D1D8L6
A	18	HIS	-	EXPRESSION TAG	UNP D1D8L6
A	19	SER	-	EXPRESSION TAG	UNP D1D8L6
A	20	SER	-	EXPRESSION TAG	UNP D1D8L6
A	21	GLY	-	EXPRESSION TAG	UNP D1D8L6
A	22	LEU	-	EXPRESSION TAG	UNP D1D8L6
A	23	VAL	-	EXPRESSION TAG	UNP D1D8L6
A	24	PRO	-	EXPRESSION TAG	UNP D1D8L6
A	25	ARG	-	EXPRESSION TAG	UNP D1D8L6
A	26	GLY	-	EXPRESSION TAG	UNP D1D8L6
A	27	SER	-	EXPRESSION TAG	UNP D1D8L6
A	28	HIS	-	EXPRESSION TAG	UNP D1D8L6
B	13	HIS	-	EXPRESSION TAG	UNP D1D8L6
B	14	HIS	-	EXPRESSION TAG	UNP D1D8L6
B	15	HIS	-	EXPRESSION TAG	UNP D1D8L6
B	16	HIS	-	EXPRESSION TAG	UNP D1D8L6
B	17	HIS	-	EXPRESSION TAG	UNP D1D8L6
B	18	HIS	-	EXPRESSION TAG	UNP D1D8L6
B	19	SER	-	EXPRESSION TAG	UNP D1D8L6
B	20	SER	-	EXPRESSION TAG	UNP D1D8L6
B	21	GLY	-	EXPRESSION TAG	UNP D1D8L6

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	22	LEU	-	EXPRESSION TAG	UNP D1D8L6
B	23	VAL	-	EXPRESSION TAG	UNP D1D8L6
B	24	PRO	-	EXPRESSION TAG	UNP D1D8L6
B	25	ARG	-	EXPRESSION TAG	UNP D1D8L6
B	26	GLY	-	EXPRESSION TAG	UNP D1D8L6
B	27	SER	-	EXPRESSION TAG	UNP D1D8L6
B	28	HIS	-	EXPRESSION TAG	UNP D1D8L6

- Molecule 2 is (3S)-3-(1H-INDOL-3-YLMETHYL)-3,4-DIHYDRO-1H-1,4-BENZODIAZEPINE-2,5-DIONE (three-letter code: 0MV) (formula: C₁₈H₁₅N₃O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			23	18	3	2		
2	B	1	Total	C	N	O	0	0
			23	18	3	2		

- Molecule 3 is TRIHYDROGEN THIODIPHOSPHATE (three-letter code: PIS) (formula: H₃O₆P₂S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	O	P	S	0	0
			9	6	2	1		
3	B	1	Total	O	P	S	0	0
			9	6	2	1		

- Molecule 4 is IMIDAZOLE (three-letter code: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			5	3	2		
4	B	1	Total	C	N	0	0
			5	3	2		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Na	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	22	Total	O	0	0
			22	22		
7	B	25	Total	O	0	0
			25	25		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cyclic dipeptide N-prenyltransferase



- Molecule 1: Cyclic dipeptide N-prenyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	137.73Å 137.73Å 170.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.83 – 2.60 29.83 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.83-2.60) 98.1 (29.83-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.196 , 0.229 0.193 , 0.223	Depositor DCC
R_{free} test set	2502 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	67.9	Xtriage
Anisotropy	0.349	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 42.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 50027 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6623	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, 0MV, PIS, IMD, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/3327	0.59	0/4522
1	B	0.52	0/3328	0.62	0/4523
All	All	0.52	0/6655	0.60	0/9045

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3242	0	3165	27	0
1	B	3243	0	3169	21	0
2	A	23	0	15	0	0
2	B	23	0	15	0	0
3	A	9	0	0	2	0
3	B	9	0	0	0	0
4	A	5	0	5	0	0
4	B	5	0	5	0	0
5	A	8	0	12	0	0
5	B	8	0	12	1	0
6	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	22	0	0	0	0
7	B	25	0	0	0	0
All	All	6623	0	6398	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68:MET:HE2	1:A:128:THR:HG23	1.62	0.81
1:A:288:TYR:OH	3:A:502:PIS:S1	2.47	0.71
1:B:309:LEU:HD12	1:B:309:LEU:O	1.93	0.68
1:B:300:GLU:HG2	1:B:318:LEU:HD11	1.78	0.66
1:B:141:VAL:HG13	1:B:141:VAL:O	1.97	0.65
1:A:432:LEU:N	1:A:432:LEU:HD12	2.13	0.63
1:B:351:ASN:HD22	1:B:351:ASN:C	2.05	0.60
1:A:148:ARG:NH2	1:A:184:GLU:OE1	2.33	0.60
1:A:300:GLU:HG2	1:A:318:LEU:HD11	1.86	0.57
1:B:351:ASN:ND2	1:B:351:ASN:C	2.58	0.56
1:B:141:VAL:O	1:B:141:VAL:CG1	2.56	0.52
1:A:179:LEU:HD12	1:A:183:ASP:HB3	1.91	0.52
1:A:305:LEU:HD21	1:A:352:TYR:CE1	2.45	0.52
1:B:234:THR:HG22	1:B:236:SER:N	2.26	0.51
1:A:144:ASP:O	1:A:145:ARG:CB	2.60	0.49
1:B:48:LEU:HD12	5:B:505:EDO:H22	1.93	0.49
1:A:160:LEU:HD22	1:B:138:GLN:HG2	1.95	0.49
1:B:89:LEU:HD21	1:B:158:LEU:HD13	1.95	0.48
1:B:351:ASN:HD22	1:B:352:TYR:N	2.11	0.48
1:B:273:LEU:HD12	1:B:286:LYS:O	2.15	0.47
1:A:376:LYS:O	1:A:380:ILE:HD13	2.14	0.47
1:A:179:LEU:HD12	1:A:183:ASP:CB	2.45	0.47
1:A:187:LEU:HD13	1:A:230:ALA:HB2	1.97	0.46
1:A:141:VAL:O	1:A:141:VAL:HG13	2.16	0.45
1:A:68:MET:HE3	1:A:119:GLN:HB2	1.97	0.45
1:B:327:GLU:HB3	1:B:380:ILE:HD12	1.98	0.45
1:A:328:LEU:HB2	1:A:330:ILE:HD12	1.99	0.45
1:B:192:LEU:HG	1:B:196:LEU:HD13	1.99	0.44
1:A:305:LEU:HD12	1:A:309:LEU:HD13	1.99	0.44
1:A:68:MET:CE	1:A:128:THR:HG23	2.42	0.43
1:B:369:LEU:HD11	1:B:420:LEU:HB2	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:372:ILE:HA	1:A:373:PRO:HD3	1.79	0.43
1:A:188:ASN:HD22	1:A:188:ASN:H	1.67	0.43
1:B:330:ILE:HD11	1:B:367:MET:HE2	2.00	0.43
1:B:234:THR:CG2	1:B:236:SER:HB3	2.49	0.43
1:A:402:GLN:HA	1:A:410:LEU:HD11	2.01	0.43
1:A:196:LEU:O	1:A:200:GLN:N	2.48	0.43
1:B:170:HIS:ND1	1:B:220:GLU:OE2	2.52	0.42
1:B:358:GLN:NE2	1:B:360:LEU:O	2.37	0.42
1:B:330:ILE:CD1	1:B:367:MET:HE2	2.50	0.41
1:A:262:HIS:HE1	1:A:302:HIS:ND1	2.18	0.41
1:A:255:LEU:HD12	1:A:255:LEU:H	1.85	0.41
1:B:234:THR:HG22	1:B:237:ASN:H	1.85	0.41
1:A:68:MET:HE2	1:A:128:THR:CG2	2.43	0.41
1:A:430:PRO:HB2	1:A:432:LEU:HD11	2.01	0.41
1:A:366:TYR:OH	3:A:502:PIS:S1	2.69	0.40
1:A:194:LYS:O	1:A:198:ASN:ND2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	406/428 (95%)	389 (96%)	15 (4%)	2 (0%)	34	60
1	B	406/428 (95%)	397 (98%)	8 (2%)	1 (0%)	52	77
All	All	812/856 (95%)	786 (97%)	23 (3%)	3 (0%)	39	65

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	145	ARG
1	B	144	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	334	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/367 (93%)	332 (97%)	9 (3%)	54	80
1	B	341/367 (93%)	331 (97%)	10 (3%)	50	77
All	All	682/734 (93%)	663 (97%)	19 (3%)	51	78

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	129	ARG
1	A	141	VAL
1	A	188	ASN
1	A	252	LEU
1	A	337	PRO
1	A	359	ARG
1	A	386	GLN
1	A	427	GLU
1	A	432	LEU
1	B	73	LYS
1	B	129	ARG
1	B	141	VAL
1	B	174	SER
1	B	187	LEU
1	B	200	GLN
1	B	252	LEU
1	B	324	LEU
1	B	342	GLU
1	B	351	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	188	ASN
1	A	233	GLN
1	A	237	ASN
1	A	262	HIS
1	B	60	HIS
1	B	237	ASN
1	B	264	GLN
1	B	351	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	0MV	A	501	-	25,26,26	3.74	9 (36%)	21,37,37	1.66	5 (23%)
3	PIS	A	502	-	4,8,8	8.62	2 (50%)	9,13,13	1.48	1 (11%)
4	IMD	A	503	-	3,5,5	0.43	0	4,5,5	0.60	0
5	EDO	A	504	-	3,3,3	0.59	0	2,2,2	0.19	0
5	EDO	A	506	-	3,3,3	0.41	0	2,2,2	0.48	0
2	0MV	B	501	-	25,26,26	3.48	9 (36%)	21,37,37	1.59	5 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PIS	B	502	-	4,8,8	7.67	1 (25%)	9,13,13	1.58	1 (11%)
5	EDO	B	503	-	3,3,3	0.53	0	2,2,2	0.48	0
4	IMD	B	504	-	3,5,5	0.47	0	4,5,5	0.64	0
5	EDO	B	505	-	3,3,3	0.47	0	2,2,2	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	0MV	A	501	-	-	0/3/20/20	0/4/4/4
3	PIS	A	502	-	-	0/5/6/6	0/0/0/0
4	IMD	A	503	-	-	0/0/0/0	0/1/1/1
5	EDO	A	504	-	-	0/1/1/1	0/0/0/0
5	EDO	A	506	-	-	0/1/1/1	0/0/0/0
2	0MV	B	501	-	-	0/3/20/20	0/4/4/4
3	PIS	B	502	-	-	0/5/6/6	0/0/0/0
5	EDO	B	503	-	-	0/1/1/1	0/0/0/0
4	IMD	B	504	-	-	0/0/0/0	0/1/1/1
5	EDO	B	505	-	-	0/1/1/1	0/0/0/0

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	PIS	P2-S1	-16.99	1.64	1.96
3	B	502	PIS	P2-S1	-15.14	1.67	1.96
2	A	501	0MV	CAT-CAP	-10.02	1.39	1.49
2	B	501	0MV	CAT-CAP	-9.36	1.40	1.49
2	A	501	0MV	CA-C	-8.25	1.41	1.52
2	B	501	0MV	CA-C	-7.65	1.42	1.52
2	A	501	0MV	CZ2-CE2	-4.35	1.33	1.41
2	B	501	0MV	CZ2-CE2	-4.17	1.34	1.41
2	A	501	0MV	CE3-CD2	-4.07	1.34	1.42
2	B	501	0MV	CE3-CD2	-3.38	1.35	1.42
2	B	501	0MV	CD1-CG	-2.91	1.33	1.38
2	B	501	0MV	CD2-CE2	-2.73	1.35	1.42
2	A	501	0MV	CD2-CE2	-2.62	1.35	1.42
2	A	501	0MV	CD1-CG	-2.33	1.34	1.38
3	A	502	PIS	P1-O2	2.20	1.62	1.54
2	B	501	0MV	CA-N	3.30	1.49	1.46
2	B	501	0MV	C-NAM	4.20	1.40	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	0MV	CA-N	4.26	1.50	1.46
2	A	501	0MV	C-NAM	4.84	1.41	1.35
2	B	501	0MV	CAP-N	8.40	1.40	1.34
2	A	501	0MV	CAP-N	8.95	1.41	1.34

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	PIS	P1-O1-P2	-4.12	118.86	132.67
3	A	502	PIS	P1-O1-P2	-4.07	119.02	132.67
2	B	501	0MV	CB-CG-CD1	-3.47	123.68	127.97
2	A	501	0MV	CB-CG-CD1	-3.19	124.03	127.97
2	A	501	0MV	CAG-CAS-NAM	-3.09	112.11	117.93
2	B	501	0MV	CAG-CAS-NAM	-2.83	112.60	117.93
2	B	501	0MV	O-C-NAM	-2.53	116.52	121.39
2	A	501	0MV	O-C-NAM	-2.33	116.91	121.39
2	B	501	0MV	CAT-CAS-NAM	2.35	126.98	123.48
2	A	501	0MV	CAT-CAS-NAM	2.53	127.25	123.48
2	B	501	0MV	CA-C-NAM	3.32	121.39	116.26
2	A	501	0MV	CA-C-NAM	3.91	122.30	116.26

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	PIS	2	0
5	B	505	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	408/428 (95%)	-0.07	16 (3%) 43 35	51, 79, 121, 142	0
1	B	408/428 (95%)	-0.19	18 (4%) 38 30	49, 72, 112, 134	0
All	All	816/856 (95%)	-0.13	34 (4%) 40 32	49, 74, 119, 142	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	335	LEU	4.7
1	B	33	PRO	4.5
1	A	440	GLU	4.5
1	B	193	THR	4.4
1	A	193	THR	4.4
1	B	192	LEU	4.3
1	A	334	PRO	4.1
1	B	335	LEU	3.9
1	A	339	ALA	3.9
1	A	197	THR	3.5
1	A	338	SER	3.3
1	B	357	GLY	3.0
1	A	336	GLU	3.0
1	B	197	THR	3.0
1	A	337	PRO	3.0
1	A	190	GLU	2.8
1	A	33	PRO	2.8
1	B	198	ASN	2.8
1	B	336	GLU	2.7
1	A	421	SER	2.4
1	B	191	GLN	2.4
1	B	196	LEU	2.4
1	B	338	SER	2.4
1	B	339	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	194	LYS	2.3
1	A	249	ASP	2.3
1	B	334	PRO	2.3
1	A	307	GLY	2.3
1	B	421	SER	2.2
1	A	356	ALA	2.2
1	A	200	GLN	2.1
1	B	342	GLU	2.1
1	B	433	SER	2.1
1	B	435	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	NA	A	505	1/1	0.77	0.76	12.80	57,57,57,57	0
5	EDO	B	503	4/4	0.89	0.48	6.68	88,88,89,90	0
5	EDO	A	504	4/4	0.81	0.39	5.61	78,78,79,79	0
5	EDO	B	505	4/4	0.94	0.26	4.54	90,90,91,91	0
5	EDO	A	506	4/4	0.95	0.16	1.40	84,84,84,85	0
2	0MV	A	501	23/23	0.89	0.18	-0.28	71,79,84,85	0
4	IMD	A	503	5/5	0.91	0.23	-0.79	77,77,78,79	0
2	0MV	B	501	23/23	0.96	0.12	-1.11	71,79,81,82	0
3	PIS	A	502	9/9	0.95	0.15	-1.20	80,83,85,87	0
3	PIS	B	502	9/9	0.96	0.13	-1.37	70,73,77,85	0
4	IMD	B	504	5/5	0.95	0.28	-	80,80,81,81	0

6.5 Other polymers ⓘ

There are no such residues in this entry.