



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:59 PM GMT

PDB ID : 4E0V
Title : Structure of L-amino acid oxidase from the B. jararacussu venom
Authors : Ullah, A.; Souza, T.A.C.B.; Betzel, C.; Murakami, M.T.; Arni, R.K.
Deposited on : 2012-03-05
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1114 (3.14-3.06)
Clashscore	102246	1222 (3.14-3.06)
Ramachandran outliers	100387	1174 (3.14-3.06)
Sidechain outliers	100360	1174 (3.14-3.06)
RSRZ outliers	91569	1119 (3.14-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	497	
1	B	497	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FAD	A	501	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7726 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-amino-acid oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	474	Total	C	N	O	S	0	0	0
			3771	2398	637	722	14			
1	B	480	Total	C	N	O	S	0	0	0
			3833	2432	653	734	14			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	7	Total 7	O 7	0	0
3	B	9	Total 9	O 9	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: L-amino-acid oxidase



• Molecule 1: L-amino-acid oxidase



F433	A371	P307	P244	K178	R116
T434	I372	L308	T245	K179	E117
P435	G373	P309	S246	Y180	R118
Y436	I374	P310	N247	D181	A119
Q437	G375	K311	Y248	T182	W120
F438	D376	K312	Q249	Y183	Y121
Q439	D377	A313	K250		F122
H440	A378	H314	I251	K186	I123
F441	I379	A315	Q252	E187	K124
S442	Y380	L316	E253	Y188	L125
	F381		K254	L189	I126
	E382	V319	V255	L190	I127
L445	A383	H320		K191	K128
T446	I384	Y321	N258	E192	R129
A447	D385	R322	A259		V130
P448	D386	S323	R260	S196	G131
V449	F387	G324	Y261	P197	E132
D450	E387	T325	I262	G198	V133
R451	D388	K326	V263	A199	N134
I452		I327	I264	V200	K135
Y453	D391	F328	Q265	D201	D136
F454	I392	L329	Q266	M202	P137
A455	V393	T330	D267	I203	G138
G456	I394	G331	V268	G204	V139
E457	N395	T332	K269	D205	L140
Y458	D396	K333	E270	L206	D141
T459	I397	R334	V271	L207	Y142
A460	S398	F335	T272	N208	P143
Q461	L399	W336	T273	E209	V144
A462	I400	E337	T274	D210	K145
H463	H401	D338	Y275	S211	P146
G464	Q402	D339	Q276	G212	S147
I465	L403	G340	T277	Y213	E148
I466	P404	I341	S278	Y214	V149
A467	K405		E279	V215	G150
S468	E406	K345	K280	S216	K151
T469	E407	S346	E281	F217	S152
I470	I408	T347	T282	I218	A153
K471	Q409	T348	L283		
S472	A410	D349	S284	L221	L156
G473	I411	L350	V285	K222	Y157
P474	C412	P351	T286	H223	
E475	R413	S352	A287	D224	S160
G476		R353	D288	D225	L161
I477	H416	F354	Y289	I226	Q162
D478	I417	I355	V290		K163
V479	Q418	Y356	I291	Y229	A164
R480	R419	Y357	V292	E230	V165
R481	N420	P358	C293	K231	E166
A482	S421	N359	T294	R232	L167
	I422	H360	T295	F233	L168
SER	D423	N361	S296	D234	R169
GLU	K424		R297	E235	R170
	Y425	N364	A298	T171	T172
	A426	G365	A299	I236	C173
	N427	V366	R300	V237	
	G428	G367	R301		S174
	Q429	V368	I302	M240	Y175
	I430	I369	K303	D241	
	T431	I370		K242	M176
	T432			L243	

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.39Å 72.19Å 101.53Å 90.00° 90.90° 90.00°	Depositor
Resolution (Å)	29.02 – 3.10 28.79 – 3.10	Depositor EDS
% Data completeness (in resolution range)	95.1 (29.02-3.10) 94.7 (28.79-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 3.11Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, R_{free}	0.181 , 0.259 0.188 , 0.262	Depositor DCC
R_{free} test set	1721 reflections (10.87%)	DCC
Wilson B-factor (Å ²)	50.9	Xtriage
Anisotropy	0.016	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 73.3	EDS
Estimated twinning fraction	0.025 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 16752 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7726	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	1/3856 (0.0%)	0.80	0/5218
1	B	0.61	0/3920	0.76	0/5305
All	All	0.64	1/7776 (0.0%)	0.78	0/10523

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	88	PRO	N-CD	11.01	1.63	1.47

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	62	LEU	Peptide
1	A	63	GLU	Peptide
1	B	449	VAL	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3771	0	3678	591	1
1	B	3833	0	3747	655	1
2	A	53	0	31	29	0
2	B	53	0	31	18	0
3	A	7	0	0	0	0
3	B	9	0	0	1	0
All	All	7726	0	7487	1242	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 82.

The worst 5 of 1242 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:15:ASP:HB3	1:A:19:PHE:CE2	1.54	1.42
1:B:315:ALA:O	1:B:319:VAL:HG23	1.24	1.35
1:A:92:PRO:HG2	1:A:95:HIS:ND1	1.39	1.34
1:B:473:GLY:HA2	1:B:475:GLU:OE1	1.18	1.30
1:A:269:LYS:HE3	1:A:269:LYS:C	1.56	1.26

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:104:LYS:O	1:B:387:GLU:OE2[2_545]	2.06	0.14

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	470/497 (95%)	326 (69%)	96 (20%)	48 (10%)	1	4
1	B	478/497 (96%)	357 (75%)	78 (16%)	43 (9%)	1	5
All	All	948/994 (95%)	683 (72%)	174 (18%)	91 (10%)	1	4

5 of 91 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	8	GLU
1	A	51	VAL
1	A	53	ALA
1	A	96	ARG
1	A	278	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	403/427 (94%)	322 (80%)	81 (20%)	1	7
1	B	412/427 (96%)	336 (82%)	76 (18%)	2	9
All	All	815/854 (95%)	658 (81%)	157 (19%)	2	8

5 of 157 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	418	GLN
1	B	35	ARG
1	B	398	SER
1	A	439	GLN
1	A	479	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	418	GLN
1	B	95	HIS
1	B	418	GLN
1	B	77	ASN
1	A	265	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	501	-	48,58,58	1.26	5 (10%)	54,89,89	3.12	17 (31%)
2	FAD	B	501	-	48,58,58	1.23	5 (10%)	54,89,89	2.13	10 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	501	-	-	0/30/50/50	0/6/6/6
2	FAD	B	501	-	-	0/30/50/50	0/6/6/6

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	FAD	C6-C5X	-3.31	1.36	1.41
2	B	501	FAD	O4B-C4B	-2.03	1.40	1.45
2	B	501	FAD	C10-N1	2.17	1.39	1.35
2	A	501	FAD	C4-N3	2.46	1.37	1.33
2	A	501	FAD	C2A-N3A	2.73	1.37	1.32

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	FAD	N3A-C2A-N1A	-13.21	118.78	128.89
2	B	501	FAD	N3A-C2A-N1A	-11.79	119.87	128.89
2	A	501	FAD	C4B-O4B-C1B	-11.41	97.18	109.72
2	A	501	FAD	C2B-C1B-N9A	-5.89	105.30	114.29
2	A	501	FAD	C1B-N9A-C4A	-5.03	119.36	126.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 47 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	FAD	29	0
2	B	501	FAD	18	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	474/497 (95%)	-0.52	1 (0%) 95 91	18, 30, 44, 53	11 (2%)
1	B	480/497 (96%)	-0.48	2 (0%) 93 85	14, 31, 49, 63	16 (3%)
All	All	954/994 (95%)	-0.50	3 (0%) 94 88	14, 30, 45, 63	27 (2%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	113	SER	2.3
1	B	9	GLU	2.1
1	B	3	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FAD	A	501	53/53	0.94	0.18	0.13	8,19,39,40	0
2	FAD	B	501	53/53	0.95	0.16	-0.27	13,25,28,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.