



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 07:05 PM GMT

PDB ID : 1E1K
Title : ADRENODOXIN REDUCTASE IN COMPLEX WITH NADP⁺ OBTAINED
BY A SOAKING EXPERIMENT
Authors : Ziegler, G.A.; Schulz, G.E.
Deposited on : 2000-05-09
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	460	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAP	A	802	X	-	-	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3959 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADRENODOXIN REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	455	Total	C	N	O	S	0	1	0
			3512	2219	637	645	11			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	346	Total	O	0	0
			346	346		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ADRENODOXIN REDUCTASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.90Å 62.40Å 78.20Å 90.00° 106.60° 90.00°	Depositor
Resolution (Å)	20.00 – 1.95 25.78 – 1.95	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-1.95) 99.8 (25.78-1.95)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 1.95Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.184 , 0.227 0.184 , 0.225	Depositor DCC
R_{free} test set	2106 reflections (5.41%)	DCC
Wilson B-factor (Å ²)	18.5	Xtriage
Anisotropy	0.462	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 41100 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3959	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/3595	1.53	60/4885 (1.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (60) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	31	ARG	CD-NE-CZ	15.61	145.46	123.60
1	A	185	ARG	NE-CZ-NH2	-14.60	113.00	120.30
1	A	48	ARG	NE-CZ-NH1	12.39	126.50	120.30
1	A	31	ARG	NE-CZ-NH1	12.16	126.38	120.30
1	A	188	ARG	NE-CZ-NH1	12.03	126.31	120.30
1	A	162	ARG	NE-CZ-NH2	-11.95	114.33	120.30
1	A	278	ARG	NE-CZ-NH2	10.49	125.55	120.30
1	A	264	ARG	NE-CZ-NH2	-10.39	115.10	120.30
1	A	70	ARG	CD-NE-CZ	9.67	137.14	123.60
1	A	445	ARG	NE-CZ-NH2	-9.52	115.54	120.30
1	A	310	ARG	NE-CZ-NH2	-9.37	115.61	120.30
1	A	264	ARG	NE-CZ-NH1	9.32	124.96	120.30
1	A	417	ARG	NE-CZ-NH2	-9.19	115.71	120.30
1	A	188	ARG	CD-NE-CZ	9.03	136.24	123.60
1	A	242	ARG	NE-CZ-NH2	-8.92	115.84	120.30
1	A	445	ARG	NE-CZ-NH1	8.59	124.59	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	162	ARG	NE-CZ-NH1	8.19	124.39	120.30
1	A	437	ARG	NE-CZ-NH2	-8.13	116.23	120.30
1	A	278	ARG	NH1-CZ-NH2	-8.08	110.52	119.40
1	A	355	ARG	NE-CZ-NH2	-8.08	116.26	120.30
1	A	188	ARG	NE-CZ-NH2	-7.89	116.36	120.30
1	A	70	ARG	NE-CZ-NH1	7.63	124.12	120.30
1	A	370	ARG	NE-CZ-NH1	7.54	124.07	120.30
1	A	242	ARG	NE-CZ-NH1	7.49	124.05	120.30
1	A	251	ARG	NE-CZ-NH2	-7.30	116.65	120.30
1	A	278	ARG	NE-CZ-NH1	7.28	123.94	120.30
1	A	124[A]	ARG	NE-CZ-NH2	7.13	123.86	120.30
1	A	124[B]	ARG	NE-CZ-NH2	7.13	123.86	120.30
1	A	106	ASP	CB-CG-OD1	-6.82	112.17	118.30
1	A	456	ARG	NE-CZ-NH2	-6.78	116.91	120.30
1	A	31	ARG	NE-CZ-NH2	-6.72	116.94	120.30
1	A	343	ASP	CB-CG-OD2	-6.71	112.26	118.30
1	A	334	ARG	N-CA-CB	-6.21	99.42	110.60
1	A	185	ARG	NH1-CZ-NH2	6.14	126.16	119.40
1	A	370	ARG	NH1-CZ-NH2	-6.12	112.67	119.40
1	A	171	LEU	CB-CG-CD1	6.10	121.37	111.00
1	A	198	ARG	NE-CZ-NH2	6.08	123.34	120.30
1	A	48	ARG	NE-CZ-NH2	-6.03	117.28	120.30
1	A	240	ARG	NE-CZ-NH2	-5.87	117.36	120.30
1	A	19	PHE	CB-CG-CD2	5.86	124.90	120.80
1	A	319	ASP	CB-CG-OD1	5.86	123.58	118.30
1	A	106	ASP	CB-CG-OD2	5.85	123.57	118.30
1	A	370	ARG	NE-CZ-NH2	5.83	123.22	120.30
1	A	352	MET	CA-CB-CG	5.76	123.09	113.30
1	A	422	VAL	CA-CB-CG2	-5.76	102.26	110.90
1	A	409	PHE	CB-CG-CD2	5.59	124.71	120.80
1	A	206	THR	CA-CB-CG2	-5.52	104.67	112.40
1	A	256	LYS	CA-CB-CG	5.48	125.45	113.40
1	A	137	ARG	NE-CZ-NH1	5.47	123.03	120.30
1	A	318	GLU	OE1-CD-OE2	-5.40	116.82	123.30
1	A	137	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	A	20	TYR	CB-CG-CD2	-5.36	117.79	121.00
1	A	403	PRO	CA-C-N	5.23	128.70	117.20
1	A	19	PHE	CB-CG-CD1	-5.19	117.17	120.80
1	A	211	ARG	NE-CZ-NH2	-5.19	117.71	120.30
1	A	110	LEU	CA-CB-CG	5.18	127.21	115.30
1	A	403	PRO	N-CA-C	5.08	125.31	112.10
1	A	57	GLU	CA-CB-CG	5.07	124.56	113.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	415	ASP	CB-CG-OD2	5.06	122.86	118.30
1	A	171	LEU	CA-CB-CG	5.02	126.85	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	330	GLY	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3512	0	3544	42	0
2	A	53	0	31	0	0
3	A	48	0	24	3	0
4	A	346	0	0	8	0
All	All	3959	0	3599	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:306:GLY:O	1:A:309:THR:HG23	1.85	0.77
1:A:124[A]:ARG:HD2	1:A:331:TYR:HE1	1.50	0.76
1:A:107:HIS:ND1	1:A:124[B]:ARG:NH2	2.36	0.73
1:A:160:VAL:HG11	1:A:326:LEU:HD13	1.71	0.73
1:A:160:VAL:HG11	1:A:326:LEU:CD1	2.19	0.72
1:A:215:GLN:HG2	4:A:2186:HOH:O	1.90	0.70
1:A:29:HIS:HD2	1:A:31:ARG:H	1.41	0.65
1:A:104:ALA:HA	1:A:333:SER:HA	1.80	0.64
1:A:124[B]:ARG:HD3	1:A:331:TYR:HE1	1.64	0.63
1:A:309:THR:C	1:A:310:ARG:HD2	2.19	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:209:GLU:OE2	3:A:802:NAP:N7N	2.33	0.61
1:A:106:ASP:O	1:A:331:TYR:HD1	1.84	0.60
1:A:124[A]:ARG:HD2	1:A:331:TYR:CE1	2.33	0.59
1:A:77:TYR:CE1	1:A:411:LYS:HE2	2.39	0.57
1:A:229:LEU:HD22	4:A:2192:HOH:O	2.05	0.56
1:A:433:GLU:HG2	1:A:457:LEU:HD13	1.89	0.55
1:A:124[B]:ARG:HD3	1:A:331:TYR:CE1	2.43	0.53
1:A:108:GLN:HE22	1:A:332:LYS:HE3	1.75	0.52
1:A:282:GLN:OE1	4:A:2243:HOH:O	2.19	0.51
1:A:10:CYS:SG	1:A:98:VAL:HG22	2.51	0.51
1:A:404:ARG:HA	4:A:2300:HOH:O	2.10	0.51
1:A:309:THR:HG21	4:A:2251:HOH:O	2.12	0.50
1:A:29:HIS:CD2	1:A:31:ARG:H	2.27	0.48
1:A:309:THR:O	1:A:310:ARG:HD2	2.13	0.48
1:A:373:THR:O	3:A:802:NAP:O3D	2.32	0.47
1:A:396:ALA:HB1	1:A:398:HIS:CE1	2.50	0.47
1:A:375:VAL:HG12	3:A:802:NAP:H2N	1.95	0.47
1:A:215:GLN:NE2	4:A:2184:HOH:O	2.48	0.47
1:A:331:TYR:HD2	4:A:2290:HOH:O	1.98	0.46
1:A:231:LEU:HD23	1:A:250:LEU:HG	1.97	0.46
1:A:259:VAL:HG23	4:A:2214:HOH:O	2.16	0.45
1:A:55:HIS:N	1:A:56:PRO:CD	2.79	0.45
1:A:29:HIS:CD2	1:A:32:ALA:H	2.36	0.43
1:A:211:ARG:HG2	1:A:215:GLN:NE2	2.34	0.42
1:A:110:LEU:O	1:A:111:ASP:HB2	2.19	0.42
1:A:117:LEU:HA	1:A:118:PRO:HD3	1.86	0.41
1:A:19:PHE:HB3	1:A:65:PHE:CD1	2.56	0.41
1:A:229:LEU:O	1:A:229:LEU:HD23	2.20	0.41
1:A:224:ASP:HA	1:A:225:PRO:HD2	1.91	0.40
1:A:240:ARG:HB2	1:A:241:PRO:HD3	2.03	0.40
1:A:194:ILE:HB	1:A:274:LEU:HD12	2.04	0.40
1:A:223:LEU:HD21	1:A:274:LEU:HD22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	454/460 (99%)	440 (97%)	13 (3%)	1 (0%)	52 43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	404	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	375/379 (99%)	363 (97%)	12 (3%)	46 33

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	ARG
1	A	106	ASP
1	A	110	LEU
1	A	171	LEU
1	A	229	LEU
1	A	233	ASP
1	A	250	LEU
1	A	256	LYS
1	A	274	LEU
1	A	352	MET
1	A	425	SER
1	A	452	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	8	GLN
1	A	29	HIS
1	A	108	GLN
1	A	215	GLN
1	A	282	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	801	-	48,58,58	1.64	8 (16%)	54,89,89	2.35	14 (25%)
3	NAP	A	802	-	42,52,52	2.08	9 (21%)	54,80,80	3.22	20 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	801	-	-	0/30/50/50	0/6/6/6
3	NAP	A	802	-	1/1/12/12	0/27/67/67	0/5/5/5

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	802	NAP	P2B-O2B	-5.26	1.44	1.60
3	A	802	NAP	P2B-O3X	-4.98	1.36	1.54
2	A	801	FAD	C10-N10	-4.95	1.33	1.39
2	A	801	FAD	PA-O2A	-3.68	1.39	1.54
3	A	802	NAP	O4D-C1D	-3.02	1.37	1.41
2	A	801	FAD	P-O2P	-3.00	1.42	1.54
3	A	802	NAP	C8A-N7A	-2.80	1.29	1.34
2	A	801	FAD	C6-C5X	-2.52	1.37	1.41
3	A	802	NAP	O2B-C2B	-2.37	1.36	1.44
3	A	802	NAP	C5N-C4N	2.63	1.44	1.38
2	A	801	FAD	C5'-C4'	2.70	1.55	1.51
2	A	801	FAD	C2A-N3A	2.79	1.37	1.32
2	A	801	FAD	O5'-C5'	3.02	1.57	1.44
3	A	802	NAP	P2B-O1X	3.76	1.63	1.51
2	A	801	FAD	O4B-C1B	4.17	1.46	1.41
3	A	802	NAP	P2B-O2X	4.91	1.72	1.54
3	A	802	NAP	C3N-C7N	5.27	1.58	1.50

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	NAP	C5N-C4N-C3N	-6.68	111.93	120.33
3	A	802	NAP	O2X-P2B-O1X	-5.41	93.18	110.58
2	A	801	FAD	C4X-C4-N3	-5.00	116.75	123.59
2	A	801	FAD	C4-C4X-C10	-4.53	117.04	119.94
3	A	802	NAP	P2B-O2B-C2B	-4.45	110.90	121.56
2	A	801	FAD	N3A-C2A-N1A	-4.35	125.56	128.89
2	A	801	FAD	C4B-O4B-C1B	-4.33	104.96	109.72
2	A	801	FAD	C4X-C10-N10	-4.07	118.12	120.52
3	A	802	NAP	O3X-P2B-O2X	-3.50	94.05	107.38
3	A	802	NAP	C4N-C3N-C7N	-3.46	111.95	121.09
3	A	802	NAP	C5N-C6N-N1N	-3.02	115.25	120.47
3	A	802	NAP	C4D-O4D-C1D	-2.90	106.53	109.72
2	A	801	FAD	C6-C5X-N5	-2.84	115.31	118.96
3	A	802	NAP	O2B-C2B-C1B	-2.64	99.73	110.02
3	A	802	NAP	C4B-O4B-C1B	-2.46	107.01	109.72
3	A	802	NAP	C3N-C7N-N7N	-2.37	115.22	117.82

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	NAP	N3A-C2A-N1A	-2.33	127.11	128.89
2	A	801	FAD	O5B-PA-O1A	-2.05	101.67	109.62
2	A	801	FAD	O4B-C1B-N9A	-2.01	103.89	108.10
3	A	802	NAP	C3N-C2N-N1N	2.09	122.78	120.36
2	A	801	FAD	O3P-PA-O5B	2.11	108.53	102.94
2	A	801	FAD	C2A-N1A-C6A	2.16	122.63	118.77
3	A	802	NAP	O5D-C5D-C4D	2.18	117.17	109.12
2	A	801	FAD	O3B-C3B-C2B	2.21	119.00	111.83
3	A	802	NAP	O4B-C4B-C5B	2.29	117.51	109.32
3	A	802	NAP	C2N-C3N-C4N	2.30	120.85	118.29
2	A	801	FAD	C4A-C5A-N7A	2.40	111.69	109.48
3	A	802	NAP	C2N-C3N-C7N	2.71	127.19	119.31
3	A	802	NAP	O7N-C7N-C3N	2.77	122.61	119.59
2	A	801	FAD	C1'-N10-C9A	4.26	123.65	118.86
3	A	802	NAP	O4B-C1B-N9A	4.97	118.50	108.10
3	A	802	NAP	C6N-C5N-C4N	5.84	128.26	119.44
2	A	801	FAD	C4-N3-C2	9.98	123.87	115.25
3	A	802	NAP	O4D-C1D-N1N	16.33	126.07	108.13

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	802	NAP	C4B

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	802	NAP	3	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	455/460 (98%)	0.05	17 (3%) 45 56	13, 23, 44, 69	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	401	SER	7.3
1	A	331	TYR	4.5
1	A	403	PRO	4.5
1	A	346	LEU	4.0
1	A	345	LYS	3.9
1	A	402	GLY	3.4
1	A	398	HIS	3.4
1	A	99	VAL	2.8
1	A	236	LYS	2.7
1	A	399	LEU	2.6
1	A	400	PRO	2.6
1	A	305	ILE	2.5
1	A	10	CYS	2.5
1	A	237	GLU	2.4
1	A	310	ARG	2.4
1	A	12	VAL	2.3
1	A	31	ARG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAP	A	802	48/48	0.87	0.18	1.49	25,42,64,66	0
2	FAD	A	801	53/53	0.97	0.09	-0.51	13,17,22,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.