



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:42 AM GMT

PDB ID : 2E1C  
Title : Structure of Putative HTH-type transcriptional regulator PH1519/DNA Complex  
Authors : Koike, H.; Suzuki, M.  
Deposited on : 2006-10-24  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1-B	17	<div> <div>71%</div> <div>82% 18%</div> </div>
1	2-B	17	<div> <div>71%</div> <div>82% 18%</div> </div>
2	1-D	17	<div> <div>71%</div> <div>82% 18%</div> </div>
2	2-D	17	<div> <div>71%</div> <div>82% 18%</div> </div>
3	1-A	171	<div> <div>36%</div> <div>73% 13% 14%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	2-A	171	 A horizontal bar chart showing the quality of chain 2-A. The bar is divided into four segments: red (36%), green (73%), yellow (13%), and grey (14%). The segments are stacked horizontally, with the red segment starting from the left and ending at 36%, the green segment starting at 36% and ending at 73%, the yellow segment starting at 73% and ending at 86%, and the grey segment starting at 86% and ending at 100%.

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3958 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(\*DAP\*DGP\*DTP\*DGP\*DAP\*DAP\*DAP\*DAP\*DTP\*DTP\*DTP\*DTP\*DTP\*DTP\*DCP\*DAP\*DCP\*DA)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	1-B	17	Total	C	N	O	P	0	0	0
			346	168	63	99	16			
1	2-B	17	Total	C	N	O	P	0	0	0
			346	168	63	99	16			

- Molecule 2 is a DNA chain called DNA (5'-D(\*DTP\*DGP\*DTP\*DGP\*DAP\*DAP\*DAP\*DAP\*DAP\*DTP\*DTP\*DTP\*DTP\*DTP\*DCP\*DAP\*DCP\*DT)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	1-D	17	Total	C	N	O	P	0	0	0
			345	168	60	101	16			
2	2-D	17	Total	C	N	O	P	0	0	0
			345	168	60	101	16			

- Molecule 3 is a protein called Putative HTH-type transcriptional regulator PH1519.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	1-A	147	Total	C	N	O	S	0	0	0
			1155	741	190	221	3			
3	2-A	147	Total	C	N	O	S	0	0	0
			1155	741	190	221	3			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP O59188
A	2	GLY	-	EXPRESSION TAG	UNP O59188
A	3	SER	-	EXPRESSION TAG	UNP O59188
A	4	SER	-	EXPRESSION TAG	UNP O59188
A	5	HIS	-	EXPRESSION TAG	UNP O59188
A	6	HIS	-	EXPRESSION TAG	UNP O59188

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	HIS	-	EXPRESSION TAG	UNP O59188
A	8	HIS	-	EXPRESSION TAG	UNP O59188
A	9	HIS	-	EXPRESSION TAG	UNP O59188
A	10	HIS	-	EXPRESSION TAG	UNP O59188
A	11	SER	-	EXPRESSION TAG	UNP O59188
A	12	SER	-	EXPRESSION TAG	UNP O59188
A	13	GLY	-	EXPRESSION TAG	UNP O59188
A	14	LEU	-	EXPRESSION TAG	UNP O59188
A	15	VAL	-	EXPRESSION TAG	UNP O59188
A	16	PRO	-	EXPRESSION TAG	UNP O59188
A	17	ARG	-	EXPRESSION TAG	UNP O59188
A	18	GLY	-	EXPRESSION TAG	UNP O59188
A	19	SER	-	EXPRESSION TAG	UNP O59188
A	20	HIS	-	EXPRESSION TAG	UNP O59188


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	1-A	92	Total O 92 92	0	0
4	2-A	90	Total O 90 90	0	0
4	1-B	36	Total O 36 36	0	0
4	2-B	4	Total O 4 4	0	0
4	1-D	5	Total O 5 5	0	0
4	2-D	39	Total O 39 39	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (5'-D(\*DAP\*DGP\*DTP\*DGP\*DAP\*DAP\*DAP\*DAP\*DTP\*DTP\*DTP\*DT\*TP\*DTP\*DCP\*DAP\*DCP\*DA)-3')




- Molecule 1: DNA (5'-D(\*DAP\*DGP\*DTP\*DGP\*DAP\*DAP\*DAP\*DAP\*DTP\*DTP\*DTP\*DT\*TP\*DTP\*DCP\*DAP\*DCP\*DA)-3')




- Molecule 2: DNA (5'-D(\*DTP\*DGP\*DTP\*DGP\*DAP\*DAP\*DAP\*DAP\*DAP\*DTP\*DTP\*DT\*TP\*DTP\*DCP\*DAP\*DCP\*DT)-3')




- Molecule 2: DNA (5'-D(\*DTP\*DGP\*DTP\*DGP\*DAP\*DAP\*DAP\*DAP\*DAP\*DTP\*DTP\*DT\*TP\*DTP\*DCP\*DAP\*DCP\*DT)-3')




- Molecule 3: Putative HTH-type transcriptional regulator PH1519





● Molecule 3: Putative HTH-type transcriptional regulator PH1519



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.41Å 92.35Å 109.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	55.00 – 2.10 54.97 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.2 (55.00-2.10) 99.2 (54.97-2.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 2.00Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.206 , 0.230 0.289 , 0.301	Depositor DCC
$R_{free}$ test set	1051 reflections (5.42%)	DCC
Wilson B-factor (Å <sup>2</sup> )	41.1	Xtriage
Anisotropy	0.612	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.64 , 87.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	3 of 23589 reflections (0.013%)	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	3958	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	1-B	0.34	0/388	0.69	0/597
1	2-B	0.34	0/388	0.69	0/597
2	1-D	0.33	0/386	0.69	0/594
2	2-D	0.33	0/386	0.69	0/594
3	1-A	0.28	0/1171	0.59	0/1578
3	2-A	0.28	0/1171	0.59	0/1578
All	All	0.30	0/3890	0.63	0/5538

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	1-B	0	1
1	2-B	0	1
2	1-D	0	1
2	2-D	0	1
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	1-B	11	DT	Sidechain
2	1-D	11	DT	Sidechain
1	2-B	11	DT	Sidechain
2	2-D	11	DT	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1-B	346	0	195	1	0
1	2-B	346	0	195	1	0
2	1-D	345	0	196	1	0
2	2-D	345	0	196	1	0
3	1-A	1155	0	1217	16	0
3	2-A	1155	0	1217	18	0
4	1-A	92	0	0	1	0
4	1-B	36	0	0	0	0
4	1-D	5	0	0	0	0
4	2-A	90	0	0	1	0
4	2-B	4	0	0	0	0
4	2-D	39	0	0	0	0
All	All	3958	0	3216	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:37:GLN:HE22	3:A:168:LEU:H	1.36	0.73
3:A:37:GLN:HE22	3:A:168:LEU:H	1.35	0.72
3:A:60:GLU:OE2	3:A:63:ARG:NH2	2.21	0.72
3:A:60:GLU:OE2	3:A:63:ARG:NH2	2.23	0.72
3:A:66:ARG:HD3	3:A:74:PHE:HE1	1.57	0.69
3:A:66:ARG:HD3	3:A:74:PHE:HE1	1.60	0.65
3:A:91:ILE:HD13	3:A:141:LEU:HD23	1.82	0.61
3:A:91:ILE:HD13	3:A:141:LEU:HD23	1.84	0.60
3:A:115:GLU:HB2	3:A:128:LYS:HB3	1.85	0.58
3:A:115:GLU:HB2	3:A:128:LYS:HB3	1.86	0.57
3:A:37:GLN:NE2	3:A:168:LEU:H	2.02	0.56
3:A:37:GLN:NE2	3:A:168:LEU:H	2.04	0.56
3:A:100:TYR:CD2	3:A:118:GLU:HG3	2.43	0.53
3:A:100:TYR:CD2	3:A:118:GLU:HG3	2.48	0.49
3:A:66:ARG:HD2	4:A:215:HOH:O	2.12	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:63:ARG:NH1	3:A:67:GLU:OE1	2.47	0.48
3:A:72:LYS:O	3:A:73:LYS:HB3	2.14	0.48
1:B:3:DT:H2"	1:B:4:DG:C8	2.50	0.46
1:B:3:DT:H2"	1:B:4:DG:C8	2.50	0.46
2:D:3:DT:H2"	2:D:4:DG:C8	2.50	0.46
2:D:3:DT:H2"	2:D:4:DG:C8	2.50	0.46
3:A:45:ARG:O	3:A:49:LYS:HG2	2.16	0.45
3:A:63:ARG:NH1	3:A:67:GLU:CD	2.71	0.44
3:A:55:GLU:HG3	4:A:185:HOH:O	2.17	0.44
3:A:63:ARG:NH1	3:A:67:GLU:CD	2.72	0.43
3:A:45:ARG:O	3:A:49:LYS:HG2	2.18	0.43
3:A:92:LEU:HD11	3:A:156:MET:HE1	2.00	0.43
3:A:91:ILE:HD13	3:A:141:LEU:CD2	2.49	0.42
3:A:66:ARG:HD3	3:A:74:PHE:CE1	2.45	0.42
3:A:50:ILE:HD13	3:A:50:ILE:HA	1.92	0.42
3:A:63:ARG:NH1	3:A:67:GLU:OE1	2.52	0.42
3:A:72:LYS:O	3:A:73:LYS:HB3	2.20	0.41
3:A:107:LEU:HD21	3:A:144:ILE:HG12	2.02	0.41
3:A:92:LEU:HD11	3:A:156:MET:CE	2.50	0.41
3:A:107:LEU:HD21	3:A:144:ILE:HG12	2.02	0.41
3:A:45:ARG:HG2	3:A:45:ARG:HH11	1.87	0.40
3:A:91:ILE:HD13	3:A:141:LEU:CD2	2.48	0.40
3:A:92:LEU:HD11	3:A:156:MET:HE1	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	1-A	145/171 (85%)	142 (98%)	3 (2%)	0	100	100
3	2-A	145/171 (85%)	142 (98%)	3 (2%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	290/342 (85%)	284 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	1-A	130/151 (86%)	130 (100%)	0	100	100
3	2-A	130/151 (86%)	130 (100%)	0	100	100
All	All	260/302 (86%)	260 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
3	1-A	37	GLN
3	2-A	37	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	1-B	17/17 (100%)	2.44	12 (70%) 0 0	33, 48, 60, 65	17 (100%)
1	2-B	17/17 (100%)	2.44	12 (70%) 0 0	33, 48, 60, 65	17 (100%)
2	1-D	17/17 (100%)	2.38	12 (70%) 0 0	33, 47, 61, 65	17 (100%)
2	2-D	17/17 (100%)	2.38	12 (70%) 0 0	33, 47, 61, 65	17 (100%)
3	1-A	147/171 (85%)	1.92	62 (42%) 0 0	33, 44, 62, 81	147 (100%)
3	2-A	147/171 (85%)	1.92	62 (42%) 0 0	33, 44, 62, 81	147 (100%)
All	All	362/410 (88%)	2.01	172 (47%) 0 0	33, 45, 63, 81	362 (100%)

All (172) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	1-A	36	LEU	4.3
3	2-A	36	LEU	4.3
3	1-A	33	ILE	4.1
3	2-A	33	ILE	4.1
2	1-D	15	DA	4.0
2	2-D	15	DA	4.0
1	1-B	15	DA	3.9
1	2-B	15	DA	3.9
3	1-A	76	ALA	3.9
3	2-A	76	ALA	3.9
3	1-A	83	LEU	3.9
3	2-A	83	LEU	3.9
3	1-A	71	ILE	3.8
3	2-A	71	ILE	3.8
3	1-A	168	LEU	3.8
3	2-A	168	LEU	3.8
3	1-A	32	ILE	3.6
3	2-A	32	ILE	3.6
3	1-A	113	ILE	3.6

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
3	2-A	113	ILE	3.6
3	1-A	74	PHE	3.5
3	1-A	140	PHE	3.5
3	2-A	74	PHE	3.5
3	2-A	140	PHE	3.5
1	1-B	6	DA	3.4
1	2-B	6	DA	3.4
2	1-D	6	DA	3.4
2	2-D	6	DA	3.4
3	1-A	35	ILE	3.4
3	1-A	129	ILE	3.4
3	2-A	35	ILE	3.4
3	2-A	129	ILE	3.4
3	1-A	107	LEU	3.3
3	2-A	107	LEU	3.3
3	1-A	116	VAL	3.3
3	2-A	116	VAL	3.3
3	1-A	127	VAL	3.3
3	2-A	127	VAL	3.3
1	1-B	5	DA	3.2
1	2-B	5	DA	3.2
2	1-D	5	DA	3.2
2	2-D	5	DA	3.2
3	1-A	77	ILE	3.1
3	1-A	170	ILE	3.1
3	2-A	77	ILE	3.1
3	2-A	170	ILE	3.1
3	1-A	85	TYR	3.1
3	2-A	85	TYR	3.1
3	1-A	53	LEU	3.1
3	2-A	53	LEU	3.1
3	1-A	62	ILE	3.0
3	2-A	62	ILE	3.0
1	1-B	1	DA	3.0
1	1-B	7	DA	3.0
1	2-B	1	DA	3.0
1	2-B	7	DA	3.0
3	1-A	126	VAL	3.0
3	2-A	126	VAL	3.0
3	1-A	78	ILE	3.0
3	2-A	78	ILE	3.0
2	1-D	7	DA	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	2-D	7	DA	3.0
3	1-A	75	THR	3.0
3	2-A	75	THR	3.0
3	1-A	110	TYR	3.0
3	2-A	110	TYR	3.0
3	1-A	44	LEU	2.9
3	2-A	44	LEU	2.9
2	1-D	4	DG	2.9
2	2-D	4	DG	2.9
3	1-A	90	PHE	2.9
3	2-A	90	PHE	2.9
3	1-A	88	LEU	2.9
3	2-A	88	LEU	2.9
1	1-B	4	DG	2.8
1	2-B	4	DG	2.8
3	1-A	25	LEU	2.8
3	2-A	25	LEU	2.8
1	1-B	17	DA	2.8
1	2-B	17	DA	2.8
3	1-A	89	ALA	2.7
3	2-A	89	ALA	2.7
1	1-B	14	DC	2.7
1	2-B	14	DC	2.7
2	1-D	14	DC	2.7
2	2-D	14	DC	2.7
3	1-A	108	ALA	2.7
3	2-A	108	ALA	2.7
3	1-A	65	LEU	2.7
3	2-A	65	LEU	2.7
2	1-D	8	DA	2.7
2	2-D	8	DA	2.7
3	1-A	117	TYR	2.7
3	2-A	117	TYR	2.7
3	1-A	47	ILE	2.6
3	2-A	47	ILE	2.6
3	1-A	158	VAL	2.6
3	2-A	158	VAL	2.6
1	1-B	8	DA	2.6
1	2-B	8	DA	2.6
3	1-A	82	ALA	2.6
3	1-A	104	ALA	2.6
3	2-A	82	ALA	2.6

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	2-A	104	ALA	2.6
2	1-D	9	DA	2.6
2	2-D	9	DA	2.6
3	1-A	147	ILE	2.6
3	2-A	147	ILE	2.6
3	1-A	40	GLY	2.5
3	2-A	40	GLY	2.5
3	1-A	143	LEU	2.5
3	2-A	143	LEU	2.5
3	1-A	91	ILE	2.5
3	2-A	91	ILE	2.5
2	1-D	1	DT	2.5
2	2-D	1	DT	2.5
1	1-B	2	DG	2.5
1	2-B	2	DG	2.5
1	1-B	16	DC	2.4
1	2-B	16	DC	2.4
2	1-D	16	DC	2.4
2	2-D	16	DC	2.4
3	1-A	165	THR	2.4
3	2-A	165	THR	2.4
3	1-A	114	VAL	2.4
3	2-A	114	VAL	2.4
3	1-A	28	ILE	2.4
3	1-A	43	PRO	2.4
3	2-A	28	ILE	2.4
3	2-A	43	PRO	2.4
3	1-A	56	SER	2.4
3	2-A	56	SER	2.4
2	1-D	2	DG	2.3
2	2-D	2	DG	2.3
3	1-A	100	TYR	2.3
3	2-A	100	TYR	2.3
1	1-B	9	DT	2.3
1	2-B	9	DT	2.3
3	1-A	70	VAL	2.3
3	2-A	70	VAL	2.3
3	1-A	51	THR	2.2
3	2-A	51	THR	2.2
3	1-A	87	MET	2.2
3	2-A	87	MET	2.2
3	1-A	57	THR	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	2-A	57	THR	2.2
3	1-A	155	THR	2.2
3	2-A	155	THR	2.2
3	1-A	50	ILE	2.2
3	2-A	50	ILE	2.2
3	1-A	156	MET	2.2
3	2-A	156	MET	2.2
3	1-A	169	PRO	2.2
3	2-A	169	PRO	2.2
3	1-A	125	MET	2.1
3	2-A	125	MET	2.1
3	1-A	123	TYR	2.1
3	2-A	123	TYR	2.1
3	1-A	58	ILE	2.1
3	2-A	58	ILE	2.1
3	1-A	152	GLY	2.1
3	2-A	152	GLY	2.1
3	1-A	161	THR	2.0
3	2-A	161	THR	2.0
3	1-A	80	PRO	2.0
3	2-A	80	PRO	2.0
3	1-A	137	LEU	2.0
3	2-A	137	LEU	2.0
3	1-A	92	LEU	2.0
3	2-A	92	LEU	2.0
2	1-D	17	DT	2.0
2	2-D	17	DT	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.