



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:17 AM GMT

PDB ID : 3E2N
Title : Engineering ascorbate peroxidase activity into cytochrome c peroxidase
Authors : Poulos, T.L.; Mehareenna, Y.T.; Oertel, P.
Deposited on : 2008-08-05
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1475 (1.34-1.26)
Clashscore	102246	1031 (1.32-1.28)
Ramachandran outliers	100387	1504 (1.34-1.26)
Sidechain outliers	100360	1503 (1.34-1.26)
RSRZ outliers	91569	1476 (1.34-1.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	287	 79% 16% . .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2910 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cytochrome c peroxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	282	2278	1451	386	434	7	0	3	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	177	ARG	ASN	ENGINEERED	UNP P00431
A	184	PHE	TRP	ENGINEERED	UNP P00431

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	589	Total 589	O 589	0	3

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cytochrome c peroxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	106.78 Å 74.49 Å 50.95 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.30 32.12 – 1.25	Depositor EDS
% Data completeness (in resolution range)	91.9 (10.00-1.30) 96.3 (32.12-1.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.01 (at 1.25 Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.181 , (Not available) 0.169 , 0.173	Depositor DCC
R_{free} test set	4875 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	13.7	Xtriage
Anisotropy	0.650	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 55.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 108553 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2910	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	4/2357 (0.2%)	1.57	55/3183 (1.7%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	260	GLU	CD-OE1	9.77	1.36	1.25
1	A	287	LEU	C-OXT	-6.79	1.10	1.23
1	A	11	GLU	CD-OE2	-6.10	1.19	1.25
1	A	264	GLU	CD-OE1	5.04	1.31	1.25

All (55) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	136	ARG	NE-CZ-NH2	17.61	129.10	120.30
1	A	153	ARG	NE-CZ-NH1	13.04	126.82	120.30
1	A	123	ARG	NE-CZ-NH2	12.31	126.46	120.30
1	A	41[A]	ARG	NE-CZ-NH2	-10.66	114.97	120.30
1	A	41[B]	ARG	NE-CZ-NH2	-10.66	114.97	120.30
1	A	153	ARG	NE-CZ-NH2	-9.45	115.57	120.30
1	A	11	GLU	OE1-CD-OE2	8.77	133.82	123.30
1	A	120	ARG	NE-CZ-NH2	-8.51	116.04	120.30
1	A	18	ASP	CB-CG-OD2	-8.50	110.65	118.30
1	A	158	ASP	CB-CG-OD2	-8.17	110.94	118.30
1	A	146	TYR	CB-CG-CD1	8.09	125.85	121.00
1	A	177	ARG	NE-CZ-NH2	-8.07	116.27	120.30
1	A	60	TYR	CD1-CE1-CZ	-7.98	112.61	119.80
1	A	136	ARG	NE-CZ-NH1	-7.94	116.33	120.30
1	A	60	TYR	CB-CG-CD2	-7.88	116.27	121.00
1	A	23	TYR	CZ-CE2-CD2	-7.79	112.79	119.80
1	A	158	ASP	CB-CG-OD1	7.68	125.21	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	244	TYR	CB-CG-CD1	-7.65	116.41	121.00
1	A	254	ASP	CB-CG-OD1	7.29	124.86	118.30
1	A	23	TYR	CG-CD2-CE2	7.24	127.09	121.30
1	A	60	TYR	CG-CD2-CE2	-7.18	115.56	121.30
1	A	141	ASP	CB-CG-OD1	7.18	124.76	118.30
1	A	264	GLU	OE1-CD-OE2	-6.59	115.39	123.30
1	A	180	TYR	CB-CG-CD1	-6.49	117.11	121.00
1	A	44	TRP	CB-CG-CD2	6.47	135.01	126.60
1	A	148	ARG	NE-CZ-NH1	6.42	123.51	120.30
1	A	119	TRP	CG-CD1-NE1	6.28	116.38	110.10
1	A	259	PHE	CB-CG-CD2	6.22	125.16	120.80
1	A	41[A]	ARG	NE-CZ-NH1	6.19	123.39	120.30
1	A	41[B]	ARG	NE-CZ-NH1	6.19	123.39	120.30
1	A	119	TRP	CD1-NE1-CE2	-6.18	103.44	109.00
1	A	244	TYR	CZ-CE2-CD2	-6.08	114.32	119.80
1	A	64	TYR	CG-CD1-CE1	5.98	126.08	121.30
1	A	119	TRP	CH2-CZ2-CE2	-5.94	111.46	117.40
1	A	119	TRP	CG-CD2-CE3	-5.77	128.71	133.90
1	A	50	TRP	CD1-NE1-CE2	5.69	114.12	109.00
1	A	125	ASP	CB-CG-OD2	-5.67	113.19	118.30
1	A	180	TYR	CG-CD1-CE1	-5.56	116.86	121.30
1	A	191	PHE	CB-CG-CD1	-5.49	116.95	120.80
1	A	23	TYR	CB-CG-CD2	5.47	124.28	121.00
1	A	50	TRP	NE1-CE2-CD2	-5.46	101.84	107.30
1	A	151	PHE	CB-CG-CD1	5.46	124.62	120.80
1	A	191	PHE	CD1-CE1-CZ	-5.42	113.60	120.10
1	A	229	TYR	CB-CG-CD2	5.36	124.22	121.00
1	A	11	GLU	CG-CD-OE1	-5.34	107.63	118.30
1	A	191	PHE	CE1-CZ-CE2	5.31	129.56	120.00
1	A	260	GLU	CG-CD-OE2	5.30	128.90	118.30
1	A	159	ARG	NE-CZ-NH1	5.23	122.92	120.30
1	A	165	MET	CG-SD-CE	-5.21	91.86	100.20
1	A	119	TRP	CD1-CG-CD2	-5.20	102.14	106.30
1	A	19	PHE	CB-CG-CD1	-5.15	117.19	120.80
1	A	66	PHE	CG-CD1-CE1	5.15	126.46	120.80
1	A	123	ARG	NH1-CZ-NH2	-5.14	113.74	119.40
1	A	259	PHE	CG-CD2-CE2	5.14	126.45	120.80
1	A	237	TYR	CD1-CE1-CZ	-5.04	115.26	119.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2278	0	2179	23	0
2	A	43	0	30	0	0
3	A	589	0	0	11	0
All	All	2910	0	2209	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:LYS:HD3	3:A:1196:HOH:O	1.91	0.69
1:A:205:LYS:HD2	3:A:1087:HOH:O	1.94	0.67
1:A:17:GLU:HG2	3:A:1123:HOH:O	1.96	0.64
1:A:271:LYS:HG2	1:A:271:LYS:O	2.04	0.56
1:A:257:LYS:HE3	3:A:1477:HOH:O	2.07	0.54
1:A:208:LYS:HD3	1:A:212:ASN:ND2	2.24	0.53
1:A:176:LYS:CD	1:A:176:LYS:H	2.22	0.52
1:A:186:ALA:HB2	1:A:222:TYR:OH	2.09	0.51
1:A:259:PHE:O	1:A:263:LEU:HG	2.12	0.50
1:A:6:HIS:N	3:A:1011:HOH:O	2.46	0.49
1:A:172:LYS:HE2	1:A:172:LYS:HB3	1.60	0.47
1:A:41[B]:ARG:NH2	3:A:1052:HOH:O	2.48	0.46
1:A:6:HIS:N	3:A:1008:HOH:O	2.48	0.46
1:A:46:THR:HG22	1:A:64:TYR:HB2	1.98	0.46
1:A:90:LYS:HG3	3:A:1307:HOH:O	2.15	0.45
1:A:253:LYS:HE2	3:A:1201:HOH:O	2.16	0.44
1:A:205:LYS:HD3	3:A:1377:HOH:O	2.16	0.43
1:A:203:ASP:HB2	1:A:219:LYS:HD3	2.01	0.43
1:A:176:LYS:H	1:A:176:LYS:HD2	1.83	0.42
1:A:219:LYS:HG2	3:A:1491:HOH:O	2.20	0.41
1:A:203:ASP:OD1	1:A:219:LYS:HE3	2.21	0.41
1:A:86:GLU:N	1:A:87:PRO:HD2	2.36	0.41
1:A:205:LYS:HA	1:A:205:LYS:HD2	1.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/287 (99%)	282 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/246 (99%)	235 (96%)	9 (4%)	41	5

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	GLU
1	A	67	LYS
1	A	172	LYS
1	A	176	LYS
1	A	193	ASN
1	A	219	LYS
1	A	253	LYS
1	A	271	LYS
1	A	274	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	152	GLN
1	A	189	ASN
1	A	193	ASN
1	A	201	ASN
1	A	212	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	296	1,3	30,50,50	2.69	10 (33%)	24,82,82	2.32	9 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	296	1,3	-	0/10/54/54	0/0/8/8

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	296	HEM	C3B-C4B	-6.98	1.45	1.51
2	A	296	HEM	C2D-C3D	-6.11	1.36	1.54
2	A	296	HEM	C3D-C4D	-5.75	1.44	1.51
2	A	296	HEM	C3B-CAB	-4.93	1.42	1.51
2	A	296	HEM	C2C-C1C	-4.19	1.44	1.52
2	A	296	HEM	C3C-CAC	-3.39	1.45	1.51
2	A	296	HEM	C2B-C1B	-2.76	1.42	1.51
2	A	296	HEM	C2D-C1D	-2.08	1.45	1.51
2	A	296	HEM	CBB-CAB	2.86	1.45	1.29
2	A	296	HEM	C4C-NC	3.24	1.40	1.36

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	296	HEM	CAA-CBA-CGA	-2.59	108.00	112.75
2	A	296	HEM	CMA-C3A-C4A	-2.38	124.43	128.36
2	A	296	HEM	C3B-C4B-CHC	2.28	126.38	123.16
2	A	296	HEM	C2D-C3D-C4D	3.26	107.02	101.50
2	A	296	HEM	CMD-C2D-C3D	3.40	129.39	114.35
2	A	296	HEM	CAD-C3D-C4D	3.74	125.67	112.47
2	A	296	HEM	CMC-C2C-C3C	4.68	128.21	116.53
2	A	296	HEM	CMB-C2B-C3B	4.70	128.26	116.53
2	A	296	HEM	CAD-C3D-C2D	4.77	126.92	113.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/287 (98%)	-0.44	2 (0%) 89 87	12, 18, 33, 52	1 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	187	ALA	2.8
1	A	170	LEU	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	HEM	A	296	43/43	0.98	0.07	-0.22	11,13,19,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.