



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 07:07 PM GMT

PDB ID : 1E4V  
Title : MUTANT G10V OF ADENYLATE KINASE FROM E. COLI, MODIFIED  
IN THE GLY-LOOP  
Authors : Mueller, C.W.; Schulz, G.E.  
Deposited on : 2000-07-12  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.



Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	214	 83% 14% •
1	B	214	 75% 21% • •

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3780 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ADENYLATE KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total	C	N	O	S	0	0	0
			1659	1043	289	320	7			
1	B	214	Total	C	N	O	S	0	0	0
			1659	1043	289	320	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	10	VAL	GLY	ENGINEERED MUTATION	UNP P05082
B	10	VAL	GLY	ENGINEERED MUTATION	UNP P05082

- Molecule 2 is BIS(ADENOSINE)-5'-PENTAPHOSPHATE (three-letter code: AP5) (formula:  $C_{20}H_{29}N_{10}O_{22}P_5$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			57	20	10	22	5		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			57	20	10	22	5		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	220	Total	O	0	0
			220	220		
3	B	128	Total	O	0	0
			128	128		

### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

#### • Molecule 1: ADENYLATE KINASE



#### • Molecule 1: ADENYLATE KINASE



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.20 Å   79.80 Å   85.00 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	10.00 – 1.85	Depositor
% Data completeness (in resolution range)	90.0 (10.00-1.85)	Depositor
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.10	Depositor
Refinement program	X-PLOR 1.5	Depositor
R, $R_{free}$	0.196 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3780	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AP5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.94	0/1683	1.61	18/2267 (0.8%)
1	B	0.82	0/1683	1.51	13/2267 (0.6%)
All	All	0.88	0/3366	1.56	31/4534 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	167	ARG	NE-CZ-NH2	-9.43	115.58	120.30
1	A	167	ARG	NE-CZ-NH1	9.15	124.88	120.30
1	A	2	ARG	NE-CZ-NH1	8.65	124.63	120.30
1	A	88	ARG	NE-CZ-NH2	-7.51	116.54	120.30
1	A	165	ARG	NE-CZ-NH1	7.35	123.98	120.30
1	B	167	ARG	NE-CZ-NH1	7.12	123.86	120.30
1	B	64	VAL	CG1-CB-CG2	-7.02	99.67	110.90
1	A	47	LYS	CG-CD-CE	-6.96	91.00	111.90
1	B	88	ARG	NE-CZ-NH2	-6.77	116.91	120.30
1	B	164	VAL	CA-CB-CG2	-6.66	100.92	110.90
1	A	111	VAL	CG1-CB-CG2	-6.56	100.41	110.90
1	A	88	ARG	NE-CZ-NH1	6.26	123.43	120.30
1	A	105	TYR	CB-CG-CD1	-6.20	117.28	121.00
1	B	165	ARG	CA-CB-CG	-6.05	100.10	113.40

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	111	VAL	CA-CB-CG2	-5.99	101.92	110.90
1	B	126	HIS	N-CA-C	-5.98	94.85	111.00
1	A	88	ARG	CG-CD-NE	-5.94	99.32	111.80
1	A	157	LYS	CA-CB-CG	5.94	126.47	113.40
1	B	124	ARG	NE-CZ-NH1	5.68	123.14	120.30
1	A	10	VAL	CG1-CB-CG2	-5.59	101.96	110.90
1	A	174	MET	CA-CB-CG	5.58	122.79	113.30
1	B	117	VAL	CG1-CB-CG2	-5.53	102.05	110.90
1	A	193	TYR	CB-CG-CD2	-5.49	117.70	121.00
1	A	165	ARG	CA-CB-CG	-5.45	101.40	113.40
1	B	111	VAL	CA-CB-CG1	5.38	118.96	110.90
1	A	156	ARG	NE-CZ-NH2	5.26	122.93	120.30
1	B	120	ILE	CA-CB-CG1	-5.24	101.05	111.00
1	A	202	VAL	CG1-CB-CG2	-5.22	102.55	110.90
1	B	123	ARG	NE-CZ-NH1	5.13	122.86	120.30
1	B	157	LYS	CA-CB-CG	5.12	124.67	113.40
1	A	118	ASP	CB-CG-OD1	5.00	122.80	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	193	TYR	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1659	0	1691	16	1
1	B	1659	0	1691	26	0
2	A	57	0	23	0	0
2	B	57	0	22	2	0
3	A	220	0	0	4	0
3	B	128	0	0	7	0
All	All	3780	0	3427	42	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 6.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:19:PHE:HZ	1:B:210:GLU:HG3	1.49	0.77
1:A:120:ILE:HG12	1:A:164:VAL:HG21	1.71	0.72
1:A:75:GLU:O	1:A:78:ARG:HG3	1.90	0.71
1:B:19:PHE:CZ	1:B:210:GLU:HG3	2.29	0.67
1:B:195:LYS:NZ	3:B:2114:HOH:O	2.31	0.63
1:B:9:PRO:O	1:B:116:ILE:HD12	1.99	0.63
1:A:29:ILE:HD11	1:A:77:CYS:SG	2.42	0.60
1:B:120:ILE:HG12	1:B:164:VAL:HG21	1.85	0.58
1:B:75:GLU:O	1:B:78:ARG:HG3	2.04	0.57
1:B:119:ARG:HA	1:B:134:HIS:HE2	1.71	0.55
1:A:156:ARG:HD3	3:A:2159:HOH:O	2.08	0.54
1:B:2:ARG:HG2	1:B:81:PHE:CE1	2.43	0.53
1:B:117:VAL:HG23	3:B:2062:HOH:O	2.09	0.52
1:A:10:VAL:HG11	1:A:120:ILE:HD12	1.91	0.52
1:B:16:GLN:HG3	1:B:109:PHE:HE2	1.76	0.50
1:A:10:VAL:HG11	1:A:120:ILE:CD1	2.41	0.50
1:B:71:ARG:NH1	1:B:77:CYS:SG	2.85	0.50
1:A:172:HIS:HB3	1:B:150:GLY:O	2.11	0.50
1:B:22:GLU:HG3	1:B:23:LYS:N	2.28	0.49
1:B:136:LYS:NZ	3:B:2071:HOH:O	2.39	0.48
1:A:148:VAL:HG11	3:A:2139:HOH:O	2.14	0.47
1:B:127:ALA:HB3	1:B:128:PRO:HD3	1.97	0.46
1:B:202:VAL:HG23	3:B:2118:HOH:O	2.16	0.46
1:B:167:ARG:NH1	3:B:2095:HOH:O	2.44	0.46
1:A:126:HIS:CD2	1:A:128:PRO:HD2	2.53	0.44
1:A:6:LEU:HG	3:A:2173:HOH:O	2.18	0.44
1:A:9:PRO:O	1:A:116:ILE:HD12	2.18	0.43
1:A:4:ILE:HD12	1:A:103:VAL:HG21	2.00	0.43
1:B:61:ASP:O	1:B:65:ILE:HG13	2.18	0.43
1:B:180:GLY:O	1:B:184:LYS:HB2	2.18	0.43
1:B:38:ALA:HB2	3:B:2013:HOH:O	2.19	0.43
1:B:206:ARG:O	1:B:209:LEU:HB2	2.18	0.42
1:A:29:ILE:HG21	1:A:29:ILE:HD13	1.82	0.42
1:B:71:ARG:O	1:B:74:GLN:HB2	2.20	0.42
1:B:55:ALA:O	1:B:166:LYS:HD2	2.20	0.41
1:A:19:PHE:HZ	1:A:210:GLU:CG	2.33	0.41
1:B:119:ARG:HA	1:B:134:HIS:NE2	2.35	0.41
1:A:88:ARG:NH2	3:A:2107:HOH:O	2.53	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186:ALA:HA	1:A:191:THR:O	2.20	0.41
1:B:59:VAL:HG12	1:B:64:VAL:HG23	2.02	0.41
1:B:137:PHE:HB3	2:B:215:AP5:H2A	2.02	0.40
2:B:215:AP5:H4J	3:B:2125:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:GLY:O	1:A:100:GLY:O[2_656]	2.04	0.16

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/214 (99%)	205 (97%)	7 (3%)	0	100	100
1	B	212/214 (99%)	207 (98%)	5 (2%)	0	100	100
All	All	424/428 (99%)	412 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/176 (100%)	170 (97%)	6 (3%)	44	24
1	B	176/176 (100%)	161 (92%)	15 (8%)	13	2
All	All	352/352 (100%)	331 (94%)	21 (6%)	24	7

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	54	ASP
1	A	78	ARG
1	A	88	ARG
1	A	120	ILE
1	A	157	LYS
1	A	168	LEU
1	B	2	ARG
1	B	43	SER
1	B	54	ASP
1	B	75	GLU
1	B	78	ARG
1	B	88	ARG
1	B	91	PRO
1	B	120	ILE
1	B	125	VAL
1	B	135	VAL
1	B	157	LYS
1	B	167	ARG
1	B	168	LEU
1	B	174	MET
1	B	192	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	138	ASN
1	B	160	GLN
1	B	172	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	AP5	A	215	-	44,62,62	1.11	5 (11%)	55,98,98	2.05	12 (21%)
2	AP5	B	215	-	44,62,62	0.90	2 (4%)	55,98,98	2.20	14 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AP5	A	215	-	-	0/36/76/76	0/6/6/6
2	AP5	B	215	-	-	0/36/76/76	0/6/6/6

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	215	AP5	C8A-N7A	-2.44	1.29	1.34
2	B	215	AP5	C8B-N7B	-2.34	1.30	1.34
2	A	215	AP5	C8B-N7B	-2.02	1.30	1.34
2	A	215	AP5	C8A-N7A	-2.01	1.30	1.34
2	A	215	AP5	O4J-C1J	2.13	1.43	1.41
2	A	215	AP5	C5F-C4F	2.14	1.58	1.51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	215	AP5	C3F-C4F	2.49	1.59	1.53

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	215	AP5	C4B-C5B-N7B	-3.77	106.01	109.48
2	A	215	AP5	C4J-O4J-C1J	-3.44	105.94	109.72
2	B	215	AP5	C4A-C5A-N7A	-2.73	106.96	109.48
2	B	215	AP5	C4J-O4J-C1J	-2.71	106.74	109.72
2	A	215	AP5	C4A-C5A-N7A	-2.67	107.03	109.48
2	B	215	AP5	PD-O3G-PG	-2.11	126.81	132.73
2	A	215	AP5	O2B-PB-O3B	2.18	114.96	105.09
2	B	215	AP5	O2G-PG-O3G	2.30	115.53	105.09
2	A	215	AP5	O3A-PA-O5F	2.50	109.57	102.94
2	A	215	AP5	O2A-PA-O3A	2.60	116.88	105.09
2	B	215	AP5	N6A-C6A-N1A	2.62	124.83	119.20
2	B	215	AP5	C1F-N9A-C4A	2.70	131.02	126.94
2	B	215	AP5	O2A-PA-O3A	2.85	118.03	105.09
2	A	215	AP5	N3A-C2A-N1A	3.28	131.41	128.89
2	B	215	AP5	N3B-C2B-N1B	3.48	131.56	128.89
2	B	215	AP5	O2B-PB-O3B	3.58	121.32	105.09
2	B	215	AP5	N3A-C2A-N1A	3.69	131.72	128.89
2	A	215	AP5	N3B-C2B-N1B	3.84	131.83	128.89
2	A	215	AP5	C2J-C1J-N9B	4.00	120.40	114.29
2	B	215	AP5	C2F-C1F-N9A	5.07	122.03	114.29
2	B	215	AP5	O4J-C1J-N9B	5.68	119.98	108.10
2	A	215	AP5	C2F-C1F-N9A	5.71	123.01	114.29
2	A	215	AP5	O4J-C1J-N9B	5.99	120.64	108.10
2	B	215	AP5	C2J-C1J-N9B	6.20	123.77	114.29
2	A	215	AP5	O4F-C1F-N9A	6.24	121.17	108.10
2	B	215	AP5	O4F-C1F-N9A	7.39	123.56	108.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	215	AP5	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.