



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:07 PM GMT

PDB ID : 1E6S  
Title : MYROSINASE FROM SINAPIS ALBA WITH BOUND GLUCO-HYDROX  
IMOLACTAM AND SULFATE  
Authors : Burmeister, W.P.  
Deposited on : 2000-08-23  
Resolution : 1.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2199 (1.40-1.32)
Clashscore	102246	2337 (1.40-1.32)
Ramachandran outliers	100387	2280 (1.40-1.32)
Sidechain outliers	100360	2279 (1.40-1.32)
RSRZ outliers	91569	2199 (1.40-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	M	501	<div> <div></div> <div>83%</div> <div>14%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	M	901	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	M	931	-	-	X	-
2	NAG	M	961	-	-	-	X
5	BMA	M	954	-	-	X	-
5	MAN	M	957	X	-	X	-
6	GOX	M	999	-	-	-	X
8	SO4	M	1503	-	-	-	X
8	SO4	M	1505[A]	-	X	-	X
8	SO4	M	1505[B]	-	-	X	-
8	SO4	M	1533	-	-	-	X
9	GOL	M	1510[A]	-	-	-	X
9	GOL	M	1510[B]	-	-	-	X
9	GOL	M	1513	-	-	-	X

## 2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 5207 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MYROSINASE MA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	M	499	Total	C	N	O	S	0	21	0
			4082	2618	660	788	16			

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	M	1	Total	C	N	O	0	0
			14	8	1	5		
2	M	1	Total	C	N	O	0	0
			14	8	1	5		
2	M	1	Total	C	N	O	0	0
			14	8	1	5		
2	M	1	Total	C	N	O	0	0
			14	8	1	5		
2	M	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	M	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	M	2	Total	C	N	O	0	0
			28	16	2	10		


- Molecule 4 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	M	5	Total	C	N	O	0	0
			58	33	2	23		

- Molecule 5 is a polymer of unknown type called SUGAR (7-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	M	7	Total	C	N	O	0	0
			80	45	2	33		

- Molecule 6 is (2S,3S,4R,5R)-6-(HYDROXYAMINO)-2-(HYDROXYMETHYL)-2,3,4,5-TE TRAHYDROPYRIDINE-3,4,5-TRIOL (three-letter code: GOX) (formula: C<sub>6</sub>H<sub>12</sub>N<sub>2</sub>O<sub>5</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	M	1	Total	C	N	O	0	0
			13	6	2	5		

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	M	1	Total	Zn	0	0
			1	1		

- Molecule 8 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	M	1	Total	O	S	0	0
			5	4	1		
8	M	1	Total	O	S	0	0
			5	4	1		
8	M	1	Total	O	S	0	1
			10	8	2		
8	M	1	Total	O	S	0	0
			5	4	1		
8	M	1	Total	O	S	0	0
			5	4	1		
8	M	1	Total	O	S	0	0
			5	4	1		
8	M	1	Total	O	S	0	0
			5	4	1		

- Molecule 9 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	M	1	Total	C	O	0	1
			7	3	4		
9	M	1	Total	C	O	0	0
			6	3	3		
9	M	1	Total	C	O	0	0
			6	3	3		
9	M	1	Total	C	O	0	0
			6	3	3		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	M	786	Total	O	0	0
			786	786		
10	N	5	Total	O	0	0
			5	5		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: MYROSINASE MA1





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	135.30Å 137.20Å 80.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.35 9.99 – 1.35	Depositor EDS
% Data completeness (in resolution range)	82.1 (10.00-1.35) 81.7 (9.99-1.35)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.55 (at 1.35Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.120 , 0.152 0.137 , 0.163	Depositor DCC
$R_{free}$ test set	6736 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	10.9	Xtriage
Anisotropy	0.269	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.51 , 62.6	EDS
Estimated twinning fraction	0.014 for -k,-h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 133304 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5207	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: XYP, GOL, ZN, BMA, NAG, SO4, FUC, MAN, GOX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	M	0.98	5/4290 (0.1%)	1.50	56/5833 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
5	M	1	0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	M	24	SER	CB-OG	7.01	1.51	1.42
1	M	178	SER	CB-OG	6.19	1.50	1.42
1	M	151	GLU	CD-OE2	-5.99	1.19	1.25
1	M	115	GLU	CD-OE2	5.23	1.31	1.25
1	M	501	PRO	N-CD	5.02	1.54	1.47

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	M	109	ARG	NE-CZ-NH2	-18.31	111.14	120.30
1	M	269	ARG	NE-CZ-NH2	-13.15	113.73	120.30
1	M	115	GLU	OE1-CD-OE2	-13.02	107.67	123.30
1	M	109	ARG	NH1-CZ-NH2	10.98	131.48	119.40
1	M	57	ARG	NE-CZ-NH2	-9.36	115.62	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	M	957	MAN	C5

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	M	4082	0	3835	91	0
2	M	84	0	77	19	0
3	M	28	0	25	5	0
4	M	58	0	50	5	0
5	M	80	0	66	17	0
6	M	13	0	12	1	0
7	M	1	0	0	0	0
8	M	45	0	0	5	0
9	M	25	0	29	1	0
10	M	786	0	0	33	1
10	N	5	0	0	0	0
All	All	5207	0	4094	108	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 108 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:21:ASN:HD21	2:M:901:NAG:C1	0.92	1.56
1:M:244:ASN:HD21	2:M:931:NAG:C1	0.96	1.54
1:M:90:ASN:HD21	2:M:911:NAG:C1	0.92	1.54
1:M:265:ASN:HD21	4:M:941:NAG:C1	0.88	1.51
1:M:218:ASN:HD21	3:M:921:NAG:C1	0.86	1.49

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:M:2064:HOH:O	10:M:2737:HOH:O[6_565]	1.95	0.25

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	M	518/501 (103%)	503 (97%)	14 (3%)	1 (0%)	52 22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	M	187	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	M	456/437 (104%)	455 (100%)	1 (0%)	95 84

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	3	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	M	218	ASN
1	M	365	HIS
1	M	244	ASN
1	M	122	HIS
1	M	265	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

14 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	M	921	1,3	14,14,15	2.57	7 (50%)	15,19,21	3.60	8 (53%)
3	NAG	M	923	3	14,14,15	2.28	3 (21%)	15,19,21	4.52	6 (40%)
4	NAG	M	941	1,4	14,14,15	1.03	1 (7%)	15,19,21	3.21	7 (46%)
4	FUC	M	942	4	10,10,11	2.00	2 (20%)	14,14,16	2.27	3 (21%)
4	NAG	M	943	4	14,14,15	1.43	1 (7%)	15,19,21	1.70	3 (20%)
4	BMA	M	944	4	11,11,12	2.01	3 (27%)	14,15,17	1.39	1 (7%)
4	XYP	M	945	4	9,9,10	1.31	1 (11%)	12,12,14	2.23	5 (41%)
5	NAG	M	951	1,5	14,14,15	1.60	2 (14%)	15,19,21	2.84	4 (26%)
5	FUC	M	952	5	10,10,11	2.21	5 (50%)	14,14,16	2.80	7 (50%)
5	NAG	M	953	5	14,14,15	1.43	2 (14%)	15,19,21	2.37	6 (40%)
5	BMA	M	954	5	11,11,12	2.54	3 (27%)	14,15,17	4.64	8 (57%)
5	XYP	M	955	5	9,9,10	2.19	2 (22%)	12,12,14	3.75	8 (66%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	MAN	M	956	5	11,11,12	1.52	3 (27%)	14,15,17	2.62	6 (42%)
5	MAN	M	957	5	11,11,12	2.74	5 (45%)	14,15,17	7.30	11 (78%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	M	921	1,3	-	0/6/23/26	0/1/1/1
3	NAG	M	923	3	-	0/6/23/26	0/1/1/1
4	NAG	M	941	1,4	-	0/6/23/26	0/1/1/1
4	FUC	M	942	4	-	0/0/17/20	0/1/1/1
4	NAG	M	943	4	-	0/6/23/26	0/1/1/1
4	BMA	M	944	4	-	0/2/19/22	0/1/1/1
4	XYP	M	945	4	-	0/0/14/17	0/1/1/1
5	NAG	M	951	1,5	-	0/6/23/26	0/1/1/1
5	FUC	M	952	5	-	0/0/17/20	0/1/1/1
5	NAG	M	953	5	-	0/6/23/26	0/1/1/1
5	BMA	M	954	5	-	0/2/19/22	0/1/1/1
5	XYP	M	955	5	-	0/0/14/17	0/1/1/1
5	MAN	M	956	5	-	0/2/19/22	0/1/1/1
5	MAN	M	957	5	1/1/4/5	0/2/19/22	0/1/1/1

The worst 5 of 40 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	M	954	BMA	C2-C3	-6.91	1.43	1.52
3	M	921	NAG	O7-C7	-4.88	1.11	1.23
4	M	944	BMA	C2-C3	-4.61	1.46	1.52
3	M	921	NAG	C4-C5	-4.29	1.44	1.53
5	M	951	NAG	C1-C2	-4.28	1.46	1.52

The worst 5 of 83 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	M	923	NAG	C1-O5-C5	-14.89	93.36	112.25
5	M	957	MAN	C1-O5-C5	-14.34	94.05	112.25
5	M	957	MAN	O5-C5-C6	-10.09	85.50	107.35
3	M	921	NAG	C1-O5-C5	-9.43	100.29	112.25
5	M	951	NAG	C1-O5-C5	-9.04	100.78	112.25

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	M	957	MAN	C5

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 27 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	M	921	NAG	5	0
4	M	941	NAG	5	0
5	M	951	NAG	2	0
5	M	954	BMA	10	0
5	M	957	MAN	12	0

## 5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 1 is monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
8	SO4	M	1503	-	4,4,4	1.55	1 (25%)	6,6,6	0.82	0
8	SO4	M	1504	-	4,4,4	0.86	0	6,6,6	0.64	0
8	SO4	M	1505[A]	-	4,4,4	2.12	2 (50%)	6,6,6	2.55	3 (50%)
8	SO4	M	1505[B]	-	4,4,4	1.14	0	6,6,6	1.23	1 (16%)
8	SO4	M	1506	-	4,4,4	0.85	0	6,6,6	0.17	0
8	SO4	M	1507	-	4,4,4	2.75	1 (25%)	6,6,6	3.12	2 (33%)
8	SO4	M	1508	-	4,4,4	1.40	1 (25%)	6,6,6	1.25	0
8	SO4	M	1509	-	4,4,4	1.02	0	6,6,6	0.47	0
9	GOL	M	1510[A]	-	5,5,5	0.86	0	5,5,5	1.18	1 (20%)
9	GOL	M	1510[B]	-	5,5,5	0.95	0	5,5,5	2.79	2 (40%)
9	GOL	M	1511	-	5,5,5	0.91	0	5,5,5	1.13	0
9	GOL	M	1512	-	5,5,5	0.75	0	5,5,5	1.46	0
9	GOL	M	1513	-	5,5,5	0.20	0	5,5,5	1.11	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
8	SO4	M	1533	-	4,4,4	0.84	0	6,6,6	0.75	0
2	NAG	M	901	1	14,14,15	1.20	2 (14%)	15,19,21	2.09	4 (26%)
2	NAG	M	911	1	14,14,15	1.43	2 (14%)	15,19,21	2.52	3 (20%)
2	NAG	M	931	1	14,14,15	1.71	3 (21%)	15,19,21	6.71	11 (73%)
2	NAG	M	961	1	14,14,15	1.36	1 (7%)	15,19,21	2.20	5 (33%)
2	NAG	M	971	1	14,14,15	1.25	1 (7%)	15,19,21	1.81	3 (20%)
2	NAG	M	991	1	14,14,15	1.21	1 (7%)	15,19,21	2.13	4 (26%)
6	GOX	M	999	-	12,13,13	1.26	2 (16%)	9,18,18	2.27	4 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	SO4	M	1503	-	-	0/0/0/0	0/0/0/0
8	SO4	M	1504	-	-	0/0/0/0	0/0/0/0
8	SO4	M	1505[A]	-	-	0/0/0/0	0/0/0/0
8	SO4	M	1505[B]	-	-	0/0/0/0	0/0/0/0
8	SO4	M	1506	-	-	0/0/0/0	0/0/0/0
8	SO4	M	1507	-	-	0/0/0/0	0/0/0/0
8	SO4	M	1508	-	-	0/0/0/0	0/0/0/0
8	SO4	M	1509	-	-	0/0/0/0	0/0/0/0
9	GOL	M	1510[A]	-	-	0/4/4/4	0/0/0/0
9	GOL	M	1510[B]	-	-	0/4/4/4	0/0/0/0
9	GOL	M	1511	-	-	0/4/4/4	0/0/0/0
9	GOL	M	1512	-	-	0/4/4/4	0/0/0/0
9	GOL	M	1513	-	-	0/4/4/4	0/0/0/0
8	SO4	M	1533	-	-	0/0/0/0	0/0/0/0
2	NAG	M	901	1	-	0/6/23/26	0/1/1/1
2	NAG	M	911	1	-	0/6/23/26	0/1/1/1
2	NAG	M	931	1	-	0/6/23/26	0/1/1/1
2	NAG	M	961	1	-	0/6/23/26	0/1/1/1
2	NAG	M	971	1	-	0/6/23/26	0/1/1/1
2	NAG	M	991	1	-	0/6/23/26	0/1/1/1
6	GOX	M	999	-	-	0/2/24/24	0/1/1/1

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	M	971	NAG	O7-C7	-4.06	1.13	1.23

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	M	911	NAG	O7-C7	-3.87	1.14	1.23
2	M	991	NAG	O7-C7	-3.71	1.14	1.23
2	M	961	NAG	O7-C7	-3.67	1.14	1.23
2	M	931	NAG	O7-C7	-3.41	1.15	1.23

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	M	931	NAG	C1-O5-C5	-10.07	99.46	112.25
2	M	931	NAG	O7-C7-N2	-9.84	101.80	121.86
2	M	911	NAG	C1-O5-C5	-7.92	102.19	112.25
8	M	1507	SO4	O2-S-O1	-6.70	88.27	109.50
2	M	931	NAG	O7-C7-C8	-5.66	111.67	122.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 26 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	M	1505[B]	SO4	5	0
9	M	1513	GOL	1	0
2	M	901	NAG	4	0
2	M	911	NAG	3	0
2	M	931	NAG	12	0
6	M	999	GOX	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	M	499/501 (99%)	-0.37	7 (1%) 78 78	8, 12, 23, 51	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	M	376	ALA	5.8
1	M	380	ASP	4.2
1	M	3	GLU	4.1
1	M	379	THR	3.2
1	M	481	ASN	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	M	921	14/15	0.94	0.08	1.32	17,21,29,30	0
4	NAG	M	941	14/15	0.94	0.07	0.72	15,18,21,21	0
5	NAG	M	951	14/15	0.93	0.07	-	15,16,20,24	0
5	BMA	M	954	11/12	0.89	0.10	-	20,25,29,30	0
3	NAG	M	923	14/15	0.57	0.33	-	36,41,47,48	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	XYP	M	955	9/10	0.53	0.26	-	34,37,40,42	0
5	FUC	M	952	10/11	0.93	0.10	-	16,18,21,23	0
4	FUC	M	942	10/11	0.85	0.20	-	24,28,33,36	0
5	NAG	M	953	14/15	0.89	0.11	-	18,20,27,29	0
4	NAG	M	943	14/15	0.93	0.09	-	20,24,29,33	0
4	BMA	M	944	11/12	0.79	0.29	-	33,37,41,43	0
5	MAN	M	957	11/12	0.69	0.34	-	32,41,47,52	0
4	XYP	M	945	9/10	0.76	0.30	-	38,39,40,41	0
5	MAN	M	956	11/12	0.70	0.25	-	22,32,38,38	0

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	M	961	14/15	0.57	0.42	59.36	28,35,43,45	0
9	GOL	M	1513	6/6	0.69	0.30	18.16	37,39,41,44	6
8	SO4	M	1505[A]	5/5	0.84	0.21	11.33	23,27,27,27	5
8	SO4	M	1533	5/5	0.99	0.13	11.10	16,20,23,25	0
8	SO4	M	1503	5/5	0.99	0.13	9.50	19,20,26,27	5
9	GOL	M	1510[B]	6/6	0.94	0.11	5.71	9,13,16,16	2
9	GOL	M	1510[A]	6/6	0.94	0.11	5.64	9,13,15,16	2
6	GOX	M	999	13/13	0.96	0.07	2.91	11,12,15,19	0
2	NAG	M	901	14/15	0.74	0.16	2.09	27,30,35,38	0
9	GOL	M	1511	6/6	0.97	0.07	1.29	11,16,18,25	6
2	NAG	M	991	14/15	0.73	0.34	-	30,35,44,46	0
9	GOL	M	1512	6/6	0.77	0.17	-	28,29,30,33	6
2	NAG	M	971	14/15	0.61	0.48	-	53,61,64,66	0
8	SO4	M	1504	5/5	0.95	0.12	-	18,24,30,30	5
8	SO4	M	1507	5/5	0.91	0.16	-	25,25,29,30	5
8	SO4	M	1506	5/5	0.99	0.05	-	14,14,22,24	0
2	NAG	M	931	14/15	0.70	0.28	-	31,40,45,47	0
7	ZN	M	1502	1/1	1.00	0.03	-	9,9,9,9	1
2	NAG	M	911	14/15	0.85	0.17	-	22,24,28,28	0
8	SO4	M	1505[B]	5/5	0.84	0.21	-	37,39,40,41	5
8	SO4	M	1508	5/5	0.73	0.22	-	21,28,30,31	5
8	SO4	M	1509	5/5	0.88	0.17	-	34,36,37,37	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.