



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:18 AM GMT

PDB ID : 3E65  
Title : Murine INOS dimer with HEME, pterin and inhibitor AR-C120011  
Authors : Rosenfeld, R.J.; Garcin, E.D.; Getzoff, E.D.  
Deposited on : 2008-08-14  
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1269 (2.04-2.04)
Ramachandran outliers	100387	1258 (2.04-2.04)
Sidechain outliers	100360	1258 (2.04-2.04)
RSRZ outliers	91569	1194 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	<div> <div>5%</div> <div>84%</div> <div>12%</div> <div>••</div> </div>
1	B	433	<div> <div>5%</div> <div>86%</div> <div>11%</div> <div>•</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	H4B	A	902	X	-	-	-
3	H4B	B	902	X	-	-	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	XXZ	B	907	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7139 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, inducible.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	417	Total	C	N	O	S	0	0	0
			3395	2177	584	614	20			
1	B	421	Total	C	N	O	S	0	0	0
			3421	2193	589	618	21			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula:  $C_9H_{15}N_5O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 1-[4-(AMINOMETHYL)BENZOYL]-5'-FLUORO-1'H-SPIRO[PIPERIDINE-4,2'-QUINAZOLIN]-4'-AMINE (three-letter code: XXZ) (formula: C<sub>20</sub>H<sub>22</sub>FN<sub>5</sub>O).



*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	F	N	O	0	0
			27	20	1	5	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	75	Total	O	0	0
			75	75		
5	B	74	Total	O	0	0
			74	74		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, inducible



- Molecule 1: Nitric oxide synthase, inducible



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	213.94Å 213.94Å 117.03Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.98 – 2.05 39.95 – 2.05	Depositor EDS
% Data completeness (in resolution range)	95.5 (19.98-2.05) 95.3 (39.95-2.05)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.15 (at 2.05Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.249 , 0.271 (Not available) , (Not available)	Depositor DCC
$R_{free}$ test set	NotAvailable	DCC
Wilson B-factor (Å <sup>2</sup> )	36.7	Xtriage
Anisotropy	0.595	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 36.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Outliers	0 of 94091 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7139	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, H4B, XXZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/3494	0.61	2/4751 (0.0%)
1	B	0.36	0/3521	0.58	1/4787 (0.0%)
All	All	0.37	0/7015	0.60	3/9538 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	102	PHE	N-CA-C	6.20	127.74	111.00
1	A	368	MET	N-CA-C	-5.06	97.33	111.00
1	B	368	MET	N-CA-C	-5.05	97.36	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	108	SER	Mainchain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3395	0	3281	34	0
1	B	3421	0	3312	30	0
2	A	43	0	30	0	0
2	B	43	0	30	0	0
3	A	17	0	14	0	0
3	B	17	0	14	0	0
4	A	27	0	22	2	0
4	B	27	0	22	1	0
5	A	75	0	0	2	0
5	B	74	0	0	0	0
All	All	7139	0	6725	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:215:GLN:HE21	1:B:219:GLN:HE21	1.23	0.86
1:A:81:ILE:HD11	1:A:475:LEU:HD13	1.62	0.81
1:B:81:ILE:HD11	1:B:475:LEU:HD13	1.62	0.79
1:A:215:GLN:HE21	1:A:219:GLN:HE21	1.33	0.76
1:A:83:ASN:ND2	1:A:85:GLY:H	1.85	0.74
1:A:366:TRP:HE1	1:A:428:MET:HE2	1.52	0.73
1:B:215:GLN:HE21	1:B:219:GLN:NE2	1.89	0.68
1:B:194:CYS:HB3	1:B:197:ARG:HD2	1.75	0.68
1:A:366:TRP:HE1	1:A:428:MET:CE	2.10	0.65
1:A:83:ASN:HD22	1:A:85:GLY:H	1.43	0.64
1:A:153:ILE:O	1:A:157:LEU:HD23	2.02	0.60
1:A:108:SER:O	1:A:110:LEU:HD12	2.01	0.60
1:A:108:SER:O	1:A:110:LEU:CD1	2.50	0.60
1:B:144:TYR:O	1:B:147:SER:HB3	2.02	0.59
1:A:188:TRP:CE3	1:A:200:TRP:HA	2.40	0.57
1:A:465:SER:O	1:A:471:HIS:HE1	1.89	0.56
1:B:104:CYS:SG	1:B:109:CYS:HA	2.47	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:442:GLN:HE21	1:A:446:ARG:HH12	1.56	0.54
1:B:188:TRP:CE3	1:B:200:TRP:HA	2.42	0.54
1:A:77:GLN:O	1:A:96:HIS:HE1	1.90	0.54
1:B:81:ILE:HD11	1:B:475:LEU:CD1	2.37	0.52
1:A:209:ARG:O	1:A:242:PRO:HG3	2.12	0.49
1:A:246:ASP:OD1	1:A:249:HIS:HD2	1.95	0.49
1:B:101:ASP:CG	1:B:102:PHE:H	2.17	0.49
1:B:194:CYS:O	1:B:197:ARG:HD3	2.14	0.48
1:B:134:LEU:HB3	1:B:135:PRO:HD3	1.96	0.48
1:A:134:LEU:HB3	1:A:135:PRO:HD3	1.96	0.48
1:A:460:PRO:HG3	5:A:1444:HOH:O	2.14	0.47
1:B:246:ASP:OD1	1:B:249:HIS:HD2	1.98	0.47
4:B:907:XXZ:H111	4:B:907:XXZ:CE1	2.45	0.46
1:B:215:GLN:NE2	1:B:219:GLN:HE21	2.02	0.46
1:A:251:PHE:O	1:A:360:ALA:HB2	2.16	0.46
1:A:195:ILE:HD12	5:A:1443:HOH:O	2.15	0.46
1:B:253:LEU:HD22	1:B:347:ALA:HB1	1.99	0.45
1:B:137:ALA:O	1:B:141:ILE:HG12	2.16	0.45
1:B:77:GLN:O	1:B:96:HIS:HE1	2.00	0.45
1:A:81:ILE:HD11	1:A:475:LEU:CD1	2.40	0.44
1:B:215:GLN:O	1:B:219:GLN:HG3	2.18	0.44
1:A:243:GLN:HB3	1:A:358:PHE:CE2	2.52	0.44
1:A:83:ASN:C	1:A:83:ASN:HD22	2.21	0.44
1:B:252:ARG:NH2	1:B:489:PRO:HD3	2.33	0.43
1:B:132:GLU:O	1:B:135:PRO:HD2	2.19	0.43
1:B:103:THR:HG22	1:B:116:PRO:HB2	1.99	0.43
1:B:239:THR:O	1:B:361:CYS:HA	2.19	0.43
4:A:906:XXZ:H111	4:A:906:XXZ:CE1	2.48	0.42
1:B:190:ASN:O	1:B:192:PRO:HD3	2.18	0.42
1:B:188:TRP:CZ3	1:B:200:TRP:HA	2.54	0.42
1:B:243:GLN:HB3	1:B:358:PHE:CE2	2.54	0.42
1:A:83:ASN:HD22	1:A:85:GLY:N	2.13	0.42
1:A:188:TRP:CZ3	1:A:200:TRP:HA	2.54	0.42
1:A:254:TRP:HB2	1:A:302:VAL:HB	2.01	0.42
1:B:290:LYS:HA	1:B:291:PRO:HD2	1.96	0.42
1:A:388:GLU:O	1:A:392:ARG:HG3	2.19	0.42
1:A:382:ARG:NH2	4:A:906:XXZ:HD2	2.35	0.41
1:A:190:ASN:O	1:A:192:PRO:HD3	2.20	0.41
1:B:361:CYS:SG	1:B:361:CYS:O	2.79	0.41
1:A:469:VAL:HG13	1:A:474:MET:CE	2.50	0.41
1:B:110:LEU:N	1:B:110:LEU:HD12	2.35	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:254:TRP:HB2	1:B:302:VAL:HB	2.02	0.41
1:B:103:THR:HG23	1:B:118:SER:OG	2.21	0.41
1:A:488:GLU:HA	1:A:489:PRO:HD2	1.87	0.41
1:B:460:PRO:HG2	1:B:467:THR:HG21	2.03	0.41
1:A:189:ARG:HD2	1:A:200:TRP:CE3	2.56	0.41
1:A:446:ARG:HH21	1:A:446:ARG:HG3	1.85	0.41
1:A:252:ARG:NH2	1:A:489:PRO:HD3	2.36	0.40
1:A:239:THR:O	1:A:361:CYS:HA	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	413/433 (95%)	396 (96%)	16 (4%)	1 (0%)	52	43
1	B	419/433 (97%)	398 (95%)	20 (5%)	1 (0%)	52	43
All	All	832/866 (96%)	794 (95%)	36 (4%)	2 (0%)	52	43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	105	LYS
1	A	100	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	363/381 (95%)	357 (98%)	6 (2%)	68	65
1	B	366/381 (96%)	362 (99%)	4 (1%)	80	79
All	All	729/762 (96%)	719 (99%)	10 (1%)	74	72

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83	ASN
1	A	103	THR
1	A	109	CYS
1	A	223	ARG
1	A	334	GLN
1	A	475	LEU
1	B	223	ARG
1	B	295	ARG
1	B	334	GLN
1	B	475	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	83	ASN
1	A	96	HIS
1	A	215	GLN
1	A	249	HIS
1	A	265	GLN
1	A	421	GLN
1	A	442	GLN
1	A	471	HIS
1	B	96	HIS
1	B	219	GLN
1	B	249	HIS
1	B	421	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	HEM	A	901	1	30,50,50	3.13	9 (30%)	24,82,82	2.26	8 (33%)
3	H4B	A	902	-	13,18,18	2.32	2 (15%)	11,26,26	2.35	5 (45%)
4	XXZ	A	906	-	26,30,30	2.94	4 (15%)	36,44,44	1.71	8 (22%)
2	HEM	B	901	1	30,50,50	3.20	10 (33%)	24,82,82	2.23	10 (41%)
3	H4B	B	902	-	13,18,18	2.30	2 (15%)	11,26,26	2.43	5 (45%)
4	XXZ	B	907	-	26,30,30	2.87	5 (19%)	36,44,44	1.69	8 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	901	1	-	0/10/54/54	0/0/8/8
3	H4B	A	902	-	1/1/3/5	0/8/17/17	0/2/2/2
4	XXZ	A	906	-	-	0/10/36/36	0/3/4/4
2	HEM	B	901	1	-	0/10/54/54	0/0/8/8
3	H4B	B	902	-	1/1/3/5	0/8/17/17	0/2/2/2
4	XXZ	B	907	-	-	0/10/36/36	0/3/4/4

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	901	HEM	C3B-C4B	-9.58	1.43	1.51
2	A	901	HEM	C2D-C3D	-6.98	1.33	1.54
2	B	901	HEM	C3B-CAB	-6.91	1.38	1.51
2	A	901	HEM	C3B-CAB	-6.88	1.38	1.51
2	A	901	HEM	C3D-C4D	-6.83	1.42	1.51
2	A	901	HEM	C3B-C4B	-6.55	1.46	1.51
3	A	902	H4B	C6-N5	-6.16	1.33	1.45
3	B	902	H4B	C6-N5	-6.13	1.33	1.45
2	B	901	HEM	C2D-C3D	-6.07	1.36	1.54
2	A	901	HEM	C3C-CAC	-5.87	1.40	1.51
2	B	901	HEM	C3C-CAC	-5.66	1.40	1.51
2	B	901	HEM	C3D-C4D	-5.38	1.44	1.51
3	B	902	H4B	C7-N8	-5.13	1.39	1.46
3	A	902	H4B	C7-N8	-5.04	1.39	1.46
2	A	901	HEM	C2C-C1C	-4.94	1.43	1.52
2	B	901	HEM	C2C-C1C	-3.82	1.45	1.52
2	A	901	HEM	C2D-C1D	-2.86	1.42	1.51
2	A	901	HEM	C2B-C1B	-2.75	1.42	1.51
2	B	901	HEM	C2B-C1B	-2.62	1.43	1.51
2	B	901	HEM	C2D-C1D	-2.08	1.45	1.51
4	A	906	XXZ	CE1-CZ	2.08	1.42	1.39
4	B	907	XXZ	C15-N12	2.23	1.39	1.34
4	B	907	XXZ	CE1-CZ	2.27	1.43	1.39
2	B	901	HEM	C1C-NC	3.19	1.40	1.36
2	A	901	HEM	C1C-NC	3.34	1.40	1.36
2	B	901	HEM	C4C-NC	3.76	1.40	1.36
4	B	907	XXZ	C7-CG	4.43	1.52	1.46
4	A	906	XXZ	C7-CG	4.59	1.52	1.46
4	A	906	XXZ	CG-N20	6.78	1.36	1.28
4	B	907	XXZ	CG-N20	6.83	1.37	1.28
4	B	907	XXZ	C6-N1	10.84	1.55	1.39
4	A	906	XXZ	C6-N1	11.34	1.56	1.39

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	906	XXZ	C10-C5-N1	-4.58	102.26	110.09
4	B	907	XXZ	C10-C5-N1	-4.49	102.42	110.09
4	B	907	XXZ	C6-C7-CG	-3.29	114.40	117.20
4	A	906	XXZ	C7-C6-N1	-3.24	116.06	119.13
4	B	907	XXZ	C14-C5-C10	-3.18	106.83	109.95
4	A	906	XXZ	C14-C5-C10	-3.14	106.87	109.95
4	A	906	XXZ	C6-C7-CG	-2.75	114.86	117.20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	907	XXZ	C13-C14-C5	-2.52	108.88	112.68
3	B	902	H4B	N3-C2-N1	-2.47	121.49	125.53
4	A	906	XXZ	C13-C14-C5	-2.40	109.05	112.68
3	A	902	H4B	N3-C2-N1	-2.33	121.72	125.53
4	B	907	XXZ	C7-C6-N1	-2.11	117.12	119.13
2	B	901	HEM	C3C-CAC-CBC	2.08	127.65	124.46
2	B	901	HEM	CAA-C2A-C1A	2.09	129.28	127.01
4	B	907	XXZ	CE1-CZ-C18	2.15	121.79	118.60
2	A	901	HEM	CMD-C2D-C3D	2.26	124.33	114.35
4	A	906	XXZ	CE1-CZ-C18	2.26	121.96	118.60
2	B	901	HEM	CMD-C2D-C3D	2.29	124.49	114.35
4	B	907	XXZ	CD2-C8-CD1	2.31	121.84	118.13
4	A	906	XXZ	CD2-C8-CD1	2.36	121.92	118.13
2	B	901	HEM	C3B-CAB-CBB	2.47	128.25	124.46
2	B	901	HEM	C3B-C4B-CHC	2.63	126.87	123.16
3	A	902	H4B	C4-C4A-C8A	2.68	116.99	114.56
3	B	902	H4B	C4-C4A-C8A	2.75	117.05	114.56
3	A	902	H4B	C2-N1-C8A	2.86	120.96	114.54
4	A	906	XXZ	C7-CG-N21	2.89	123.11	120.13
3	B	902	H4B	C2-N1-C8A	2.96	121.19	114.54
2	B	901	HEM	CMB-C2B-C3B	3.16	124.42	116.53
4	B	907	XXZ	C7-CG-N21	3.24	123.48	120.13
2	A	901	HEM	C3B-C4B-CHC	3.31	127.82	123.16
2	A	901	HEM	C3C-CAC-CBC	3.35	129.59	124.46
3	B	902	H4B	C4-N3-C2	3.37	120.61	115.94
3	A	902	H4B	C4-N3-C2	3.40	120.66	115.94
2	A	901	HEM	C2D-C3D-C4D	3.41	107.28	101.50
2	B	901	HEM	C2D-C3D-C4D	3.57	107.55	101.50
2	B	901	HEM	CAD-C3D-C4D	3.60	125.17	112.47
2	A	901	HEM	CMB-C2B-C3B	3.65	125.64	116.53
2	A	901	HEM	CAD-C3D-C4D	4.01	126.63	112.47
2	A	901	HEM	CAD-C3D-C2D	4.33	125.66	113.22
3	A	902	H4B	C7-C6-N5	4.57	119.94	110.45
3	B	902	H4B	C7-C6-N5	4.75	120.31	110.45
2	B	901	HEM	CAD-C3D-C2D	4.78	126.95	113.22
2	B	901	HEM	CMC-C2C-C3C	5.00	129.00	116.53
2	A	901	HEM	CMC-C2C-C3C	5.06	129.15	116.53

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	902	H4B	C6

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atom
3	B	902	H4B	C6

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	906	XXZ	2	0
4	B	907	XXZ	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	417/433 (96%)	0.21	20 (4%) 34 40	29, 43, 68, 88	0
1	B	421/433 (97%)	0.23	22 (5%) 31 36	27, 41, 67, 89	0
All	All	838/866 (96%)	0.22	42 (5%) 32 37	27, 42, 68, 89	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	102	PHE	15.0
1	B	102	PHE	15.0
1	B	99	THR	11.0
1	B	103	THR	10.9
1	A	103	THR	10.0
1	B	106	SER	9.8
1	B	108	SER	9.6
1	B	101	ASP	7.3
1	A	101	ASP	7.1
1	A	108	SER	7.0
1	A	100	SER	6.8
1	A	110	LEU	5.5
1	A	497	ASN	5.3
1	B	497	ASN	4.9
1	B	104	CYS	4.8
1	B	494	ILE	4.6
1	A	111	GLY	4.2
1	B	105	LYS	4.2
1	B	107	LYS	4.2
1	B	109	CYS	3.7
1	A	109	CYS	3.7
1	B	130	LEU	3.5
1	B	149	LYS	3.4
1	B	78	TYR	3.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	110	LEU	3.3
1	A	78	TYR	3.2
1	A	149	LYS	3.2
1	A	494	ILE	3.1
1	A	495	TRP	3.1
1	B	150	GLU	3.0
1	A	150	GLU	3.0
1	A	446	ARG	2.8
1	B	459	VAL	2.7
1	A	91	HIS	2.6
1	A	457	TRP	2.4
1	A	290	LYS	2.3
1	B	267	PRO	2.3
1	B	112	SER	2.2
1	A	496	GLN	2.2
1	A	334	GLN	2.1
1	B	148	PHE	2.1
1	B	456	ILE	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	XXZ	B	907	27/27	0.94	0.20	2.96	28,31,34,36	0
4	XXZ	A	906	27/27	0.93	0.20	1.94	26,29,35,38	0
2	HEM	B	901	43/43	0.97	0.18	1.51	25,29,32,39	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	HEM	A	901	43/43	0.97	0.18	1.05	28,31,34,39	0
3	H4B	A	902	17/17	0.96	0.18	0.50	30,33,36,36	0
3	H4B	B	902	17/17	0.96	0.18	0.41	24,30,33,35	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.