



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:02 PM GMT

PDB ID : 4E7I  
Title : PFV intasome freeze-trapped prior to 3'-processing, Mn-bound form (UI-Mn)  
Authors : Hare, S.; Cherepanov, P.  
Deposited on : 2012-03-17  
Resolution : 2.53 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4549 (2.58-2.50)
Clashscore	102246	5292 (2.58-2.50)
Ramachandran outliers	100387	5194 (2.58-2.50)
Sidechain outliers	100360	5196 (2.58-2.50)
RSRZ outliers	91569	4561 (2.58-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	<div> <div>5%</div> <div>82%</div> <div>11%</div> <div>7%</div> </div>
1	B	395	<div> <div>3%</div> <div>40%</div> <div>7%</div> <div>53%</div> </div>
2	C	19	<div> <div>21%</div> <div>63%</div> <div>16%</div> </div>
3	D	19	<div> <div>47%</div> <div>47%</div> <div>5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	GOL	A	407	-	-	-	X
7	GOL	A	409	-	-	-	X
7	GOL	B	404	-	-	-	X
7	GOL	D	102	-	-	-	X
8	MES	A	412	-	-	-	X
9	HEZ	B	403	-	-	-	X

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 5483 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pro-Pol polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	368	Total	C	N	O	S	0	0	0
			2922	1873	515	530	4			
1	B	184	Total	C	N	O	S	0	0	0
			1423	922	233	267	1			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP P14350
A	-1	PRO	-	EXPRESSION TAG	UNP P14350
A	0	GLY	-	EXPRESSION TAG	UNP P14350
A	217	SER	GLY	VARIANT	UNP P14350
A	218	GLY	SER	VARIANT	UNP P14350
B	-2	GLY	-	EXPRESSION TAG	UNP P14350
B	-1	PRO	-	EXPRESSION TAG	UNP P14350
B	0	GLY	-	EXPRESSION TAG	UNP P14350
B	217	SER	GLY	VARIANT	UNP P14350
B	218	GLY	SER	VARIANT	UNP P14350

- Molecule 2 is a DNA chain called DNA (5'-D(\*AP\*TP\*TP\*GP\*TP\*CP\*AP\*TP\*GP\*GP\*AP\*AP\*TP\*TP\*TP\*CP\*GP\*CP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	19	Total	C	N	O	P	0	0	0
			387	187	68	114	18			

- Molecule 3 is a DNA chain called DNA (5'-D(\*TP\*GP\*CP\*GP\*AP\*AP\*AP\*TP\*TP\*CP\*CP\*AP\*TP\*GP\*AP\*CP\*AP\*AP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	19	Total	C	N	O	P	0	0	0
			386	186	72	110	18			

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		

- Molecule 5 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mn	0	0
			1	1		
5	A	2	Total	Mn	0	0
			2	2		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		
7	D	1	Total	C	O	0	0
			6	3	3		
7	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 9 is HEXANE-1,6-DIOL (three-letter code: HEZ) (formula:  $C_6H_{14}O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			8	6	2		



- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	160	Total 160	O 160	0	0
10	B	50	Total 50	O 50	0	0
10	C	35	Total 35	O 35	0	0
10	D	22	Total 22	O 22	0	0





- Molecule 1: Pro-Pol polypeptide



- Molecule 3: DNA (5'-D(\*TP\*GP\*CP\*GP\*AP\*AP\*AP\*TP\*TP\*CP\*CP\*AP\*TP\*GP\*AP\*CP\*AP\*AP\*T)-3')

Chain D:  47% 47% 5%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	160.15Å 160.15Å 124.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.84 – 2.53 38.84 – 2.53	Depositor EDS
% Data completeness (in resolution range)	99.8 (38.84-2.53) 99.8 (38.84-2.53)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.02 (at 2.54Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, $R_{free}$	0.183 , 0.211 0.179 , 0.209	Depositor DCC
$R_{free}$ test set	2746 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	59.0	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 44.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 54244 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5483	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.41% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, MN, HEZ, SO4, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.50	0/3001	0.65	0/4096
1	B	0.51	0/1463	0.62	0/2003
2	C	0.98	0/433	1.94	17/667 (2.5%)
3	D	1.00	0/433	1.98	12/666 (1.8%)
All	All	0.61	0/5330	1.01	29/7432 (0.4%)

There are no bond length outliers.

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	7	DA	O4'-C1'-N9	-13.30	98.69	108.00
3	D	17	DA	O4'-C1'-N9	12.64	116.84	108.00
3	D	12	DA	O4'-C1'-N9	-10.30	100.79	108.00
2	C	15	DT	O4'-C1'-N1	-9.86	101.10	108.00
3	D	13	DT	O4'-C1'-N1	-9.10	101.63	108.00
3	D	18	DA	O4'-C1'-N9	-9.00	101.70	108.00
2	C	8	DT	O4'-C4'-C3'	-8.68	100.79	106.00
3	D	1	DT	O4'-C1'-N1	8.42	113.89	108.00
3	D	10	DC	O4'-C1'-N1	-7.31	102.88	108.00
2	C	5	DT	O4'-C1'-N1	-7.22	102.94	108.00
3	D	17	DA	C1'-O4'-C4'	-6.99	103.11	110.10
3	D	1	DT	C3'-C2'-C1'	-6.72	94.44	102.50
2	C	2	DT	N3-C2-O2	-6.61	118.33	122.30
2	C	14	DT	O4'-C1'-N1	-6.43	103.50	108.00
2	C	17	DG	O4'-C1'-N9	-6.41	103.51	108.00
2	C	11	DA	O4'-C1'-N9	-6.33	103.57	108.00
2	C	19	DA	O4'-C1'-C2'	-6.26	100.89	105.90
2	C	7	DA	O4'-C1'-N9	-6.18	103.67	108.00
2	C	12	DA	O4'-C1'-N9	-5.90	103.87	108.00
2	C	5	DT	N1-C2-O2	-5.73	118.52	123.10

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	4	DG	O4'-C1'-N9	-5.58	104.10	108.00
3	D	9	DT	O4'-C1'-N1	-5.54	104.12	108.00
2	C	16	DC	O4'-C1'-N1	-5.51	104.14	108.00
2	C	18	DC	N1-C2-O2	-5.48	115.61	118.90
2	C	15	DT	C5-C4-O4	-5.40	121.12	124.90
3	D	1	DT	C1'-O4'-C4'	-5.39	104.70	110.10
2	C	1	DA	N9-C1'-C2'	5.21	122.49	112.60
2	C	15	DT	N3-C4-O4	5.08	122.95	119.90
2	C	2	DT	N1-C2-O2	5.08	127.16	123.10

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2922	0	2952	26	0
1	B	1423	0	1384	14	0
2	C	387	0	218	6	0
3	D	386	0	216	2	0
4	A	1	0	0	0	0
5	A	2	0	0	0	0
5	B	1	0	0	0	0
6	A	15	0	0	0	0
6	B	5	0	0	1	0
7	A	30	0	40	4	0
7	B	6	0	8	2	0
7	C	6	0	8	0	0
7	D	12	0	16	0	0
8	A	12	0	13	4	0
9	B	8	0	14	1	0
10	A	160	0	0	4	1
10	B	50	0	0	0	0
10	C	35	0	0	0	0
10	D	22	0	0	1	0
All	All	5483	0	4869	51	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:1:DA:H4'	2:C:2:DT:H5'	1.43	1.01
7:A:408:GOL:HO1	7:A:408:GOL:HO3	0.92	0.89
1:A:137:GLN:H	7:A:406:GOL:H2	1.46	0.81
1:B:250:GLN:HE22	9:B:403:HEZ:H22	1.52	0.74
1:B:219:LYS:HE2	1:B:261:LEU:HD12	1.71	0.72
1:A:161:PRO:HG3	8:A:412:MES:H52	1.73	0.69
1:A:98:GLN:OE1	10:A:548:HOH:O	2.11	0.68
2:C:1:DA:H1'	2:C:2:DT:H73	1.83	0.61
1:A:14:GLN:HB3	1:A:21:ILE:HD11	1.86	0.58
1:A:358:LEU:N	1:A:359:GLY:HA2	2.18	0.57
1:B:222:ARG:NE	6:B:402:SO4:O4	2.33	0.56
8:A:412:MES:H61	10:A:642:HOH:O	2.05	0.56
1:A:69:ARG:HD2	1:A:70:GLU:OE2	2.05	0.56
1:A:163:THR:H	7:A:409:GOL:H32	1.71	0.55
1:B:286:LEU:HD12	1:B:286:LEU:H	1.71	0.55
3:D:18:DA:OP2	10:D:203:HOH:O	2.18	0.54
2:C:1:DA:H8	2:C:2:DT:H72	1.71	0.54
1:A:73:LEU:HD22	1:A:86:ARG:CZ	2.42	0.49
1:B:141:TYR:CE2	1:B:161:PRO:HD3	2.48	0.49
1:A:185:ASP:HB2	3:D:19:DT:OP2	2.13	0.48
1:B:127:ILE:HA	1:B:144:VAL:O	2.13	0.48
1:A:358:LEU:H	1:A:359:GLY:HA2	1.78	0.48
1:A:295:LEU:HD21	1:A:299:ARG:NH2	2.28	0.48
7:A:406:GOL:O1	7:A:406:GOL:O3	2.32	0.47
1:A:73:LEU:O	1:A:73:LEU:HG	2.14	0.46
1:B:231:LEU:HD23	1:B:231:LEU:HA	1.79	0.46
1:A:111:PRO:O	1:A:350:ARG:HD3	2.15	0.46
1:B:264:THR:OG1	1:B:267:GLN:HG3	2.15	0.46
1:B:287:THR:O	1:B:291:GLU:HG3	2.16	0.46
1:B:267:GLN:HB3	7:B:404:GOL:H12	1.99	0.45
1:A:315:TRP:CE2	1:A:371:PRO:HD3	2.52	0.45
1:B:263:TYR:CE1	7:B:404:GOL:H32	2.52	0.44
1:A:219:LYS:HE3	2:C:6:DC:OP1	2.16	0.44
1:A:25:PRO:HB2	1:A:28:TYR:HD2	1.83	0.44
1:A:325:GLU:HA	1:A:369:LEU:HD23	2.00	0.44
1:A:292:LEU:O	1:A:296:GLN:HG3	2.18	0.44
1:A:161:PRO:O	1:A:189:ALA:HB2	2.17	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:341:SER:HB2	1:A:355:LEU:O	2.17	0.44
8:A:412:MES:H22	8:A:412:MES:H71	1.84	0.42
1:B:284:LEU:HA	1:B:284:LEU:HD12	1.71	0.42
1:A:219:LYS:NZ	10:A:511:HOH:O	2.51	0.42
1:A:156:TYR:OH	1:A:173:LEU:HB2	2.20	0.42
2:C:1:DA:C8	2:C:2:DT:H72	2.53	0.42
1:B:258:SER:O	1:B:261:LEU:O	2.38	0.42
1:B:283:THR:C	1:B:285:ASP:H	2.23	0.42
2:C:8:DT:H2'	2:C:9:DG:C8	2.55	0.41
1:A:122:PHE:O	1:A:179:PRO:HA	2.20	0.41
1:A:108:ALA:O	1:A:314:SER:HA	2.20	0.41
1:A:284:LEU:HA	1:A:284:LEU:HD12	1.81	0.41
1:A:322:LEU:HD23	1:A:322:LEU:HA	1.81	0.41
8:A:412:MES:H71	10:A:642:HOH:O	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:A:528:HOH:O	10:A:632:HOH:O[8_554]	2.12	0.08

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	366/395 (93%)	356 (97%)	10 (3%)	0	100	100
1	B	182/395 (46%)	168 (92%)	11 (6%)	3 (2%)	12	20
All	All	548/790 (69%)	524 (96%)	21 (4%)	3 (0%)	34	54

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	216	SER
1	B	260	VAL
1	B	298	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	327/354 (92%)	318 (97%)	9 (3%)	51	76
1	B	154/354 (44%)	148 (96%)	6 (4%)	39	64
All	All	481/708 (68%)	466 (97%)	15 (3%)	47	73

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	ASP
1	A	55	GLN
1	A	149	MET
1	A	246	LEU
1	A	284	LEU
1	A	306	SER
1	A	312	SER
1	A	332	SER
1	A	364	VAL
1	B	149	MET
1	B	215	GLN
1	B	274	SER
1	B	282	ASP
1	B	286	LEU
1	B	288	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.



### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 19 ligands modelled in this entry, 4 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	SO4	A	404	-	4,4,4	0.18	0	6,6,6	0.27	0
6	SO4	A	405	-	4,4,4	0.16	0	6,6,6	0.25	0
7	GOL	A	406	-	5,5,5	0.31	0	5,5,5	0.73	0
7	GOL	A	407	-	5,5,5	0.36	0	5,5,5	0.68	0
7	GOL	A	408	-	5,5,5	0.34	0	5,5,5	0.86	0
7	GOL	A	409	-	5,5,5	0.45	0	5,5,5	0.90	0
6	SO4	A	410	-	4,4,4	0.28	0	6,6,6	0.43	0
7	GOL	A	411	-	5,5,5	0.34	0	5,5,5	0.23	0
8	MES	A	412	-	11,12,12	1.44	3 (27%)	14,16,16	1.35	2 (14%)
6	SO4	B	402	-	4,4,4	0.22	0	6,6,6	0.21	0
9	HEZ	B	403	-	7,7,7	0.31	0	6,6,6	0.97	0
7	GOL	B	404	-	5,5,5	0.44	0	5,5,5	0.47	0
7	GOL	C	101	-	5,5,5	0.30	0	5,5,5	0.83	0
7	GOL	D	101	-	5,5,5	0.36	0	5,5,5	0.43	0
7	GOL	D	102	-	5,5,5	0.24	0	5,5,5	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	SO4	A	404	-	-	0/0/0/0	0/0/0/0
6	SO4	A	405	-	-	0/0/0/0	0/0/0/0
7	GOL	A	406	-	-	0/4/4/4	0/0/0/0
7	GOL	A	407	-	-	0/4/4/4	0/0/0/0
7	GOL	A	408	-	-	0/4/4/4	0/0/0/0
7	GOL	A	409	-	-	0/4/4/4	0/0/0/0
6	SO4	A	410	-	-	0/0/0/0	0/0/0/0
7	GOL	A	411	-	-	0/4/4/4	0/0/0/0
8	MES	A	412	-	-	0/6/14/14	0/1/1/1
6	SO4	B	402	-	-	0/0/0/0	0/0/0/0
9	HEZ	B	403	-	-	0/5/5/5	0/0/0/0
7	GOL	B	404	-	-	0/4/4/4	0/0/0/0
7	GOL	C	101	-	-	0/4/4/4	0/0/0/0
7	GOL	D	101	-	-	0/4/4/4	0/0/0/0
7	GOL	D	102	-	-	0/4/4/4	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	412	MES	O1S-S	2.63	1.53	1.45
8	A	412	MES	O2S-S	2.73	1.53	1.45
8	A	412	MES	O3S-S	2.73	1.53	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	412	MES	O2S-S-C8	2.91	109.39	106.91
8	A	412	MES	O1S-S-C8	3.01	109.48	106.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	406	GOL	2	0
7	A	408	GOL	1	0
7	A	409	GOL	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	412	MES	4	0
6	B	402	SO4	1	0
9	B	403	HEZ	1	0
7	B	404	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	368/395 (93%)	0.03	20 (5%) 29 34	38, 51, 91, 126	0
1	B	184/395 (46%)	-0.03	13 (7%) 19 22	41, 58, 124, 141	0
2	C	19/19 (100%)	-0.27	0 100 100	42, 56, 84, 97	0
3	D	19/19 (100%)	-0.59	0 100 100	43, 51, 91, 96	0
All	All	590/828 (71%)	-0.02	33 (5%) 28 33	38, 53, 103, 141	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	8	LEU	7.6
1	B	280	ASN	4.7
1	B	260	VAL	4.3
1	B	294	LEU	3.9
1	A	15	LEU	3.8
1	A	17	GLN	3.5
1	B	279	ALA	3.3
1	A	357	HIS	3.3
1	B	293	SER	3.2
1	B	297	GLU	2.9
1	B	292	LEU	2.7
1	B	296	GLN	2.6
1	B	116	ASP	2.6
1	A	153	THR	2.6
1	A	32	LEU	2.6
1	A	265	PRO	2.5
1	A	16	LEU	2.5
1	A	358	LEU	2.5
1	A	19	HIS	2.5
1	A	9	ASP	2.5
1	A	21	ILE	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	12	LEU	2.4
1	B	298	ILE	2.4
1	B	295	LEU	2.3
1	A	254	ASN	2.3
1	B	281	GLN	2.1
1	A	253	LEU	2.1
1	A	13	ASP	2.1
1	A	146	VAL	2.1
1	B	261	LEU	2.1
1	A	256	THR	2.1
1	A	145	VAL	2.0
1	A	359	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
7	GOL	A	407	6/6	0.83	0.26	12.39	71,76,93,95	0
9	HEZ	B	403	8/8	0.86	0.26	4.24	65,73,79,79	0
7	GOL	A	409	6/6	0.86	0.22	4.08	65,77,81,82	0
7	GOL	B	404	6/6	0.92	0.35	3.54	64,72,76,90	0
7	GOL	D	102	6/6	0.90	0.22	3.26	67,70,85,92	0
8	MES	A	412	12/12	0.88	0.28	2.85	48,74,89,96	12
6	SO4	A	405	5/5	0.92	0.21	1.96	57,68,72,80	5
6	SO4	A	410	5/5	0.90	0.22	1.37	64,66,74,82	5
4	ZN	A	401	1/1	1.00	0.13	0.25	47,47,47,47	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
6	SO4	B	402	5/5	0.96	0.20	-0.14	78,78,90,97	0
7	GOL	C	101	6/6	0.97	0.14	-0.36	49,52,60,70	0
5	MN	A	403	1/1	0.99	0.09	-1.54	55,55,55,55	0
7	GOL	A	411	6/6	0.92	0.37	-	76,78,82,91	0
7	GOL	A	406	6/6	0.92	0.13	-	66,73,81,91	0
5	MN	A	402	1/1	1.00	0.12	-	44,44,44,44	0
7	GOL	D	101	6/6	0.88	0.24	-	59,76,78,80	0
5	MN	B	401	1/1	1.00	0.13	-	58,58,58,58	0
6	SO4	A	404	5/5	0.99	0.17	-	66,71,88,106	0
7	GOL	A	408	6/6	0.85	0.24	-	56,64,78,89	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.