



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 24, 2016 – 04:27 PM EDT

PDB ID : 5ECI
Title : Crystal Structure of FIN219-FIP1 complex with JA, ATP and Mg
Authors : Chen, C.Y.; Cheng, Y.S.
Deposited on : 2015-10-20
Resolution : 1.56 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1665 (1.58-1.54)
Clashscore	102246	1014 (1.56-1.56)
Ramachandran outliers	100387	1704 (1.58-1.54)
Sidechain outliers	100360	1702 (1.58-1.54)
RSRZ outliers	91569	1668 (1.58-1.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	575	<div> <div>13%</div> <div>23% 64% 10% ..</div> </div>
1	D	575	<div> <div>13%</div> <div>23% 65% 10% ..</div> </div>
2	B	223	<div> <div>12%</div> <div>27% 60% 8% .</div> </div>
2	C	223	<div> <div>9%</div> <div>30% 60% 6% .</div> </div>
2	E	223	<div> <div>19%</div> <div>20% 63% 13% .</div> </div>
2	F	223	<div> <div>10%</div> <div>34% 55% 7% .</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ATP	A	602	-	-	X	-
4	ATP	D	602	-	-	X	-
6	GSH	B	301	-	-	X	-
6	GSH	E	301	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 17930 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Jasmonic acid-amido synthetase JAR1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	569	Total	C	N	O	S	0	0	0
			4479	2859	748	850	22			
1	D	569	Total	C	N	O	S	0	0	0
			4479	2859	748	850	22			

- Molecule 2 is a protein called Glutathione S-transferase U20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	214	Total	C	N	O	S	0	0	0
			1748	1136	284	323	5			
2	C	214	Total	C	N	O	S	0	0	0
			1748	1136	284	323	5			
2	E	214	Total	C	N	O	S	0	0	0
			1748	1136	284	323	5			
2	F	214	Total	C	N	O	S	0	0	0
			1748	1136	284	323	5			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	HIS	-	expression tag	UNP Q8L7C9
B	-4	HIS	-	expression tag	UNP Q8L7C9
B	-3	HIS	-	expression tag	UNP Q8L7C9
B	-2	HIS	-	expression tag	UNP Q8L7C9
B	-1	HIS	-	expression tag	UNP Q8L7C9
B	0	HIS	-	expression tag	UNP Q8L7C9
C	-5	HIS	-	expression tag	UNP Q8L7C9
C	-4	HIS	-	expression tag	UNP Q8L7C9
C	-3	HIS	-	expression tag	UNP Q8L7C9
C	-2	HIS	-	expression tag	UNP Q8L7C9
C	-1	HIS	-	expression tag	UNP Q8L7C9
C	0	HIS	-	expression tag	UNP Q8L7C9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-5	HIS	-	expression tag	UNP Q8L7C9
E	-4	HIS	-	expression tag	UNP Q8L7C9
E	-3	HIS	-	expression tag	UNP Q8L7C9
E	-2	HIS	-	expression tag	UNP Q8L7C9
E	-1	HIS	-	expression tag	UNP Q8L7C9
E	0	HIS	-	expression tag	UNP Q8L7C9
F	-5	HIS	-	expression tag	UNP Q8L7C9
F	-4	HIS	-	expression tag	UNP Q8L7C9
F	-3	HIS	-	expression tag	UNP Q8L7C9
F	-2	HIS	-	expression tag	UNP Q8L7C9
F	-1	HIS	-	expression tag	UNP Q8L7C9
F	0	HIS	-	expression tag	UNP Q8L7C9

- Molecule 3 is {(1R,2R)-3-oxo-2-[(2Z)-pent-2-en-1-yl]cyclopentyl}acetic acid (three-letter code: JAA) (formula: C₁₂H₁₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			15	12	3		
3	D	1	Total	C	O	0	0
			15	12	3		

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	D	1	Total 31	C 10	N 5	O 13	P 3	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0
5	D	1	Total Mg 1 1	0	0

- Molecule 6 is GLUTATHIONE (three-letter code: GSH) (formula: $\text{C}_{10}\text{H}_{17}\text{N}_3\text{O}_6\text{S}$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	B	1	Total 20	C 10	N 3	O 6	S 1	0	0
6	C	1	Total 20	C 10	N 3	O 6	S 1	0	0
6	E	1	Total 20	C 10	N 3	O 6	S 1	0	0
6	F	1	Total 20	C 10	N 3	O 6	S 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	424	Total O 424 424	0	0
7	B	219	Total O 219 219	0	0
7	C	203	Total O 203 203	0	0
7	D	474	Total O 474 474	0	0
7	E	260	Total O 260 260	0	0
7	F	226	Total O 226 226	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Jasmonic acid-amido synthetase JAR1




- Molecule 1: Jasmonic acid-amido synthetase JAR1







• Molecule 2: Glutathione S-transferase U20




• Molecule 2: Glutathione S-transferase U20



• Molecule 2: Glutathione S-transferase U20





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	53.89Å 53.86Å 195.40Å 92.52° 97.02° 113.58°	Depositor
Resolution (Å)	24.57 – 1.56 24.57 – 1.56	Depositor EDS
% Data completeness (in resolution range)	97.2 (24.57-1.56) 93.8 (24.57-1.56)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 1.56Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692+SVN)	Depositor
R, R_{free}	0.212 , 0.225 0.212 , 0.225	Depositor DCC
R_{free} test set	27489 reflections (9.99%)	DCC
Wilson B-factor (Å ²)	2.0	Xtriage
Anisotropy	0.438	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.58 , 296.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.061 for k,h,-h-k-l 0.018 for -k,-h,l 0.014 for -h,-k,h+k+l	Xtriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	17930	wwPDB-VP
Average B, all atoms (Å ²)	3.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.49 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.9588e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GSH, MG, ATP, JAA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.58	2/4581 (0.0%)	0.85	8/6219 (0.1%)
1	D	0.55	2/4581 (0.0%)	0.80	4/6219 (0.1%)
2	B	0.51	0/1799	0.70	2/2428 (0.1%)
2	C	0.50	0/1799	0.74	1/2428 (0.0%)
2	E	0.55	0/1799	0.75	0/2428
2	F	0.49	0/1799	0.68	0/2428
All	All	0.54	4/16358 (0.0%)	0.78	15/22150 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	D	0	1
2	C	0	1
All	All	0	5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	520	LEU	CG-CD2	-7.42	1.24	1.51
1	A	450	LYS	CD-CE	-6.33	1.35	1.51
1	A	546	LYS	CG-CD	-5.69	1.33	1.52
1	D	522	VAL	CB-CG2	-5.07	1.42	1.52

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	14	MET	CG-SD-CE	-19.82	68.49	100.20
1	A	425	ARG	NE-CZ-NH1	-11.86	114.37	120.30
1	A	546	LYS	CA-CB-CG	10.61	136.74	113.40
1	D	107	ARG	NE-CZ-NH2	-10.02	115.29	120.30
1	D	118	MET	CG-SD-CE	-8.45	86.68	100.20

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	427	LEU	Peptide
1	A	545	PHE	Peptide
1	A	546	LYS	Peptide
2	C	140	GLY	Peptide
1	D	427	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4479	0	4434	668	2
1	D	4479	0	4434	634	5
2	B	1748	0	1704	225	1
2	C	1748	0	1704	191	1
2	E	1748	0	1704	257	2
2	F	1748	0	1704	174	2
3	A	15	0	0	3	0
3	D	15	0	0	1	0
4	A	31	0	7	12	0
4	D	31	0	8	14	0
5	A	1	0	0	0	0
5	D	1	0	0	0	0
6	B	20	0	15	8	0
6	C	20	0	15	0	0
6	E	20	0	15	12	0
6	F	20	0	15	2	0
7	A	424	0	0	115	2
7	B	219	0	0	40	2

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	C	203	0	0	47	4
7	D	474	0	0	107	10
7	E	260	0	0	55	6
7	F	226	0	0	48	5
All	All	17930	0	15759	2054	26

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 64.

The worst 5 of 2054 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:529:ARG:NH1	1:A:533:GLU:OE2	1.81	1.14
1:A:101:SER:HA	1:A:546:LYS:HD3	1.27	1.14
2:B:26:LYS:NZ	2:B:82:ASN:O	1.83	1.12
1:D:176:LYS:NZ	1:D:190:ASP:OD2	1.84	1.11
1:A:437:THR:HG22	1:A:440:ASP:H	1.18	1.06

The worst 5 of 26 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:107:ARG:NH1	1:D:139:ASP:OD2[1_545]	1.73	0.47
1:A:114:THR:OG1	2:F:141:ASP:OD2[1_454]	1.79	0.41
2:E:38:SER:O	2:F:29:GLU:N[1_565]	1.98	0.22
7:D:717:HOH:O	7:E:427:HOH:O[1_565]	2.04	0.16
7:E:605:HOH:O	7:F:538:HOH:O[1_565]	2.04	0.16

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	567/575 (99%)	511 (90%)	36 (6%)	20 (4%)	4 0
1	D	567/575 (99%)	512 (90%)	39 (7%)	16 (3%)	6 0
2	B	212/223 (95%)	193 (91%)	17 (8%)	2 (1%)	21 4
2	C	212/223 (95%)	196 (92%)	16 (8%)	0	100 100
2	E	212/223 (95%)	192 (91%)	14 (7%)	6 (3%)	6 0
2	F	212/223 (95%)	191 (90%)	18 (8%)	3 (1%)	14 1
All	All	1982/2042 (97%)	1795 (91%)	140 (7%)	47 (2%)	7 0

5 of 47 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	368	GLU
1	A	437	THR
1	A	513	LYS
1	A	540	SER
1	A	542	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	499/505 (99%)	435 (87%)	64 (13%)	5 0
1	D	499/505 (99%)	442 (89%)	57 (11%)	7 0
2	B	187/195 (96%)	162 (87%)	25 (13%)	5 0
2	C	187/195 (96%)	174 (93%)	13 (7%)	19 2
2	E	187/195 (96%)	157 (84%)	30 (16%)	3 0
2	F	187/195 (96%)	169 (90%)	18 (10%)	10 0
All	All	1746/1790 (98%)	1539 (88%)	207 (12%)	6 0

5 of 207 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	80	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	133	ASN
2	F	29	GLU
2	C	122	GLU
1	D	88	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	60	ASN
2	C	51	HIS
2	E	110	GLN
2	B	69	ASN
2	B	72	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	JAA	A	601	-	12,15,15	5.70	6 (50%)	12,19,19	4.12	7 (58%)
4	ATP	A	602	5	26,33,33	4.09	15 (57%)	26,52,52	3.98	15 (57%)
6	GSH	B	301	-	13,19,19	1.58	4 (30%)	15,24,24	3.11	4 (26%)
6	GSH	C	301	-	13,19,19	1.60	3 (23%)	15,24,24	3.61	6 (40%)
3	JAA	D	601	-	12,15,15	5.62	7 (58%)	12,19,19	3.75	6 (50%)
4	ATP	D	602	5	26,33,33	3.91	16 (61%)	26,52,52	4.07	10 (38%)
6	GSH	E	301	-	13,19,19	1.66	4 (30%)	15,24,24	4.05	7 (46%)
6	GSH	F	301	-	13,19,19	1.76	3 (23%)	15,24,24	3.11	6 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	JAA	A	601	-	-	0/7/22/22	0/1/1/1
4	ATP	A	602	5	-	0/18/38/38	0/3/3/3
6	GSH	B	301	-	-	0/18/24/24	0/0/0/0
6	GSH	C	301	-	-	0/18/24/24	0/0/0/0
3	JAA	D	601	-	-	0/7/22/22	0/1/1/1
4	ATP	D	602	5	-	0/18/38/38	0/3/3/3
6	GSH	E	301	-	-	0/18/24/24	0/0/0/0
6	GSH	F	301	-	-	0/18/24/24	0/0/0/0

The worst 5 of 58 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	601	JAA	C05-C08	-12.77	1.31	1.52
3	A	601	JAA	C05-C08	-12.50	1.31	1.52
3	A	601	JAA	C06-C04	-11.22	1.23	1.53
3	D	601	JAA	C06-C04	-10.72	1.25	1.53
4	A	602	ATP	C3'-C4'	-9.46	1.27	1.53

The worst 5 of 61 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	E	301	GSH	CA2-CB2-SG2	-12.71	100.03	113.99
4	A	602	ATP	C4'-O4'-C1'	-10.02	99.02	109.64
4	A	602	ATP	N3-C2-N1	-9.92	121.08	128.87
4	D	602	ATP	C4'-O4'-C1'	-9.82	99.23	109.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	301	GSH	CA2-CB2-SG2	-9.72	103.31	113.99

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 51 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	601	JAA	3	0
4	A	602	ATP	12	0
6	B	301	GSH	8	0
3	D	601	JAA	1	0
4	D	602	ATP	14	0
6	E	301	GSH	12	0
6	F	301	GSH	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	569/575 (98%)	1.01	77 (13%) 4 4	3, 3, 3, 16	0
1	D	569/575 (98%)	0.99	74 (13%) 5 4	3, 3, 3, 16	0
2	B	214/223 (95%)	1.02	27 (12%) 5 5	2, 3, 7, 11	0
2	C	214/223 (95%)	0.82	20 (9%) 11 11	3, 3, 3, 4	0
2	E	214/223 (95%)	1.21	42 (19%) 1 1	2, 3, 7, 11	0
2	F	214/223 (95%)	0.80	23 (10%) 8 7	3, 3, 3, 4	0
All	All	1994/2042 (97%)	0.98	263 (13%) 4 4	2, 3, 5, 16	0

The worst 5 of 263 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	77	ALA	6.9
1	A	546	LYS	6.6
2	E	13	SER	5.6
1	D	259	VAL	5.5
1	D	493	CYS	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	JAA	D	601	15/15	0.92	0.14	0.58	3,3,3,3	0
3	JAA	A	601	15/15	0.92	0.15	0.30	3,3,3,3	0
6	GSH	C	301	20/20	0.95	0.11	-0.51	2,3,3,3	0
6	GSH	F	301	20/20	0.95	0.11	-0.70	2,3,3,3	0
6	GSH	B	301	20/20	0.96	0.11	-0.71	2,2,3,3	0
6	GSH	E	301	20/20	0.96	0.10	-0.72	2,2,3,3	0
4	ATP	D	602	31/31	0.96	0.10	-1.38	3,3,3,3	0
4	ATP	A	602	31/31	0.96	0.09	-1.69	3,3,3,3	0
5	MG	A	603	1/1	0.99	0.07	-2.54	3,3,3,3	0
5	MG	D	603	1/1	0.99	0.07	-2.82	3,3,3,3	0

6.5 Other polymers [i](#)

There are no such residues in this entry.