



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:22 AM GMT

PDB ID : 3EET
Title : Crystal structure of putative GntR-family transcriptional regulator
Authors : Chang, C.; Xu, X.; Zheng, H.; Savchenko, A.; Edwards, A.; Joachimiak, A.;
Midwest Center for Structural Genomics (MCSG)
Deposited on : 2008-09-05
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 76%, yellow 76%, yellow 89%, green 89%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 76% 14% • 10% </div> </div>
1	B	272	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 7%, orange 7%, orange 69%, yellow 69%, yellow 86%, green 86%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 7% 69% 17% • 13% </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3903 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative GntR-family transcriptional regulator.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	245	Total	C	N	O	S	Se	0	6	0
			1927	1202	359	358	1	7			
1	B	236	Total	C	N	O	S	Se	0	0	0
			1761	1108	315	330	1	7			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MSE	-	expression tag	UNP Q82IF8
A	-20	GLY	-	expression tag	UNP Q82IF8
A	-19	SER	-	expression tag	UNP Q82IF8
A	-18	SER	-	expression tag	UNP Q82IF8
A	-17	HIS	-	expression tag	UNP Q82IF8
A	-16	HIS	-	expression tag	UNP Q82IF8
A	-15	HIS	-	expression tag	UNP Q82IF8
A	-14	HIS	-	expression tag	UNP Q82IF8
A	-13	HIS	-	expression tag	UNP Q82IF8
A	-12	HIS	-	expression tag	UNP Q82IF8
A	-11	SER	-	expression tag	UNP Q82IF8
A	-10	SER	-	expression tag	UNP Q82IF8
A	-9	GLY	-	expression tag	UNP Q82IF8
A	-8	ARG	-	expression tag	UNP Q82IF8
A	-7	GLU	-	expression tag	UNP Q82IF8
A	-6	ASN	-	expression tag	UNP Q82IF8
A	-5	LEU	-	expression tag	UNP Q82IF8
A	-4	TYR	-	expression tag	UNP Q82IF8
A	-3	PHE	-	expression tag	UNP Q82IF8
A	-2	GLN	-	expression tag	UNP Q82IF8
A	-1	GLY	-	expression tag	UNP Q82IF8
A	0	HIS	-	expression tag	UNP Q82IF8
B	-21	MSE	-	expression tag	UNP Q82IF8
B	-20	GLY	-	expression tag	UNP Q82IF8
B	-19	SER	-	expression tag	UNP Q82IF8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-18	SER	-	expression tag	UNP Q82IF8
B	-17	HIS	-	expression tag	UNP Q82IF8
B	-16	HIS	-	expression tag	UNP Q82IF8
B	-15	HIS	-	expression tag	UNP Q82IF8
B	-14	HIS	-	expression tag	UNP Q82IF8
B	-13	HIS	-	expression tag	UNP Q82IF8
B	-12	HIS	-	expression tag	UNP Q82IF8
B	-11	SER	-	expression tag	UNP Q82IF8
B	-10	SER	-	expression tag	UNP Q82IF8
B	-9	GLY	-	expression tag	UNP Q82IF8
B	-8	ARG	-	expression tag	UNP Q82IF8
B	-7	GLU	-	expression tag	UNP Q82IF8
B	-6	ASN	-	expression tag	UNP Q82IF8
B	-5	LEU	-	expression tag	UNP Q82IF8
B	-4	TYR	-	expression tag	UNP Q82IF8
B	-3	PHE	-	expression tag	UNP Q82IF8
B	-2	GLN	-	expression tag	UNP Q82IF8
B	-1	GLY	-	expression tag	UNP Q82IF8
B	0	HIS	-	expression tag	UNP Q82IF8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	149	Total O 149 149	0	0
2	B	66	Total O 66 66	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative GntR-family transcriptional regulator



- Molecule 1: Putative GntR-family transcriptional regulator



4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	87.16 Å 87.16 Å 123.75 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.20 – 1.97 36.20 – 1.97	Depositor EDS
% Data completeness (in resolution range)	99.7 (36.20-1.97) 99.7 (36.20-1.97)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.23 (at 1.97 Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.176 , 0.218 0.180 , 0.205	Depositor DCC
R_{free} test set	1953 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	34.0	Xtriage
Anisotropy	0.217	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 38.6	EDS
Estimated twinning fraction	0.164 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 39005 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3903	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/1955	0.48	0/2638
1	B	0.28	0/1785	0.49	1/2417 (0.0%)
All	All	0.29	0/3740	0.48	1/5055 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	27	PRO	N-CA-CB	6.04	110.55	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1927	0	1910	29	0
1	B	1761	0	1725	31	0
2	A	149	0	0	3	0
2	B	66	0	0	1	0
All	All	3903	0	3635	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:210:VAL:HG22	1:A:213:HIS:CD2	2.13	0.84
1:A:49:LEU:HD12	2:A:282:HOH:O	1.83	0.78
1:B:100:ASP:O	1:B:103:VAL:HG12	1.83	0.78
1:B:44:SER:HB3	1:B:47:VAL:HG23	1.64	0.77
1:B:32:PRO:HB3	1:B:36:ARG:HD3	1.67	0.77
1:A:56:MSE:HG2	1:A:61:VAL:HG23	1.67	0.76
1:B:38:ARG:HH11	1:B:45:ASP:CG	1.96	0.69
1:A:52:ARG:HD3	1:A:56:MSE:HE2	1.78	0.64
1:B:31:LEU:HD12	1:B:32:PRO:HD2	1.86	0.57
1:A:171:VAL:O	1:A:171:VAL:HG22	2.04	0.57
1:B:16:LEU:O	1:B:20:ILE:HG13	2.04	0.57
1:A:38:ARG:HD2	1:A:45:ASP:HA	1.87	0.56
1:B:98:GLN:CG	1:B:103:VAL:HG11	2.37	0.55
1:B:171:VAL:HG11	1:B:179:ARG:HA	1.88	0.54
1:B:145:VAL:HG12	1:B:207:LEU:HD23	1.91	0.53
1:B:26:PRO:HG2	1:B:29:THR:OG1	2.08	0.53
1:A:140:ARG:HD2	2:A:323:HOH:O	2.10	0.51
1:A:64:ARG:HG3	1:A:70:TYR:CD1	2.46	0.51
1:A:109:SER:OG	1:A:167:GLU:HG3	2.11	0.51
1:A:14:GLY:HA2	1:A:17:ARG:NH1	2.26	0.51
1:B:237:PRO:HG3	1:B:240:ARG:NH2	2.27	0.50
1:B:239:ASP:OD1	1:B:240:ARG:HG3	2.13	0.49
1:A:26:PRO:O	1:A:29:THR:HB	2.14	0.48
1:B:157:VAL:HG22	1:B:227:ARG:NH1	2.29	0.47
1:B:157:VAL:CG2	1:B:227:ARG:HH12	2.29	0.46
1:B:157:VAL:HG22	1:B:227:ARG:HH12	1.80	0.46
1:A:38:ARG:NH1	1:A:45:ASP:OD1	2.49	0.46
1:B:139:PHE:HB2	1:B:147:MSE:HB3	1.97	0.46
1:B:50:GLU:OE2	1:B:53:LYS:HD2	2.17	0.45
1:B:74:ARG:HA	1:B:75:PRO:HD3	1.81	0.45
1:A:109:SER:HA	1:A:136:LYS:O	2.16	0.45
1:B:109:SER:OG	1:B:167:GLU:HG3	2.17	0.45
1:A:155:LEU:HD22	1:A:164:MSE:HE3	1.98	0.45
1:B:246:HIS:HD2	2:B:270:HOH:O	1.99	0.44
1:A:64:ARG:CG	1:A:70:TYR:CD1	3.00	0.44
1:B:243:VAL:HG12	1:B:245:TYR:CE2	2.53	0.44
1:A:201:ALA:O	1:A:205:LEU:HG	2.17	0.44
1:A:56:MSE:HG2	1:A:61:VAL:CG2	2.42	0.44
1:B:185:VAL:HG11	1:B:229:VAL:HG11	1.99	0.44
1:B:93:PRO:HD2	1:B:107:TRP:CZ2	2.54	0.43
1:A:74:ARG:HA	1:A:75:PRO:HD2	1.69	0.42
1:A:210:VAL:HG22	1:A:213:HIS:NE2	2.32	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:98:GLN:HG2	1:B:103:VAL:HG11	2.00	0.42
1:A:227[B]:ARG:HG3	2:A:365:HOH:O	2.18	0.42
1:B:92:THR:HB	1:B:107:TRP:CH2	2.55	0.42
1:B:38:ARG:NH1	1:B:45:ASP:CG	2.70	0.42
1:A:163:VAL:HG12	1:A:171:VAL:CG1	2.50	0.42
1:A:204:LEU:HD13	1:A:210:VAL:O	2.20	0.42
1:A:166:PRO:HA	1:A:179:ARG:NE	2.35	0.42
1:A:6:GLN:O	1:A:11[A]:ARG:NH1	2.54	0.41
1:A:86:ARG:NH1	1:A:178:GLU:OE2	2.53	0.41
1:A:109:SER:HB3	1:A:137:TYR:CD1	2.56	0.41
1:B:204:LEU:O	1:B:208:GLY:N	2.51	0.41
1:A:188:ASP:OD1	1:A:189:ASN:ND2	2.53	0.41
1:B:167:GLU:O	1:B:168:GLU:HG3	2.21	0.41
1:A:171:VAL:HG13	1:A:179:ARG:HG2	2.03	0.40
1:A:10:LEU:HA	1:A:10:LEU:HD23	1.96	0.40
1:B:98:GLN:HG3	1:B:103:VAL:HG11	2.03	0.40
1:B:25:LEU:HA	1:B:26:PRO:HD2	1.91	0.40
1:B:99:ALA:O	1:B:101:GLY:N	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/272 (89%)	239 (98%)	4 (2%)	0	100	100
1	B	228/272 (84%)	224 (98%)	3 (1%)	1 (0%)	39	27
All	All	471/544 (87%)	463 (98%)	7 (2%)	1 (0%)	52	43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	100	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/210 (94%)	195 (98%)	3 (2%)	72	68
1	B	177/210 (84%)	175 (99%)	2 (1%)	80	77
All	All	375/420 (89%)	370 (99%)	5 (1%)	74	72

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	ARG
1	A	74	ARG
1	A	95	ARG
1	B	36	ARG
1	B	49	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	238/272 (87%)	-0.03	2 (0%)	87 92	23, 37, 66, 92	0
1	B	229/272 (84%)	0.40	20 (8%)	13 20	29, 51, 104, 135	0
All	All	467/544 (85%)	0.18	22 (4%)	35 46	23, 42, 87, 135	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	72	ARG	6.2
1	B	10	LEU	5.8
1	B	61	VAL	3.9
1	B	29	THR	3.0
1	B	-3	PHE	2.9
1	B	16	LEU	2.8
1	B	55	LEU	2.6
1	B	21	VAL	2.6
1	B	70	TYR	2.6
1	B	12	VAL	2.5
1	A	87	PRO	2.5
1	B	31	LEU	2.4
1	B	11	ARG	2.4
1	B	71	VAL	2.4
1	B	18	LYS	2.3
1	B	20	ILE	2.3
1	B	62	GLU	2.3
1	A	104	ARG	2.3
1	B	59	GLY	2.2
1	B	9	TYR	2.1
1	B	-4	TYR	2.0
1	B	25	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.