



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:24 AM GMT

PDB ID : 3EJJ
Title : Structure of M-CSF bound to the first three domains of FMS
Authors : Chen, X.; Liu, H.; Focia, P.J.; Shim, A.; He, X.
Deposited on : 2008-09-18
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	155	
1	B	155	
2	X	289	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	X	306	X	-	-	-
3	NAG	X	307	X	-	-	-
3	NAG	X	6	X	-	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5213 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Colony stimulating factor-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	148	Total	C	N	O	S	0	0	0
			1204	752	203	239	10			
1	B	145	Total	C	N	O	S	0	0	0
			1184	740	200	234	10			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ALA	-	EXPRESSION TAG	UNP Q3U395
A	2	ASP	-	EXPRESSION TAG	UNP Q3U395
A	3	PRO	-	EXPRESSION TAG	UNP Q3U395
A	149	HIS	-	EXPRESSION TAG	UNP Q3U395
A	150	HIS	-	EXPRESSION TAG	UNP Q3U395
A	151	HIS	-	EXPRESSION TAG	UNP Q3U395
A	152	HIS	-	EXPRESSION TAG	UNP Q3U395
A	153	HIS	-	EXPRESSION TAG	UNP Q3U395
A	154	HIS	-	EXPRESSION TAG	UNP Q3U395
A	155	HIS	-	EXPRESSION TAG	UNP Q3U395
B	1	ALA	-	EXPRESSION TAG	UNP Q3U395
B	2	ASP	-	EXPRESSION TAG	UNP Q3U395
B	3	PRO	-	EXPRESSION TAG	UNP Q3U395
B	149	HIS	-	EXPRESSION TAG	UNP Q3U395
B	150	HIS	-	EXPRESSION TAG	UNP Q3U395
B	151	HIS	-	EXPRESSION TAG	UNP Q3U395
B	152	HIS	-	EXPRESSION TAG	UNP Q3U395
B	153	HIS	-	EXPRESSION TAG	UNP Q3U395
B	154	HIS	-	EXPRESSION TAG	UNP Q3U395
B	155	HIS	-	EXPRESSION TAG	UNP Q3U395

- Molecule 2 is a protein called Macrophage colony-stimulating factor 1 receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	X	272	Total 2115	C 1336	N 364	O 406	S 9	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	17	ALA	-	EXPRESSION TAG	UNP P09581
X	18	ASP	-	EXPRESSION TAG	UNP P09581
X	19	PRO	-	EXPRESSION TAG	UNP P09581
X	297	GLU	-	EXPRESSION TAG	UNP P09581
X	298	SER	-	EXPRESSION TAG	UNP P09581
X	299	HIS	-	EXPRESSION TAG	UNP P09581
X	300	HIS	-	EXPRESSION TAG	UNP P09581
X	301	HIS	-	EXPRESSION TAG	UNP P09581
X	302	HIS	-	EXPRESSION TAG	UNP P09581
X	303	HIS	-	EXPRESSION TAG	UNP P09581
X	304	HIS	-	EXPRESSION TAG	UNP P09581
X	305	HIS	-	EXPRESSION TAG	UNP P09581

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	X	2	Total 28	C 16	N 2	O 10	0	0
3	X	2	Total 28	C 16	N 2	O 10	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	187	Total 187	O 187	0	0
4	B	114	Total 114	O 114	0	0
4	X	353	Total 353	O 353	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Colony stimulating factor-1



• Molecule 1: Colony stimulating factor-1



• Molecule 2: Macrophage colony-stimulating factor 1 receptor





4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	158.85Å 158.85Å 237.95Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.40 24.19 – 2.40	Depositor EDS
% Data completeness (in resolution range)	5.0 (20.00-2.40) 99.0 (24.19-2.40)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 2.41Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.238 , 0.265 0.258 , 0.258	Depositor DCC
R_{free} test set	2249 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	55.9	Xtriage
Anisotropy	0.228	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 74.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 44808 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5213	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/1226	0.77	1/1653 (0.1%)
1	B	0.41	0/1205	0.62	0/1623
2	X	0.51	0/2159	0.85	6/2942 (0.2%)
All	All	0.52	0/4590	0.77	7/6218 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	X	3	0

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	90	CYS	CA-CB-SG	-8.58	98.55	114.00
2	X	214	ARG	N-CA-C	-8.06	89.23	111.00
2	X	218	GLU	N-CA-C	8.02	132.66	111.00
2	X	213	VAL	N-CA-C	-7.12	91.77	111.00
2	X	25	PRO	N-CA-C	-6.12	96.19	112.10

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	X	6	NAG	C1
3	X	306	NAG	C1
3	X	307	NAG	C1

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1204	0	1148	55	0
1	B	1184	0	1129	71	0
2	X	2115	0	2105	139	0
3	X	56	0	50	5	0
4	A	187	0	0	6	0
4	B	114	0	0	9	0
4	X	353	0	0	28	1
All	All	5213	0	4432	263	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.

The worst 5 of 263 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:X:138:VAL:HG12	2:X:162:PHE:CZ	1.63	1.32
2:X:138:VAL:CG1	2:X:162:PHE:CZ	2.32	1.13
2:X:54:ILE:HG12	2:X:58:TRP:HD1	1.11	1.09
2:X:138:VAL:HG12	2:X:162:PHE:HZ	0.90	1.06
2:X:138:VAL:CG1	2:X:162:PHE:CE1	2.46	0.99

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:X:315:HOH:O	4:X:315:HOH:O[4_555]	0.30	1.90

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	146/155 (94%)	136 (93%)	7 (5%)	3 (2%)	9	10
1	B	143/155 (92%)	122 (85%)	18 (13%)	3 (2%)	9	10
2	X	268/289 (93%)	239 (89%)	24 (9%)	5 (2%)	10	12
All	All	557/599 (93%)	497 (89%)	49 (9%)	11 (2%)	9	11

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	X	55	SER
2	X	102	LYS
1	A	24	ASP
1	A	32	GLN
1	B	112	GLN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	139/146 (95%)	116 (84%)	23 (16%)	3	3
1	B	137/146 (94%)	124 (90%)	13 (10%)	11	15
2	X	239/253 (94%)	220 (92%)	19 (8%)	15	23
All	All	515/545 (94%)	460 (89%)	55 (11%)	8	11

5 of 55 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	52	LYS
1	B	98	GLN
2	X	245	GLU
1	B	56	LEU
1	B	93	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	141	ASN
1	B	70	ASN
2	X	147	GLN
1	A	119	ASN
2	X	98	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	X	306	3,2	14,14,15	0.79	1 (7%)	15,19,21	1.03	0
3	NAG	X	307	3	14,14,15	0.72	0	15,19,21	1.46	3 (20%)
3	NAG	X	5	3,2	14,14,15	0.58	0	15,19,21	0.70	1 (6%)
3	NAG	X	6	3	14,14,15	0.58	0	15,19,21	1.04	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	X	306	3,2	1/1/5/7	0/6/23/26	0/1/1/1
3	NAG	X	307	3	1/1/5/7	0/6/23/26	0/1/1/1
3	NAG	X	5	3,2	-	0/6/23/26	0/1/1/1
3	NAG	X	6	3	1/1/5/7	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	X	306	NAG	C1-C2	2.20	1.55	1.52

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	X	307	NAG	C2-N2-C7	-2.53	119.78	123.04
3	X	6	NAG	C2-N2-C7	-2.17	120.25	123.04
3	X	5	NAG	C2-N2-C7	-2.08	120.37	123.04
3	X	6	NAG	C3-C4-C5	2.17	113.99	110.20
3	X	307	NAG	C4-C3-C2	2.59	115.25	111.23

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	X	307	NAG	C1
3	X	306	NAG	C1
3	X	6	NAG	C1

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	X	306	NAG	1	0
3	X	307	NAG	1	0
3	X	5	NAG	2	0
3	X	6	NAG	1	0

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	148/155 (95%)	-0.03	5 (3%) 49 49	24, 37, 81, 138	0
1	B	145/155 (93%)	1.39	35 (24%) 1 1	40, 98, 142, 150	0
2	X	272/289 (94%)	0.36	26 (9%) 10 10	28, 56, 125, 155	0
All	All	565/599 (94%)	0.52	66 (11%) 6 6	24, 56, 134, 155	0

The worst 5 of 66 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	148	SER	13.5
1	B	95	TYR	9.4
1	B	97	GLU	8.2
1	B	135	PHE	6.9
1	B	133	ASN	6.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	X	5	14/15	0.69	0.69	-	82,88,96,102	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAG	X	307	14/15	0.60	0.67	-	81,83,89,91	0
3	NAG	X	306	14/15	0.59	0.46	-	79,83,88,88	0
3	NAG	X	6	14/15	0.70	0.76	-	84,88,93,95	0

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.