



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 28, 2016 – 07:31 PM EDT

PDB ID : 5EJS
Title : Structure of Dictyostelium Discoideum Myosin VII MyTH4-FERM MF2 domain, mutant 1
Authors : Planelles-Herrero, V.J.; Sirkia, H.; Sourigues, Y.; Titus, M.A.; Houdusse, A.
Deposited on : 2015-11-02
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027790
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027790


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	496	<div> <div>2%</div> <div> <div></div> <div>93%</div> <div>6%</div> <div>.</div> </div> </div>
1	B	496	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div>7%</div> <div>.</div> <div>.</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8269 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myosin-I heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	496	Total	C	N	O	S	0	0	0
			4005	2554	659	770	22			
1	B	485	Total	C	N	O	S	0	0	0
			3922	2504	647	750	21			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	GLN	deletion	UNP Q9U1M8
A	?	-	GLN	deletion	UNP Q9U1M8
A	?	-	ALA	deletion	UNP Q9U1M8
A	?	-	THR	deletion	UNP Q9U1M8
A	?	-	LEU	deletion	UNP Q9U1M8
A	?	-	LYS	deletion	UNP Q9U1M8
A	?	-	ARG	deletion	UNP Q9U1M8
A	?	-	LYS	deletion	UNP Q9U1M8
A	?	-	ALA	deletion	UNP Q9U1M8
A	?	-	PRO	deletion	UNP Q9U1M8
A	?	-	VAL	deletion	UNP Q9U1M8
A	1909	GLU	LYS	engineered mutation	UNP Q9U1M8
A	1912	GLU	LYS	engineered mutation	UNP Q9U1M8
A	1913	GLU	LYS	engineered mutation	UNP Q9U1M8
A	2358	LEU	-	expression tag	UNP Q9U1M8
A	2359	GLU	-	expression tag	UNP Q9U1M8
A	2360	HIS	-	expression tag	UNP Q9U1M8
B	?	-	GLN	deletion	UNP Q9U1M8
B	?	-	GLN	deletion	UNP Q9U1M8
B	?	-	ALA	deletion	UNP Q9U1M8
B	?	-	THR	deletion	UNP Q9U1M8
B	?	-	LEU	deletion	UNP Q9U1M8
B	?	-	LYS	deletion	UNP Q9U1M8
B	?	-	ARG	deletion	UNP Q9U1M8
B	?	-	LYS	deletion	UNP Q9U1M8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	ALA	deletion	UNP Q9U1M8
B	?	-	PRO	deletion	UNP Q9U1M8
B	?	-	VAL	deletion	UNP Q9U1M8
B	1909	GLU	LYS	engineered mutation	UNP Q9U1M8
B	1912	GLU	LYS	engineered mutation	UNP Q9U1M8
B	1913	GLU	LYS	engineered mutation	UNP Q9U1M8
B	2359	LEU	-	expression tag	UNP Q9U1M8
B	2360	GLU	-	expression tag	UNP Q9U1M8
B	2361	HIS	-	expression tag	UNP Q9U1M8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	202	Total O 202 202	0	0
2	B	140	Total O 140 140	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Myosin-I heavy chain



- Molecule 1: Myosin-I heavy chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	40.17Å 158.84Å 194.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.91 – 2.70 38.91 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.2 (38.91-2.70) 99.7 (38.91-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 2.69Å)	Xtriage
Refinement program	BUSTER-TNT	Depositor
R, R_{free}	0.205 , 0.235 0.205 , 0.233	Depositor DCC
R_{free} test set	1756 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	59.6	Xtriage
Anisotropy	0.415	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 59.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8269	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/4087	0.67	0/5526
1	B	0.50	0/4002	0.68	0/5408
All	All	0.50	0/8089	0.67	0/10934

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4005	0	3967	12	0
1	B	3922	0	3892	20	0
2	A	202	0	0	0	0
2	B	140	0	0	1	0
All	All	8269	0	7859	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 32 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2331:ASN:O	1:B:2331:ASN:ND2	1.93	1.02

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2329:THR:OG1	1:B:2336:ILE:O	2.09	0.71
1:B:2329:THR:OG1	1:B:2336:ILE:HG23	1.93	0.69
1:A:2312:GLN:HE22	1:A:2358:LEU:HD12	1.67	0.59
1:A:2317:VAL:HG12	1:A:2326:ILE:HG22	1.85	0.57

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	492/496 (99%)	477 (97%)	14 (3%)	1 (0%)	52	80
1	B	479/496 (97%)	461 (96%)	17 (4%)	1 (0%)	52	80
All	All	971/992 (98%)	938 (97%)	31 (3%)	2 (0%)	52	80

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2207	PRO
1	B	2207	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	455/458 (99%)	437 (96%)	18 (4%)	38	69
1	B	445/458 (97%)	425 (96%)	20 (4%)	34	65
All	All	900/916 (98%)	862 (96%)	38 (4%)	36	68

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2358	LEU
1	B	1916	LYS
1	B	2329	THR
1	B	1871	SER
1	B	1938	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2208	ASN
1	A	2312	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	1875:GLN	C	1887:ASP	N	16.94

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	496/496 (100%)	-0.13	10 (2%) 68 69	36, 67, 116, 205	0
1	B	485/496 (97%)	0.14	26 (5%) 29 28	38, 82, 144, 187	0
All	All	981/992 (98%)	0.00	36 (3%) 45 45	36, 72, 137, 205	0

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2330	GLY	5.6
1	A	1888	PRO	5.4
1	B	1868	LYS	4.8
1	A	1874	ASN	4.6
1	B	1855	GLN	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.