



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:13 PM GMT

PDB ID : 1EK4  
Title : BETA-KETOACYL [ACYL CARRIER PROTEIN] SYNTHASE I IN COMPLEX WITH DODECANOIC ACID TO 1.85 RESOLUTION  
Authors : Olsen, J.G.; Kadziola, A.; Siggaard-Andersen, M.; von Wettstein-Knowles, P.; Larsen, S.  
Deposited on : 2000-03-06  
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	418	 87% 10% . .
1	B	418	 91% 5% . .
1	C	418	 91% 6% . .
1	D	418	 87% 9% . .

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12770 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA-KETOACYL [ACYL CARRIER PROTEIN] SYNTHASE I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	406	Total	C	N	O	S	0	0	0
			2983	1855	519	587	22			
1	B	406	Total	C	N	O	S	0	0	0
			2983	1855	519	587	22			
1	C	406	Total	C	N	O	S	0	0	0
			2983	1855	519	587	22			
1	D	406	Total	C	N	O	S	0	0	0
			2983	1855	519	587	22			

There are 56 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	EXPRESSION TAG	UNP P0A953
A	-10	ARG	-	EXPRESSION TAG	UNP P0A953
A	-9	GLY	-	EXPRESSION TAG	UNP P0A953
A	-8	SER	-	EXPRESSION TAG	UNP P0A953
A	-7	HIS	-	EXPRESSION TAG	UNP P0A953
A	-6	HIS	-	EXPRESSION TAG	UNP P0A953
A	-5	HIS	-	EXPRESSION TAG	UNP P0A953
A	-4	HIS	-	EXPRESSION TAG	UNP P0A953
A	-3	HIS	-	EXPRESSION TAG	UNP P0A953
A	-2	HIS	-	EXPRESSION TAG	UNP P0A953
A	-1	GLY	-	EXPRESSION TAG	UNP P0A953
A	0	SER	-	EXPRESSION TAG	UNP P0A953
A	4	VAL	ALA	CONFLICT	UNP P0A953
A	163	SER	CYS	ENGINEERED	UNP P0A953
B	-11	MET	-	EXPRESSION TAG	UNP P0A953
B	-10	ARG	-	EXPRESSION TAG	UNP P0A953
B	-9	GLY	-	EXPRESSION TAG	UNP P0A953
B	-8	SER	-	EXPRESSION TAG	UNP P0A953
B	-7	HIS	-	EXPRESSION TAG	UNP P0A953
B	-6	HIS	-	EXPRESSION TAG	UNP P0A953

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	HIS	-	EXPRESSION TAG	UNP P0A953
B	-4	HIS	-	EXPRESSION TAG	UNP P0A953
B	-3	HIS	-	EXPRESSION TAG	UNP P0A953
B	-2	HIS	-	EXPRESSION TAG	UNP P0A953
B	-1	GLY	-	EXPRESSION TAG	UNP P0A953
B	0	SER	-	EXPRESSION TAG	UNP P0A953
B	4	VAL	ALA	CONFLICT	UNP P0A953
B	163	SER	CYS	ENGINEERED	UNP P0A953
C	-11	MET	-	EXPRESSION TAG	UNP P0A953
C	-10	ARG	-	EXPRESSION TAG	UNP P0A953
C	-9	GLY	-	EXPRESSION TAG	UNP P0A953
C	-8	SER	-	EXPRESSION TAG	UNP P0A953
C	-7	HIS	-	EXPRESSION TAG	UNP P0A953
C	-6	HIS	-	EXPRESSION TAG	UNP P0A953
C	-5	HIS	-	EXPRESSION TAG	UNP P0A953
C	-4	HIS	-	EXPRESSION TAG	UNP P0A953
C	-3	HIS	-	EXPRESSION TAG	UNP P0A953
C	-2	HIS	-	EXPRESSION TAG	UNP P0A953
C	-1	GLY	-	EXPRESSION TAG	UNP P0A953
C	0	SER	-	EXPRESSION TAG	UNP P0A953
C	4	VAL	ALA	CONFLICT	UNP P0A953
C	163	SER	CYS	ENGINEERED	UNP P0A953
D	-11	MET	-	EXPRESSION TAG	UNP P0A953
D	-10	ARG	-	EXPRESSION TAG	UNP P0A953
D	-9	GLY	-	EXPRESSION TAG	UNP P0A953
D	-8	SER	-	EXPRESSION TAG	UNP P0A953
D	-7	HIS	-	EXPRESSION TAG	UNP P0A953
D	-6	HIS	-	EXPRESSION TAG	UNP P0A953
D	-5	HIS	-	EXPRESSION TAG	UNP P0A953
D	-4	HIS	-	EXPRESSION TAG	UNP P0A953
D	-3	HIS	-	EXPRESSION TAG	UNP P0A953
D	-2	HIS	-	EXPRESSION TAG	UNP P0A953
D	-1	GLY	-	EXPRESSION TAG	UNP P0A953
D	0	SER	-	EXPRESSION TAG	UNP P0A953
D	4	VAL	ALA	CONFLICT	UNP P0A953
D	163	SER	CYS	ENGINEERED	UNP P0A953

- Molecule 2 is LAURIC ACID (three-letter code: DAO) (formula: C<sub>12</sub>H<sub>24</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	12	1		
2	B	1	Total	C	O	0	0
			13	12	1		
2	C	1	Total	C	O	0	0
			13	12	1		
2	D	1	Total	C	O	0	0
			13	12	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	188	Total	O	0	0
			188	188		
3	B	204	Total	O	0	0
			204	204		
3	C	213	Total	O	0	0
			213	213		
3	D	181	Total	O	0	0
			181	181		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


#### • Molecule 1: BETA-KETOACYL [ACYL CARRIER PROTEIN] SYNTHASE I




#### • Molecule 1: BETA-KETOACYL [ACYL CARRIER PROTEIN] SYNTHASE I



#### • Molecule 1: BETA-KETOACYL [ACYL CARRIER PROTEIN] SYNTHASE I



#### • Molecule 1: BETA-KETOACYL [ACYL CARRIER PROTEIN] SYNTHASE I



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.64 Å   142.26 Å   213.74 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	28.92 – 1.85	Depositor
% Data completeness (in resolution range)	(Not available) (28.92-1.85)	Depositor
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS 0.9	Depositor
R, $R_{free}$	0.173 , 0.196	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	12770	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DAO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.32	0/3031	0.60	0/4093
1	B	0.31	0/3031	0.61	0/4093
1	C	0.32	0/3031	0.61	0/4093
1	D	0.32	0/3031	0.61	0/4093
All	All	0.32	0/12124	0.61	0/16372

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2983	0	2942	34	0
1	B	2983	0	2942	23	0
1	C	2983	0	2942	31	0
1	D	2983	0	2942	39	0
2	A	13	0	23	0	0
2	B	13	0	23	1	0
2	C	13	0	23	6	0
2	D	13	0	23	2	0
3	A	188	0	0	3	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	204	0	0	1	0
3	C	213	0	0	4	0
3	D	181	0	0	3	0
All	All	12770	0	11860	124	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 124 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:257:ILE:HG12	3:D:1068:HOH:O	1.59	1.01
1:B:405:LYS:H	1:B:405:LYS:HD2	1.29	0.96
1:C:405:LYS:H	1:C:405:LYS:HD2	1.31	0.91
1:A:3:ARG:HH22	1:A:405:LYS:HE2	1.36	0.91
1:D:51:ASN:HD22	1:D:51:ASN:H	1.19	0.89

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	404/418 (97%)	389 (96%)	14 (4%)	1 (0%)	52	36
1	B	404/418 (97%)	391 (97%)	11 (3%)	2 (0%)	34	17
1	C	404/418 (97%)	389 (96%)	13 (3%)	2 (0%)	34	17
1	D	404/418 (97%)	389 (96%)	13 (3%)	2 (0%)	34	17
All	All	1616/1672 (97%)	1558 (96%)	51 (3%)	7 (0%)	39	22

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	299	GLY
1	B	299	GLY
1	C	299	GLY
1	D	299	GLY
1	B	301	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/320 (97%)	305 (98%)	5 (2%)	70	57
1	B	310/320 (97%)	306 (99%)	4 (1%)	76	65
1	C	310/320 (97%)	307 (99%)	3 (1%)	82	76
1	D	310/320 (97%)	305 (98%)	5 (2%)	70	57
All	All	1240/1280 (97%)	1223 (99%)	17 (1%)	74	63

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	51	ASN
1	B	405	LYS
1	D	49	TRP
1	B	49	TRP
1	D	51	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	94	GLN
1	C	18	ASN
1	D	51	ASN
1	B	18	ASN
1	B	51	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	DAO	A	901	1	12,12,13	0.47	0	10,11,13	0.56	0
2	DAO	B	901	1	12,12,13	0.52	0	10,11,13	0.53	0
2	DAO	C	901	1	12,12,13	0.45	0	10,11,13	0.53	0
2	DAO	D	901	1	12,12,13	0.43	0	10,11,13	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DAO	A	901	1	-	0/9/10/11	0/0/0/0
2	DAO	B	901	1	-	0/9/10/11	0/0/0/0
2	DAO	C	901	1	-	0/9/10/11	0/0/0/0
2	DAO	D	901	1	-	0/9/10/11	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	901	DAO	1	0
2	C	901	DAO	6	0
2	D	901	DAO	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.