



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 29, 2016 – 05:02 PM EST

PDB ID : 5EKD  
Title : Human mitochondrial tryptophanyl-tRNA synthetase bound by indolmycin and Mn\*ATP.  
Authors : Williams, T.L.; Carter Jr., C.W.  
Deposited on : 2015-11-03  
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028320  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028320


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	5422 (1.84-1.80)
Clashscore	102246	6347 (1.84-1.80)
Ramachandran outliers	100387	6276 (1.84-1.80)
Sidechain outliers	100360	6276 (1.84-1.80)
RSRZ outliers	91569	5439 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 84%, grey 10%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>1%</span> <span>84%</span> <span>10%</span> <span>• 5%</span> </div> </div>
1	B	343	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 2%, yellow 15%, green 78%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>2%</span> <span>78%</span> <span>15%</span> <span>• 5%</span> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	5BX	A	401	-	-	-	X
5	GOL	B	405	-	-	-	X

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 11183 atoms, of which 5378 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tryptophan-tRNA ligase, mitochondrial.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	327	Total	C	H	N	O	S	0	0	0
			5225	1637	2660	442	477	9			
1	B	327	Total	C	H	N	O	S	0	4	0
			5261	1649	2678	443	482	9			

- Molecule 2 is (5S)-5-[(1R)-1-(1H-indol-3-yl)ethyl]-2-(methylamino)-1,3-oxazol-4(5H)-one (three-letter code: 5BX) (formula: C<sub>14</sub>H<sub>15</sub>N<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			19	14	3	2		
2	B	1	Total	C	N	O	0	0
			19	14	3	2		

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total 43	C 10	H 12	N 5	O 13	P 3	0	0
3	B	1	Total 43	C 10	H 12	N 5	O 13	P 3	0	0

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	2	Total Mn 2 2	0	0
4	A	2	Total Mn 2 2	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $\text{C}_3\text{H}_8\text{O}_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	H	O	0	0
			14	3	8	3		
5	B	1	Total	C	H	O	0	0
			14	3	8	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	283	Total	O	0	0
			283	283		
6	B	258	Total	O	0	0
			258	258		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tryptophan-tRNA ligase, mitochondrial



- Molecule 1: Tryptophan-tRNA ligase, mitochondrial



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.29Å 78.09Å 154.00Å 90.00° 96.55° 90.00°	Depositor
Resolution (Å)	29.06 – 1.82 29.06 – 1.82	Depositor EDS
% Data completeness (in resolution range)	98.4 (29.06-1.82) 98.4 (29.06-1.82)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.36 (at 1.82Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, $R_{free}$	0.170 , 0.207 0.171 , 0.193	Depositor DCC
$R_{free}$ test set	3069 reflections (5.07%)	DCC
Wilson B-factor (Å <sup>2</sup> )	21.6	Xtrriage
Anisotropy	0.134	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 54.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	11183	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MN, 5BX, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.57	11/2612 (0.4%)	0.81	2/3534 (0.1%)
1	B	1.63	10/2640 (0.4%)	0.85	4/3570 (0.1%)
All	All	1.60	21/5252 (0.4%)	0.83	6/7104 (0.1%)

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	89[A]	ARG	CA-C	7.23	1.71	1.52
1	B	89[B]	ARG	CA-C	7.23	1.71	1.52
1	A	312	GLU	CD-OE1	-6.39	1.18	1.25
1	A	128	SER	CB-OG	-5.98	1.34	1.42
1	B	200	TYR	CE1-CZ	-5.82	1.30	1.38
1	A	312	GLU	CD-OE2	-5.81	1.19	1.25
1	B	115	SER	CB-OG	-5.61	1.34	1.42
1	B	109	SER	CB-OG	-5.59	1.34	1.42
1	B	58	SER	CB-OG	-5.58	1.35	1.42
1	B	124	SER	CB-OG	-5.51	1.35	1.42
1	B	322	GLU	CD-OE2	-5.51	1.19	1.25
1	A	125	TRP	NE1-CE2	-5.50	1.30	1.37
1	A	264	TYR	CE1-CZ	-5.45	1.31	1.38
1	A	246	GLU	CD-OE1	-5.31	1.19	1.25
1	B	347	TYR	CE1-CZ	-5.29	1.31	1.38
1	A	320	GLU	CD-OE1	-5.27	1.19	1.25
1	A	109	SER	CB-OG	-5.20	1.35	1.42
1	B	39	SER	CB-OG	-5.14	1.35	1.42
1	A	127	LEU	C-O	-5.12	1.13	1.23
1	A	66	TYR	CE1-CZ	-5.02	1.32	1.38
1	A	246	GLU	CD-OE2	-5.01	1.20	1.25

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	89	ARG	NE-CZ-NH1	6.10	123.35	120.30
1	A	292	ARG	NE-CZ-NH2	-5.52	117.54	120.30
1	B	221	ARG	NE-CZ-NH1	5.41	123.00	120.30
1	B	206	VAL	C-N-CD	5.35	139.64	128.40
1	B	89[A]	ARG	CA-C-N	-5.05	106.09	117.20
1	B	89[B]	ARG	CA-C-N	-5.05	106.09	117.20

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2565	2660	2650	20	1
1	B	2583	2678	2677	49	1
2	A	19	0	15	0	0
2	B	19	0	15	0	0
3	A	31	12	12	0	0
3	B	31	12	12	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	B	12	16	16	2	0
6	A	283	0	0	2	1
6	B	258	0	0	2	1
All	All	5805	5378	5397	69	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance ( $\text{\AA}$ )	Clash overlap ( $\text{\AA}$ )
1:B:130:MET:CE	6:B:526:HOH:O	1.86	1.24
1:B:29:LEU:HD22	1:B:208[B]:GLU:CD	1.67	1.16
1:B:249:GLN:OE1	1:B:252:ARG:NH1	1.88	1.06

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:130:MET:HE2	6:B:526:HOH:O	1.55	0.93
1:B:61:ARG:NH1	1:B:65:GLU:OE2	2.14	0.81
1:B:61:ARG:NH1	1:B:65:GLU:CD	2.40	0.75
1:B:29:LEU:CD2	1:B:208[B]:GLU:CD	2.52	0.74
1:B:251:PHE:O	1:B:303:LYS:HE3	1.93	0.69
1:B:141:GLN:O	1:B:145:LYS:HD2	1.93	0.69
1:B:29:LEU:HD22	1:B:208[A]:GLU:HG3	1.75	0.68
1:B:234:LYS:HE3	1:B:250:LYS:HZ1	1.59	0.68
1:A:43:PRO:O	1:A:91:SER:HB3	1.95	0.67
1:A:234:LYS:HD2	1:A:250:LYS:HZ1	1.60	0.66
1:B:255:VAL:O	1:B:256:THR:HG22	1.97	0.64
1:B:29:LEU:CD2	1:B:208[B]:GLU:OE2	2.47	0.63
1:B:234:LYS:HE3	1:B:250:LYS:NZ	2.14	0.63
1:B:29:LEU:HD22	1:B:208[B]:GLU:OE2	1.98	0.62
1:B:255:VAL:O	1:B:256:THR:CG2	2.46	0.62
1:A:69:VAL:HG12	1:A:109:SER:HB3	1.84	0.60
1:B:328:LYS:O	1:B:332:GLU:HG3	2.02	0.58
1:B:61:ARG:NH1	1:B:65:GLU:OE1	2.36	0.58
1:B:255:VAL:C	1:B:256:THR:HG23	2.24	0.57
1:B:43:PRO:HD2	1:B:78[B]:SER:OG	2.03	0.57
1:A:234:LYS:HD2	1:A:250:LYS:NZ	2.19	0.57
1:B:255:VAL:C	1:B:256:THR:CG2	2.73	0.57
1:B:29:LEU:HD22	1:B:208[B]:GLU:CG	2.32	0.55
1:A:130:MET:HE3	6:A:556:HOH:O	2.07	0.54
1:B:292:ARG:HG3	1:B:292:ARG:HH11	1.73	0.54
1:B:347:TYR:CZ	5:B:406:GOL:H11	2.42	0.53
1:A:46:ILE:N	1:A:46:ILE:HD12	2.24	0.53
1:A:234:LYS:CD	1:A:250:LYS:HZ1	2.23	0.51
1:A:132:ARG:HG2	1:A:132:ARG:HH11	1.74	0.51
1:A:47:LEU:HD11	1:A:56:ILE:HG13	1.93	0.49
1:B:234:LYS:HE3	1:B:250:LYS:CE	2.43	0.49
1:A:37:VAL:HB	1:A:69:VAL:HG22	1.94	0.48
1:A:231:ASP:OD1	1:A:232:PRO:HD2	2.13	0.48
1:A:160:TYR:N	1:A:161:PRO:CD	2.76	0.47
1:A:113:GLN:HB2	1:A:335:LEU:HD13	1.96	0.47
1:B:29:LEU:HD22	1:B:208[B]:GLU:OE1	2.12	0.47
1:B:141:GLN:HB3	1:B:145:LYS:HE2	1.95	0.47
1:A:55:ALA:O	1:A:58:SER:HB2	2.14	0.46
1:A:234:LYS:HD2	1:A:250:LYS:CE	2.46	0.46
1:B:29:LEU:HB3	1:B:208[A]:GLU:OE2	2.17	0.45
1:B:61:ARG:HH11	1:B:65:GLU:CD	2.09	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:347:TYR:CE1	5:B:406:GOL:H11	2.52	0.45
1:A:46:ILE:H	1:A:46:ILE:HD12	1.81	0.45
1:B:29:LEU:HD21	1:B:208[B]:GLU:OE2	2.17	0.45
1:B:69:VAL:CG1	1:B:71:TYR:CZ	3.00	0.45
1:B:249:GLN:CD	1:B:252:ARG:HH11	2.20	0.44
1:B:61:ARG:HG2	1:B:61:ARG:NH1	2.31	0.44
1:B:315:ALA:N	1:B:316:PRO:CD	2.81	0.44
1:B:231:ASP:OD1	1:B:232:PRO:HD2	2.18	0.43
1:B:69:VAL:C	1:B:70:LEU:HD12	2.38	0.43
1:B:89[B]:ARG:HD2	1:B:89[B]:ARG:HA	1.87	0.43
1:B:289:GLU:O	1:B:293:ARG:HG3	2.19	0.43
1:B:160:TYR:N	1:B:161:PRO:CD	2.82	0.43
1:B:248:VAL:HG13	1:B:304:LEU:HD22	2.02	0.42
1:B:292:ARG:HG3	1:B:292:ARG:NH1	2.35	0.42
1:B:145:LYS:HD3	1:B:157:LEU:HD21	2.02	0.42
1:B:165:ALA:HB2	1:B:189:VAL:HG22	2.03	0.41
1:A:70:LEU:HG	1:A:110:ILE:HB	2.03	0.41
1:B:234:LYS:CE	1:B:250:LYS:HZ1	2.31	0.41
1:B:105:ASN:HA	1:B:106:PRO:HD2	1.91	0.41
1:B:133:LEU:HB3	1:B:134:PRO:HD3	2.03	0.41
1:B:37:VAL:HB	1:B:69:VAL:HG22	2.03	0.41
1:A:74:VAL:HB	1:A:77:HIS:HD2	1.84	0.40
1:A:116:GLN:HG3	6:A:595:HOH:O	2.21	0.40
1:B:145:LYS:HE3	1:B:160:TYR:HE2	1.87	0.40
1:A:69:VAL:HG12	1:A:109:SER:CB	2.49	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:ARG:HH11	1:B:33:SER:HG[3_455]	1.21	0.39
6:A:736:HOH:O	6:B:701:HOH:O[3_455]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/343 (95%)	317 (98%)	8 (2%)	0	100	100
1	B	327/343 (95%)	321 (98%)	6 (2%)	0	100	100
All	All	652/686 (95%)	638 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/296 (97%)	283 (99%)	3 (1%)	82	77
1	B	290/296 (98%)	278 (96%)	12 (4%)	37	18
All	All	576/592 (97%)	561 (97%)	15 (3%)	54	37

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	SER
1	A	160	TYR
1	A	323	LYS
1	B	29	LEU
1	B	30	GLN
1	B	31	LYS
1	B	34	LYS
1	B	47	LEU
1	B	58	SER
1	B	63	GLN
1	B	145	LYS
1	B	160	TYR
1	B	234	LYS
1	B	297	MET
1	B	323	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	122	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	5BX	A	401	-	20,21,21	2.73	9 (45%)	13,30,30	1.83	1 (7%)
3	ATP	A	402	4	26,33,33	0.92	1 (3%)	26,52,52	1.95	2 (7%)
2	5BX	B	401	-	20,21,21	2.77	9 (45%)	13,30,30	1.67	2 (15%)
3	ATP	B	402	4	26,33,33	0.91	1 (3%)	26,52,52	1.87	2 (7%)
5	GOL	B	405	-	5,5,5	0.45	0	5,5,5	0.14	0
5	GOL	B	406	-	5,5,5	0.33	0	5,5,5	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5BX	A	401	-	-	0/7/22/22	0/3/3/3
3	ATP	A	402	4	-	0/18/38/38	0/3/3/3
2	5BX	B	401	-	-	0/7/22/22	0/3/3/3
3	ATP	B	402	4	-	0/18/38/38	0/3/3/3
5	GOL	B	405	-	-	0/4/4/4	0/0/0/0
5	GOL	B	406	-	-	0/4/4/4	0/0/0/0

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	5BX	C07-C05	-3.53	1.48	1.52
2	B	401	5BX	C07-C05	-3.46	1.48	1.52
2	B	401	5BX	O08-C07	-2.48	1.41	1.45
2	A	401	5BX	O08-C07	-2.37	1.41	1.45
2	A	401	5BX	C17-C18	2.22	1.41	1.36
2	B	401	5BX	C17-C18	2.35	1.42	1.36
3	A	402	ATP	C5-C4	2.68	1.46	1.40
3	B	402	ATP	C5-C4	2.73	1.46	1.40
2	B	401	5BX	C01-N02	2.75	1.50	1.45
2	A	401	5BX	C01-N02	2.82	1.50	1.45
2	B	401	5BX	C16-C17	2.99	1.45	1.38
2	A	401	5BX	C16-C17	3.01	1.45	1.38
2	A	401	5BX	C11-C19	3.04	1.44	1.40
2	A	401	5BX	C03-N04	3.28	1.42	1.33
2	B	401	5BX	C03-N04	3.33	1.42	1.33
2	B	401	5BX	C11-C19	3.51	1.44	1.40
2	B	401	5BX	O08-C03	4.68	1.41	1.36
2	A	401	5BX	O08-C03	4.70	1.41	1.36
2	A	401	5BX	C03-N02	7.28	1.42	1.32
2	B	401	5BX	C03-N02	7.33	1.42	1.32

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	ATP	N3-C2-N1	-8.15	122.47	128.87
3	B	402	ATP	N3-C2-N1	-8.03	122.56	128.87
3	A	402	ATP	C1'-N9-C4	-2.30	124.24	126.81
3	B	402	ATP	C1'-N9-C4	-2.18	124.37	126.81
2	B	401	5BX	C10-C09-C11	2.31	114.65	110.40
2	B	401	5BX	C10-C09-C07	4.89	116.37	111.26
2	A	401	5BX	C10-C09-C07	5.80	117.32	111.26

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	406	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	327/343 (95%)	-0.15	4 (1%) 81 78	16, 25, 40, 51	0
1	B	327/343 (95%)	-0.10	8 (2%) 62 57	16, 26, 44, 53	0
All	All	654/686 (95%)	-0.13	12 (1%) 71 67	16, 26, 42, 53	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	29	LEU	4.5
1	B	147	THR	3.9
1	A	326	LEU	3.8
1	B	295	ALA	3.1
1	A	292	ARG	2.7
1	B	292	ARG	2.7
1	B	291	VAL	2.5
1	B	260	SER	2.5
1	A	66	TYR	2.3
1	B	261	GLU	2.2
1	A	163	LEU	2.1
1	B	146	THR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	B	405	6/6	0.76	0.49	11.50	25,32,39,46	14
2	5BX	A	401	19/19	0.96	0.20	2.62	15,17,21,21	0
2	5BX	B	401	19/19	0.97	0.19	1.48	13,17,19,19	0
3	ATP	B	402	31/31	0.96	0.11	0.11	15,20,27,34	0
3	ATP	A	402	31/31	0.97	0.11	0.07	13,19,25,40	0
4	MN	B	403	1/1	1.00	0.07	-0.74	18,18,18,18	0
4	MN	A	403	1/1	1.00	0.08	-	16,16,16,16	0
4	MN	B	404	1/1	0.98	0.06	-	30,30,30,30	1
4	MN	A	404	1/1	0.99	0.10	-	25,25,25,25	1
5	GOL	B	406	6/6	0.81	0.23	-	39,48,58,58	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.