



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 27, 2016 – 12:35 PM EDT

PDB ID : 5EOW  
Title : Crystal Structure of 6-Hydroxynicotinic Acid 3-Monooxygenase from *Pseudomonas putida* KT2440  
Authors : Yuen, M.E.; Zhen, W.; Gerwig, T.J.; Story, R.W.; Kopp, M.; Nakamoto, K.; Snider, M.J.; Hicks, K.A.  
Deposited on : 2015-11-10  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027790  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027790


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	405	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2828 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 6-hydroxynicotinate 3-monooxygenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	353	2723	1727	487	495	4	10	0	0	0

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	MSE	-	initiating methionine	UNP Q88FY2
A	-21	GLY	-	expression tag	UNP Q88FY2
A	-20	SER	-	expression tag	UNP Q88FY2
A	-19	ASP	-	expression tag	UNP Q88FY2
A	-18	LYS	-	expression tag	UNP Q88FY2
A	-17	ILE	-	expression tag	UNP Q88FY2
A	-16	HIS	-	expression tag	UNP Q88FY2
A	-15	HIS	-	expression tag	UNP Q88FY2
A	-14	HIS	-	expression tag	UNP Q88FY2
A	-13	HIS	-	expression tag	UNP Q88FY2
A	-12	HIS	-	expression tag	UNP Q88FY2
A	-11	HIS	-	expression tag	UNP Q88FY2
A	-10	SER	-	expression tag	UNP Q88FY2
A	-9	SER	-	expression tag	UNP Q88FY2
A	-8	GLY	-	expression tag	UNP Q88FY2
A	-7	GLU	-	expression tag	UNP Q88FY2
A	-6	ASN	-	expression tag	UNP Q88FY2
A	-5	LEU	-	expression tag	UNP Q88FY2
A	-4	TYR	-	expression tag	UNP Q88FY2
A	-3	PHE	-	expression tag	UNP Q88FY2
A	-2	GLN	-	expression tag	UNP Q88FY2
A	-1	GLY	-	expression tag	UNP Q88FY2
A	0	HIS	-	expression tag	UNP Q88FY2

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	52	Total	O	0	0
			52	52		



- Molecule 1: 6-hydroxynicotinate 3-monooxygenase

ALA ALA	A174	MSE
	P175	GLY
	I16	SER
	Y177	ASP
V181	HIS	
	A182	LYS
H183	ILE	
	H182	HIS
H188	HIS	
	HIS	HIS
GLY	HIS	
	VAL	HIS
ASN	ASN	
	ASN	SER
LEU	LEU	
	LEU	SER
ALA	ALA	
	ALA	GLY
Q194	GLU	
	ASN	ASN
Y215	LEU	
	LEU	LEU
V227	TYR	
	PHE	PHE
H232	GLN	
	GLY	GLN
ALA	HIS	
	HIS	HIS
TRP	NSE	
	ARG	ARG
ASP	GLY	
	GLY	GLY
PHE	ARG	
	ARG	ARG
GLN	Q5	
	K6	K6
GLY	I7	
	I7	I7
ALA	M53	
	K54	K54
PHE	I55	
	I55	I55
V242	R58	
	D243	M59
D244	M59	
	M59	M59
P280	E62	
	L281	E62
L289	Q63	
	Q63	Q63
C286	K64	
	L65	L65
H297	M68	
	M68	M68
P298	W75	
	W75	W75
E323	T105	
	T324	T105
G325	I133	
	L326	I133
L326	VAL	
	ASP	ASP
A332	GLU	
	GLU	GLU
Y372	GLY	
	D138	D138
L376	I165	
	I165	I165
GLU	A172	
	SER	A172
GLY	A172	
	A172	A172

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.69 Å 69.47 Å 115.43 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.39 – 2.10 44.39 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.5 (44.39-2.10) 97.6 (44.39-2.10)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.93 (at 2.10 Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.206 , 0.253 0.207 , 0.255	Depositor DCC
$R_{free}$ test set	1953 reflections (8.67%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.2	Xtriage
Anisotropy	0.457	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 44.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2828	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/2778	0.54	0/3750

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2723	0	2605	16	0
2	A	53	0	30	0	0
3	A	52	0	0	1	0
All	All	2828	0	2635	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:MSE:HE3	1:A:62:GLU:HG3	1.69	0.74

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:165:ILE:HG23	1:A:289:LEU:HD23	1.77	0.67
1:A:323:GLU:HG3	1:A:376:LEU:HD12	1.89	0.55
1:A:64:LYS:HG3	1:A:68:MSE:HE2	1.89	0.53
1:A:58:ARG:HG3	1:A:372:TYR:CZ	2.46	0.50
1:A:183:HIS:O	1:A:227:VAL:HA	2.13	0.49
1:A:181:VAL:HG23	3:A:514:HOH:O	2.13	0.49
1:A:138:ASP:OD1	1:A:138:ASP:N	2.45	0.48
1:A:282:PRO:HA	1:A:296:CYS:O	2.14	0.47
1:A:324:THR:OG1	1:A:332:ALA:HB2	2.15	0.47
1:A:75:TRP:C	1:A:75:TRP:CD1	2.89	0.46
1:A:105:THR:HB	1:A:215:TYR:CD2	2.51	0.46
1:A:55:ILE:HG22	1:A:59:MSE:HE3	1.99	0.44
1:A:65:LEU:HD23	1:A:68:MSE:CE	2.47	0.44
1:A:7:ILE:HD11	1:A:326:LEU:HD21	2.02	0.42
1:A:280:PRO:HD2	1:A:298:PRO:HG2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	343/405 (85%)	333 (97%)	9 (3%)	1 (0%)	46 45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	173	GLU



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	269/311 (86%)	267 (99%)	2 (1%)	88	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	75	TRP
1	A	244	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FAD	A	401	-	52,58,58	4.00	18 (34%)	52,89,89	2.72	13 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	401	-	-	0/30/50/50	0/6/6/6

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	FAD	C2B-C1B	-13.43	1.32	1.53
2	A	401	FAD	O4B-C4B	-5.70	1.32	1.45
2	A	401	FAD	C5A-C4A	-2.82	1.34	1.40
2	A	401	FAD	O3B-C3B	-2.77	1.36	1.43
2	A	401	FAD	O4-C4	-2.50	1.18	1.24
2	A	401	FAD	C4X-C10	2.53	1.45	1.40
2	A	401	FAD	O2B-C2B	2.62	1.49	1.43
2	A	401	FAD	C6A-N6A	2.90	1.46	1.34
2	A	401	FAD	C10-N10	4.34	1.44	1.39
2	A	401	FAD	C2-N1	4.41	1.47	1.38
2	A	401	FAD	C4-C4X	4.76	1.51	1.41
2	A	401	FAD	C2-N3	4.78	1.48	1.38
2	A	401	FAD	C10-N1	5.69	1.45	1.35
2	A	401	FAD	C4-N3	7.21	1.46	1.33
2	A	401	FAD	C9A-N10	7.35	1.49	1.38
2	A	401	FAD	C4X-N5	7.75	1.45	1.33
2	A	401	FAD	C5X-N5	9.36	1.50	1.35
2	A	401	FAD	O4B-C1B	13.21	1.60	1.41

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	FAD	N3A-C2A-N1A	-11.40	119.92	128.87
2	A	401	FAD	N6A-C6A-N1A	-6.00	108.45	118.52
2	A	401	FAD	C7M-C7-C6	-5.70	104.23	120.33
2	A	401	FAD	N3-C2-N1	-4.74	119.71	127.69

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	FAD	C8M-C8-C7	-3.73	112.71	120.73
2	A	401	FAD	C4X-C4-N3	-3.27	119.25	123.52
2	A	401	FAD	C4B-O4B-C1B	-3.13	106.33	109.64
2	A	401	FAD	C5B-C4B-C3B	-2.12	106.99	115.20
2	A	401	FAD	C4X-N5-C5X	2.05	119.14	116.72
2	A	401	FAD	C8M-C8-C9	2.38	127.04	120.33
2	A	401	FAD	C5X-C9A-N10	4.03	120.59	117.58
2	A	401	FAD	C4-N3-C2	6.14	120.28	115.16
2	A	401	FAD	C7M-C7-C8	6.48	134.68	120.73

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	343/405 (84%)	0.04	2 (0%) 90 92	8, 16, 29, 38	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	172	ALA	2.2
1	A	327	SER	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	FAD	A	401	53/53	0.96	0.12	0.33	7,12,19,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.