



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:25 PM GMT

PDB ID : 4EU8
Title : Succinyl-CoA:acetate CoA-transferase (AarCH6-S71A) in complex with CoA
Authors : Mullins, E.A.; Kappock, T.J.
Deposited on : 2012-04-25
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	514	 4% 90% 8% •
1	B	514	 4% 87% 11% •

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8513 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Succinyl-CoA:acetate coenzyme A transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	504	Total	C	N	O	S	0	5	0
			3867	2418	694	735	20			
1	B	504	Total	C	N	O	S	0	3	0
			3857	2410	694	733	20			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	71	ALA	SER	ENGINEERED MUTATION	UNP B3EY95
A	506	SER	-	EXPRESSION TAG	UNP B3EY95
A	507	LEU	-	EXPRESSION TAG	UNP B3EY95
A	508	GLU	-	EXPRESSION TAG	UNP B3EY95
A	509	HIS	-	EXPRESSION TAG	UNP B3EY95
A	510	HIS	-	EXPRESSION TAG	UNP B3EY95
A	511	HIS	-	EXPRESSION TAG	UNP B3EY95
A	512	HIS	-	EXPRESSION TAG	UNP B3EY95
A	513	HIS	-	EXPRESSION TAG	UNP B3EY95
A	514	HIS	-	EXPRESSION TAG	UNP B3EY95
B	71	ALA	SER	ENGINEERED MUTATION	UNP B3EY95
B	506	SER	-	EXPRESSION TAG	UNP B3EY95
B	507	LEU	-	EXPRESSION TAG	UNP B3EY95
B	508	GLU	-	EXPRESSION TAG	UNP B3EY95
B	509	HIS	-	EXPRESSION TAG	UNP B3EY95
B	510	HIS	-	EXPRESSION TAG	UNP B3EY95
B	511	HIS	-	EXPRESSION TAG	UNP B3EY95
B	512	HIS	-	EXPRESSION TAG	UNP B3EY95
B	513	HIS	-	EXPRESSION TAG	UNP B3EY95
B	514	HIS	-	EXPRESSION TAG	UNP B3EY95

- Molecule 2 is COENZYME A (three-letter code: COA) (formula: C₂₁H₃₆N₇O₁₆P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
2	B	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Cl	0	0
			2	2		
3	A	2	Total	Cl	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	356	Total	O	0	0
			356	356		
4	B	333	Total	O	0	0
			333	333		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Succinyl-CoA:acetate coenzyme A transferase



- Molecule 1: Succinyl-CoA:acetate coenzyme A transferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.28Å 109.67Å 120.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.69 – 1.81 34.69 – 1.81	Depositor EDS
% Data completeness (in resolution range)	99.8 (34.69-1.81) 97.4 (34.69-1.81)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.12 (at 1.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.180 , 0.226 0.178 , 0.224	Depositor DCC
R_{free} test set	4076 reflections (5.39%)	DCC
Wilson B-factor (Å ²)	16.5	Xtriage
Anisotropy	0.349	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 53.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 81595 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8513	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: COA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/3954	0.53	0/5351
1	B	0.35	0/3938	0.54	0/5329
All	All	0.35	0/7892	0.54	0/10680

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3867	0	3853	21	0
1	B	3857	0	3838	32	0
2	A	48	0	32	1	0
2	B	48	0	32	3	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	356	0	0	2	1
4	B	333	0	0	2	0
All	All	8513	0	7755	53	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:270:VAL:HG22	2:B:2598:COA:HN4	1.56	0.69
1:B:202:ASP:HB3	1:B:311:ILE:HG21	1.75	0.69
1:B:222:ARG:NH1	4:B:1127:HOH:O	2.27	0.68
1:B:7:ASN:ND2	1:B:141:GLU:O	2.27	0.66
1:A:58[A]:ARG:NH2	4:A:976:HOH:O	2.27	0.66
1:B:294:GLU:OE1	2:B:2598:COA:S1P	2.56	0.64
1:B:447:VAL:HG13	1:B:470:PHE:HZ	1.65	0.61
1:B:36:THR:HA	1:B:71:ALA:HB3	1.84	0.59
1:B:270:VAL:HG22	2:B:2598:COA:N4P	2.19	0.57
1:A:269:GLY:HA2	2:A:2598:COA:H22	1.87	0.55
1:B:300:MET:HA	1:B:303:MET:HE2	1.88	0.55
1:B:296:ILE:HG23	1:B:300:MET:HB3	1.89	0.53
1:B:247:LEU:HD13	1:B:286:PHE:HE2	1.73	0.53
1:A:282:LYS:HA	1:A:308:ARG:HD3	1.92	0.52
1:A:238:THR:HG23	1:A:435:GLU:HB3	1.93	0.50
1:B:317:PHE:CE1	1:B:338:ILE:HD13	2.46	0.50
1:A:36:THR:HG23	1:A:71:ALA:HB3	1.94	0.49
1:B:147:PRO:HG3	1:B:210:LEU:HG	1.97	0.47
1:B:317:PHE:HE1	1:B:338:ILE:HD13	1.79	0.47
1:A:372:VAL:HB	1:A:424:HIS:HB2	1.97	0.46
1:A:368:ILE:O	1:A:450:ALA:HA	2.15	0.46
1:A:13:LYS:NZ	1:A:214:PRO:O	2.49	0.46
1:B:237:GLU:H	1:B:237:GLU:CD	2.20	0.46
1:A:230:ALA:HA	1:A:231:PRO:HD3	1.81	0.45
1:B:372:VAL:HB	1:B:424:HIS:HB2	1.98	0.45
1:B:368:ILE:O	1:B:450:ALA:HA	2.17	0.44
1:A:447:VAL:HG13	1:A:470:PHE:HZ	1.82	0.44
1:B:135:GLU:HG2	1:B:148:THR:HG21	1.99	0.44
1:A:12:SER:HB3	4:A:927:HOH:O	2.17	0.44
1:B:7:ASN:HB2	1:B:140:THR:O	2.18	0.44
1:B:114:LEU:HG	1:B:387:GLY:C	2.38	0.44
1:B:143:GLY:HA3	4:B:1081:HOH:O	2.17	0.44
1:B:331:MET:O	1:B:335:ARG:HB2	2.19	0.43
1:A:147:PRO:HG2	1:A:180:HIS:CD2	2.54	0.43
1:A:243:ALA:O	1:A:247:LEU:HD23	2.17	0.43
1:B:46:LYS:HG2	1:B:79:GLU:HG3	1.99	0.43
1:A:246:LEU:HD22	1:A:402:LEU:HD21	2.00	0.43
1:B:201:ALA:HB3	1:B:290:VAL:HG11	2.00	0.43
1:A:91:PRO:HG3	1:A:117:VAL:HG21	2.01	0.42
1:A:286:PHE:O	1:A:308:ARG:NH2	2.51	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:49:ALA:O	1:B:53:GLU:HG3	2.19	0.42
1:B:41:PRO:HG3	1:B:133:LEU:HG	2.00	0.42
1:B:365:GLU:HG2	1:B:373:ASN:HB3	2.01	0.42
1:B:148:THR:OG1	1:B:149:SER:N	2.53	0.42
1:B:319:LEU:HB3	1:B:323:ALA:HB3	2.01	0.42
1:A:411:LYS:HD3	1:A:411:LYS:HA	1.89	0.41
1:B:300:MET:HA	1:B:303:MET:CE	2.51	0.41
1:A:235:PRO:HB3	1:A:276:ALA:HB2	2.03	0.40
1:B:327:ILE:HG13	1:B:334:PHE:CD2	2.56	0.40
1:A:365:GLU:HG2	1:A:373:ASN:HB3	2.03	0.40
1:A:371:ASN:OD1	1:A:426:MET:HG3	2.22	0.40
1:A:332:ASP:N	1:A:332:ASP:OD1	2.54	0.40
1:B:9:ALA:O	1:B:12:SER:OG	2.33	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1158:HOH:O	4:A:1162:HOH:O[1_655]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	507/514 (99%)	487 (96%)	18 (4%)	2 (0%)	39	23
1	B	505/514 (98%)	489 (97%)	15 (3%)	1 (0%)	52	35
All	All	1012/1028 (98%)	976 (96%)	33 (3%)	3 (0%)	46	29

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	504	PRO
1	A	36	THR
1	B	70	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	413/418 (99%)	408 (99%)	5 (1%)	78	71
1	B	411/418 (98%)	407 (99%)	4 (1%)	82	77
All	All	824/836 (99%)	815 (99%)	9 (1%)	80	74

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	LYS
1	A	62	TYR
1	A	257	ASN
1	A	279	GLU
1	A	308	ARG
1	B	62	TYR
1	B	301	LEU
1	B	308	ARG
1	B	332	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	COA	A	2598	-	40,50,50	1.72	5 (12%)	50,75,75	1.70	6 (12%)
2	COA	B	2598	-	40,50,50	1.65	4 (10%)	50,75,75	2.09	9 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	A	2598	-	-	0/44/64/64	0/3/3/3
2	COA	B	2598	-	-	0/44/64/64	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2598	COA	C2B-C3B	-4.92	1.41	1.53
2	A	2598	COA	C2B-C3B	-4.85	1.42	1.53
2	A	2598	COA	C3B-C4B	-2.15	1.46	1.52
2	B	2598	COA	C6A-N6A	3.01	1.44	1.34
2	A	2598	COA	C6A-N6A	3.20	1.44	1.34
2	B	2598	COA	C5P-N4P	3.95	1.42	1.33
2	A	2598	COA	C5P-N4P	4.91	1.45	1.33
2	B	2598	COA	C9P-N8P	5.53	1.45	1.33
2	A	2598	COA	C9P-N8P	5.85	1.45	1.33

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	2598	COA	N3A-C2A-N1A	-9.25	121.81	128.89
2	A	2598	COA	N3A-C2A-N1A	-9.14	121.89	128.89
2	B	2598	COA	C7P-C6P-C5P	-4.72	104.53	112.31
2	B	2598	COA	C7P-N8P-C9P	-4.53	113.56	122.53
2	B	2598	COA	C2B-C1B-N9A	-4.03	108.13	114.29
2	B	2598	COA	C3P-N4P-C5P	-3.00	116.90	122.79
2	B	2598	COA	C1B-N9A-C4A	-2.33	123.42	126.94
2	A	2598	COA	C7P-N8P-C9P	-2.31	117.95	122.53
2	A	2598	COA	C1B-N9A-C4A	-2.26	123.53	126.94
2	B	2598	COA	C4A-C5A-N7A	-2.08	107.56	109.48
2	A	2598	COA	O3A-P1A-O5B	2.00	108.25	102.94
2	A	2598	COA	O5B-C5B-C4B	2.11	116.91	109.12
2	A	2598	COA	CDP-CBP-CCP	2.14	111.28	108.50
2	B	2598	COA	O6A-CCP-CBP	2.23	114.13	110.55
2	B	2598	COA	CDP-CBP-CCP	2.37	111.58	108.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2598	COA	1	0
2	B	2598	COA	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	504/514 (98%)	0.06	20 (3%) 42 36	7, 16, 38, 61	0
1	B	504/514 (98%)	0.21	21 (4%) 40 34	8, 20, 38, 52	0
All	All	1008/1028 (98%)	0.14	41 (4%) 41 35	7, 18, 38, 61	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	505	SER	7.3
1	A	2	THR	6.9
1	A	232	PHE	6.5
1	B	2	THR	5.4
1	B	8	VAL	4.4
1	A	329	ASN	4.1
1	B	333	PHE	3.9
1	A	272	ASN	3.8
1	B	310	ARG	3.7
1	B	505	SER	3.5
1	A	269	GLY	3.5
1	B	332	ASP	3.4
1	A	283	GLU	3.4
1	A	332	ASP	3.2
1	B	329	ASN	3.1
1	A	231	PRO	3.1
1	B	325	GLU	3.0
1	A	270	VAL	3.0
1	A	328	ASN	3.0
1	A	333	PHE	2.8
1	B	142	ASP	2.8
1	A	325	GLU	2.7
1	A	323	ALA	2.7
1	B	54	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	50	GLN	2.6
1	B	175	MET	2.5
1	B	327	ILE	2.5
1	B	307	GLY	2.5
1	A	285	PRO	2.5
1	B	58	ARG	2.4
1	A	326	GLU	2.3
1	A	230	ALA	2.3
1	B	328	ASN	2.3
1	B	57	ASP	2.3
1	B	174	PRO	2.3
1	B	15	CYS	2.3
1	B	230	ALA	2.3
1	A	305	ASP	2.2
1	A	271	GLY	2.1
1	A	279	GLU	2.1
1	B	171	TRP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	COA	B	2598	48/48	0.93	0.14	1.23	13,25,49,69	0
2	COA	A	2598	48/48	0.93	0.12	-0.33	8,24,47,60	0
3	CL	A	515	1/1	0.99	0.07	-0.45	15,15,15,15	0
3	CL	B	516	1/1	0.97	0.07	-2.23	30,30,30,30	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CL	B	515	1/1	0.99	0.05	-2.37	22,22,22,22	0
3	CL	A	516	1/1	0.97	0.04	-	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.