



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 6, 2016 – 02:44 AM EST

PDB ID : 5F0P
Title : Structure of retromer VPS26-VPS35 subunits bound to SNX3 and DMT1(L557M) (SeMet labeled)
Authors : Lucas, M.; Gershlick, D.; Vidaurrezaga, A.; Rojas, A.L.; Bonifacino, J.S.; Hierro, A.
Deposited on : 2015-11-27
Resolution : 2.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028442
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3004 (2.80-2.76)
Clashscore	102246	3480 (2.80-2.76)
Ramachandran outliers	100387	3423 (2.80-2.76)
Sidechain outliers	100360	3425 (2.80-2.76)
RSRZ outliers	91569	3016 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	462	<div> <div>3%</div> <div>97%</div> <div>..</div> </div>
2	B	321	<div> <div>2%</div> <div>87%</div> <div>9%</div> </div>
3	C	167	<div> <div>18%</div> <div>86%</div> <div>12%</div> </div>
4	D	18	<div> <div>72%</div> <div>11%</div> <div>17%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	SO4	C	202	-	-	-	X
7	EDO	A	509	-	-	-	X
7	EDO	B	406	-	-	-	X
7	EDO	B	407	-	-	-	X
7	EDO	B	409	-	-	-	X

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 7630 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vacuolar protein sorting-associated protein 35.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	458	3709	2359	631	697	22	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	GLY	-	expression tag	UNP Q96QK1
A	10	ALA	-	expression tag	UNP Q96QK1
A	11	MET	-	expression tag	UNP Q96QK1
A	12	GLY	-	expression tag	UNP Q96QK1
A	13	SER	-	expression tag	UNP Q96QK1

- Molecule 2 is a protein called Vacuolar protein sorting-associated protein 26A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
2	B	293	2410	1554	400	447	2	7	0	0	0

- Molecule 3 is a protein called Sorting nexin-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	147	1204	764	217	221	2	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-4	GLY	-	expression tag	UNP O60493
C	-3	ALA	-	expression tag	UNP O60493
C	-2	MET	-	expression tag	UNP O60493
C	-1	GLY	-	expression tag	UNP O60493
C	0	SER	-	expression tag	UNP O60493


- Molecule 4 is a protein called Natural resistance-associated macrophage protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	15	Total	C	N	O	Se	0	0	0
			130	81	25	22	2			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	557	MSE	LEU	engineered mutation	UNP P49281
D	561	SER	-	expression tag	UNP P49281
D	562	HIS	-	expression tag	UNP P49281
D	563	HIS	-	expression tag	UNP P49281
D	564	HIS	-	expression tag	UNP P49281
D	565	HIS	-	expression tag	UNP P49281
D	566	HIS	-	expression tag	UNP P49281

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		
7	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	Zn	0	0
			1	1		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	12	Total	O	0	0
			12	12		
9	B	22	Total	O	0	0
			22	22		
9	C	8	Total	O	0	0
			8	8		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Vacuolar protein sorting-associated protein 35

Chain A: 





- Molecule 2: Vacuolar protein sorting-associated protein 26A

Chain B: 



- Molecule 3: Sorting nexin-3

Chain C: 



- Molecule 4: Natural resistance-associated macrophage protein 2

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	371.29Å 75.32Å 57.27Å 90.00° 97.66° 90.00°	Depositor
Resolution (Å)	56.76 – 2.78 56.76 – 2.78	Depositor EDS
% Data completeness (in resolution range)	95.1 (56.76-2.78) 95.2 (56.76-2.78)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 2.77Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.219 , 0.261 0.225 , 0.260	Depositor DCC
R_{free} test set	1900 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	55.2	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7630	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/3770	0.57	1/5099 (0.0%)
2	B	0.37	0/2452	0.63	0/3293
3	C	0.43	0/1230	0.62	0/1662
4	D	0.42	0/133	0.65	0/176
All	All	0.39	0/7585	0.60	1/10230 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	ARG	NE-CZ-NH2	5.19	122.89	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3709	0	3762	5	0
2	B	2410	0	2436	8	0
3	C	1204	0	1203	2	0
4	D	130	0	115	0	0
5	A	10	0	0	0	0
5	B	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	15	0	0	0	0
6	A	12	0	16	0	0
6	B	12	0	16	0	0
7	A	32	0	48	0	0
7	B	44	0	66	0	0
7	C	4	0	6	0	0
8	B	1	0	0	0	0
9	A	12	0	0	0	0
9	B	22	0	0	1	0
9	C	8	0	0	0	0
All	All	7630	0	7668	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:174:LEU:HD21	2:B:278:LEU:HD22	1.75	0.68
1:A:355:ILE:HG13	1:A:372:VAL:HG11	1.86	0.57
3:C:69:ARG:NH2	3:C:147:ASP:O	2.37	0.55
1:A:123:PRO:HB2	1:A:172:THR:HG22	1.95	0.48
1:A:411:ASN:O	1:A:414:THR:HG22	2.12	0.48
2:B:16:ILE:HD12	2:B:147:TYR:HB2	1.96	0.48
1:A:257:LEU:O	1:A:257:LEU:HD12	2.14	0.48
2:B:121:TYR:CD1	2:B:261:MSE:HE1	2.50	0.46
2:B:212:ILE:HD12	2:B:275:ASN:HB2	1.99	0.45
2:B:285:ARG:NH1	9:B:501:HOH:O	2.47	0.44
2:B:16:ILE:HD11	2:B:134:VAL:HG23	2.00	0.43
2:B:219:ILE:O	2:B:219:ILE:HD12	2.19	0.42
1:A:253:CYS:HA	3:C:22:TYR:CG	2.54	0.42
2:B:261:MSE:HE3	2:B:270:VAL:HB	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	456/462 (99%)	450 (99%)	6 (1%)	0	100	100
2	B	291/321 (91%)	280 (96%)	11 (4%)	0	100	100
3	C	145/167 (87%)	139 (96%)	6 (4%)	0	100	100
4	D	13/18 (72%)	13 (100%)	0	0	100	100
All	All	905/968 (94%)	882 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	422/424 (100%)	419 (99%)	3 (1%)	88	97
2	B	267/284 (94%)	266 (100%)	1 (0%)	93	98
3	C	133/148 (90%)	133 (100%)	0	100	100
4	D	15/15 (100%)	13 (87%)	2 (13%)	5	13
All	All	837/871 (96%)	831 (99%)	6 (1%)	88	97

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65	GLU
1	A	81	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	257	LEU
2	B	33	ASP
4	D	560	MSE
4	D	563	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	456	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 31 ligands modelled in this entry, 1 is monoatomic - leaving 30 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	A	501	-	4,4,4	0.39	0	6,6,6	0.19	0
5	SO4	A	502	-	4,4,4	0.39	0	6,6,6	0.15	0
6	GOL	A	503	-	5,5,5	0.30	0	5,5,5	0.31	0
6	GOL	A	504	-	5,5,5	0.32	0	5,5,5	0.58	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	EDO	A	505	-	3,3,3	0.55	0	2,2,2	0.12	0
7	EDO	A	506	-	3,3,3	0.38	0	2,2,2	0.41	0
7	EDO	A	507	-	3,3,3	0.48	0	2,2,2	0.28	0
7	EDO	A	508	-	3,3,3	0.51	0	2,2,2	0.23	0
7	EDO	A	509	-	3,3,3	0.44	0	2,2,2	0.40	0
7	EDO	A	510	-	3,3,3	0.44	0	2,2,2	0.43	0
7	EDO	A	511	-	3,3,3	0.49	0	2,2,2	0.21	0
7	EDO	A	512	-	3,3,3	0.47	0	2,2,2	0.31	0
5	SO4	B	401	-	4,4,4	0.33	0	6,6,6	0.20	0
6	GOL	B	402	-	5,5,5	0.30	0	5,5,5	0.32	0
7	EDO	B	403	-	3,3,3	0.48	0	2,2,2	0.39	0
7	EDO	B	404	-	3,3,3	0.42	0	2,2,2	0.40	0
7	EDO	B	405	-	3,3,3	0.49	0	2,2,2	0.23	0
7	EDO	B	406	-	3,3,3	0.48	0	2,2,2	0.18	0
7	EDO	B	407	-	3,3,3	0.46	0	2,2,2	0.41	0
7	EDO	B	408	-	3,3,3	0.43	0	2,2,2	0.47	0
7	EDO	B	409	-	3,3,3	0.43	0	2,2,2	0.45	0
7	EDO	B	410	-	3,3,3	0.44	0	2,2,2	0.31	0
7	EDO	B	411	-	3,3,3	0.47	0	2,2,2	0.32	0
7	EDO	B	412	-	3,3,3	0.53	0	2,2,2	0.16	0
7	EDO	B	413	-	3,3,3	0.51	0	2,2,2	0.21	0
6	GOL	B	414	-	5,5,5	0.18	0	5,5,5	0.12	0
5	SO4	C	201	-	4,4,4	0.37	0	6,6,6	0.17	0
5	SO4	C	202	-	4,4,4	0.32	0	6,6,6	0.17	0
5	SO4	C	203	-	4,4,4	0.33	0	6,6,6	0.10	0
7	EDO	C	204	-	3,3,3	0.48	0	2,2,2	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SO4	A	501	-	-	0/0/0/0	0/0/0/0
5	SO4	A	502	-	-	0/0/0/0	0/0/0/0
6	GOL	A	503	-	-	0/4/4/4	0/0/0/0
6	GOL	A	504	-	-	0/4/4/4	0/0/0/0
7	EDO	A	505	-	-	0/1/1/1	0/0/0/0
7	EDO	A	506	-	-	0/1/1/1	0/0/0/0
7	EDO	A	507	-	-	0/1/1/1	0/0/0/0
7	EDO	A	508	-	-	0/1/1/1	0/0/0/0
7	EDO	A	509	-	-	0/1/1/1	0/0/0/0
7	EDO	A	510	-	-	0/1/1/1	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	A	511	-	-	0/1/1/1	0/0/0/0
7	EDO	A	512	-	-	0/1/1/1	0/0/0/0
5	SO4	B	401	-	-	0/0/0/0	0/0/0/0
6	GOL	B	402	-	-	0/4/4/4	0/0/0/0
7	EDO	B	403	-	-	0/1/1/1	0/0/0/0
7	EDO	B	404	-	-	0/1/1/1	0/0/0/0
7	EDO	B	405	-	-	0/1/1/1	0/0/0/0
7	EDO	B	406	-	-	0/1/1/1	0/0/0/0
7	EDO	B	407	-	-	0/1/1/1	0/0/0/0
7	EDO	B	408	-	-	0/1/1/1	0/0/0/0
7	EDO	B	409	-	-	0/1/1/1	0/0/0/0
7	EDO	B	410	-	-	0/1/1/1	0/0/0/0
7	EDO	B	411	-	-	0/1/1/1	0/0/0/0
7	EDO	B	412	-	-	0/1/1/1	0/0/0/0
7	EDO	B	413	-	-	0/1/1/1	0/0/0/0
6	GOL	B	414	-	-	0/4/4/4	0/0/0/0
5	SO4	C	201	-	-	0/0/0/0	0/0/0/0
5	SO4	C	202	-	-	0/0/0/0	0/0/0/0
5	SO4	C	203	-	-	0/0/0/0	0/0/0/0
7	EDO	C	204	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	458/462 (99%)	0.30	14 (3%) 52 45	36, 71, 123, 169	0
2	B	286/321 (89%)	0.07	7 (2%) 62 55	30, 47, 99, 132	0
3	C	147/167 (88%)	1.04	30 (20%) 1 1	35, 104, 133, 150	0
4	D	13/18 (72%)	0.80	0 100 100	56, 83, 114, 117	0
All	All	904/968 (93%)	0.35	51 (5%) 28 20	30, 64, 124, 169	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	451	ILE	6.1
3	C	51	ILE	5.7
3	C	29	LEU	5.0
3	C	146	GLN	5.0
3	C	39	VAL	5.0
3	C	88	VAL	4.7
1	A	387	HIS	4.6
2	B	33	ASP	4.5
3	C	145	LEU	4.3
3	C	140	CYS	4.3
3	C	141	LEU	4.0
2	B	219	ILE	3.7
1	A	450	GLU	3.6
3	C	74	PHE	3.6
1	A	386	GLU	3.5
3	C	53	VAL	3.5
3	C	49	TYR	3.4
2	B	34	GLY	3.4
1	A	456	GLN	3.3
2	B	221	PRO	3.3
1	A	424	LEU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	67	VAL	3.0
3	C	43	ARG	3.0
3	C	90	PRO	3.0
1	A	385	LEU	3.0
3	C	81	LEU	3.0
1	A	168	THR	2.9
3	C	85	SER	2.8
3	C	54	LYS	2.8
3	C	33	VAL	2.7
1	A	448	ASN	2.7
3	C	83	ARG	2.7
2	B	8	PHE	2.7
1	A	171	GLU	2.7
3	C	134	LEU	2.5
3	C	64	GLU	2.5
3	C	129	VAL	2.5
3	C	144	PHE	2.5
1	A	461	MET	2.5
1	A	364	ASP	2.3
3	C	137	ASN	2.3
3	C	44	GLY	2.3
3	C	82	GLU	2.3
2	B	172	ASP	2.3
1	A	390	THR	2.3
3	C	65	SER	2.2
3	C	68	ARG	2.2
1	A	464	VAL	2.1
3	C	135	ALA	2.1
3	C	149	ILE	2.0
2	B	218	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
7	EDO	B	409	4/4	0.82	0.47	6.34	71,72,72,73	0
7	EDO	B	406	4/4	0.77	0.34	5.12	80,81,83,83	0
7	EDO	A	509	4/4	0.91	0.39	4.10	63,63,64,66	0
7	EDO	B	407	4/4	0.94	0.28	2.21	57,58,58,58	0
5	SO4	C	202	5/5	0.60	0.34	2.04	141,144,146,147	0
7	EDO	B	412	4/4	0.84	0.21	1.97	75,79,81,81	0
7	EDO	B	408	4/4	0.92	0.36	1.63	70,70,72,73	0
7	EDO	A	506	4/4	0.94	0.24	1.54	52,52,52,54	0
7	EDO	A	505	4/4	0.90	0.19	1.41	49,50,53,55	0
5	SO4	C	203	5/5	0.83	0.32	1.14	124,127,127,130	0
7	EDO	B	405	4/4	0.94	0.20	0.97	54,55,56,56	0
6	GOL	A	504	6/6	0.89	0.19	-0.08	66,67,68,68	0
7	EDO	B	403	4/4	0.91	0.19	-0.21	49,51,51,51	0
5	SO4	B	401	5/5	0.96	0.15	-0.94	72,73,74,76	0
5	SO4	C	201	5/5	0.91	0.13	-1.81	94,95,97,100	0
7	EDO	A	508	4/4	0.79	0.20	-	75,76,77,78	0
7	EDO	B	411	4/4	0.79	0.20	-	74,76,78,79	0
7	EDO	A	511	4/4	0.83	0.37	-	82,85,86,87	0
6	GOL	B	414	6/6	0.85	0.28	-	57,62,66,66	0
5	SO4	A	502	5/5	0.86	0.16	-	115,118,120,120	0
7	EDO	A	507	4/4	0.86	0.12	-	79,79,79,80	0
6	GOL	A	503	6/6	0.86	0.17	-	90,94,94,95	0
7	EDO	A	510	4/4	0.81	0.17	-	85,86,86,86	0
5	SO4	A	501	5/5	0.88	0.17	-	106,106,108,108	0
7	EDO	B	413	4/4	0.61	0.34	-	79,80,81,81	0
8	ZN	B	415	1/1	0.99	0.15	-	64,64,64,64	0
7	EDO	B	404	4/4	0.93	0.23	-	68,70,72,73	0
7	EDO	A	512	4/4	0.80	0.27	-	99,100,100,101	0
7	EDO	B	410	4/4	0.88	0.21	-	71,72,72,73	0
7	EDO	C	204	4/4	0.78	0.36	-	82,84,84,85	0
6	GOL	B	402	6/6	0.89	0.13	-	51,52,55,57	0

6.5 Other polymers ⓘ

There are no such residues in this entry.