



# Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 02:15 PM BST

PDB ID : 1F3R  
Title : COMPLEX BETWEEN FV ANTIBODY FRAGMENT AND AN ANALOGUE OF THE MAIN IMMUNOGENIC REGION OF THE ACETYLCHOLINE RECEPTOR  
Authors : Kleinjung, J.; Petit, M.-C.; Orlewski, P.; Mamalaki, A.; Tzartos, S.-J.; Tsikaris, V.; Sakarellos-Daitsiotis, M.; Sakarellos, C.; Marraud, M.; Cung, M.-T.  
Deposited on : 2000-06-06

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	10	<div> <div>80%</div> <div>20%</div> </div>
2	B	257	<div> <div>81%</div> <div>17%</div> <div>.</div> </div>

## 2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3977 atoms, of which 1943 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called ACETYLCHOLINE RECEPTOR ALPHA.

Mol	Chain	Residues	Atoms					Trace
1	A	10	Total	C	H	N	O	0
			146	51	68	12	15	

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	70	GLY	ALA	ENGINEERED	GB 223528
A	76	NLE	LYS	ENGINEERED	GB 223528


- Molecule 2 is a protein called FV ANTIBODY FRAGMENT.


Mol	Chain	Residues	Atoms						Trace
2	B	257	Total	C	H	N	O	S	0
			3831	1232	1875	323	393	8	

## 4 Residue-property plots


These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


### • Molecule 1: ACETYLCHOLINE RECEPTOR ALPHA

Chain A: 



### • Molecule 2: FV ANTIBODY FRAGMENT

Chain B: 



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry simulated annealing docking molecular dynamics*.

Of the 1 calculated structures, 1 were deposited, based on the following criterion: *all calculated structures submitted*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	1.2
DISCOVER3	refinement	97

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality ⓘ

### 6.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NLE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.37	0/72 (0.0%)	1.47	1/98 (1.0%)
2	B	1.44	9/2003 (0.4%)	1.47	43/2715 (1.6%)
All	All	1.44	9/2075 (0.4%)	1.47	44/2813 (1.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
2	B	0	2
All	All	0	2

All bond outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	247	GLU	CD-OE2	9.97	1.36	1.25
2	B	180	GLU	CD-OE2	9.94	1.36	1.25
2	B	242	GLU	CD-OE2	9.88	1.36	1.25
2	B	253	GLU	CD-OE2	9.84	1.36	1.25
2	B	254	GLU	CD-OE2	9.76	1.36	1.25
2	B	219	GLU	CD-OE2	9.76	1.36	1.25
2	B	6	GLU	CD-OE2	9.71	1.36	1.25
2	B	46	GLU	CD-OE2	9.62	1.36	1.25
2	B	16	GLU	CD-OE2	9.54	1.36	1.25

All angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	71	ARG	NE-CZ-NH1	8.80	124.70	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	97	ARG	NE-CZ-NH1	8.76	124.68	120.30
2	B	66	ARG	NE-CZ-NH1	8.54	124.57	120.30
2	B	38	ARG	NE-CZ-NH1	8.37	124.49	120.30
2	B	13	ARG	NE-CZ-NH1	8.30	124.45	120.30
2	B	199	ARG	NE-CZ-NH1	8.30	124.45	120.30
2	B	156	ARG	NE-CZ-NH1	8.13	124.36	120.30
2	B	100	TYR	CB-CG-CD2	-8.00	116.20	121.00
2	B	72	ASP	CB-CG-OD2	-7.47	111.58	118.30
2	B	72	ASP	CB-CG-OD1	7.35	124.92	118.30
2	B	155	ASP	CB-CG-OD1	7.21	124.79	118.30
2	B	155	ASP	CB-CG-OD2	-7.08	111.93	118.30
2	B	54	ASP	CB-CG-OD2	-6.98	112.02	118.30
2	B	54	ASP	CB-CG-OD1	6.89	124.50	118.30
2	B	220	ASP	CB-CG-OD1	6.86	124.47	118.30
2	B	50	ARG	NE-CZ-NH1	6.82	123.71	120.30
2	B	208	ASP	CB-CG-OD2	-6.76	112.22	118.30
1	A	67	TRP	CD1-NE1-CE2	-6.57	103.09	109.00
2	B	220	ASP	CB-CG-OD2	-6.51	112.44	118.30
2	B	199	ARG	NE-CZ-NH2	-6.47	117.06	120.30
2	B	139	ASP	CB-CG-OD2	-6.47	112.48	118.30
2	B	88	ASP	CB-CG-OD2	-6.46	112.49	118.30
2	B	208	ASP	CB-CG-OD1	6.45	124.11	118.30
2	B	139	ASP	CB-CG-OD1	6.45	124.10	118.30
2	B	66	ARG	NE-CZ-NH2	-6.26	117.17	120.30
2	B	13	ARG	NE-CZ-NH2	-6.09	117.25	120.30
2	B	88	ASP	CB-CG-OD1	6.07	123.76	118.30
2	B	98	ASP	CB-CG-OD2	-6.05	112.85	118.30
2	B	100	TYR	CB-CG-CD1	5.92	124.55	121.00
2	B	255	ASP	CB-CG-OD1	5.91	123.61	118.30
2	B	255	ASP	CB-CG-OD2	-5.87	113.01	118.30
2	B	89	ASP	CB-CG-OD2	-5.83	113.05	118.30
2	B	52	TRP	CD1-NE1-CE2	-5.81	103.77	109.00
2	B	47	TRP	CD1-NE1-CE2	-5.67	103.89	109.00
2	B	97	ARG	NE-CZ-NH2	-5.65	117.48	120.30
2	B	173	TRP	CD1-NE1-CE2	-5.64	103.92	109.00
2	B	108	TRP	CD1-NE1-CE2	-5.64	103.93	109.00
2	B	113	TRP	CD1-NE1-CE2	-5.63	103.93	109.00
2	B	89	ASP	CB-CG-OD1	5.57	123.31	118.30
2	B	156	ARG	NE-CZ-NH2	-5.55	117.52	120.30
2	B	36	TRP	CD1-NE1-CE2	-5.46	104.08	109.00
2	B	38	ARG	NE-CZ-NH2	-5.28	117.66	120.30
2	B	71	ARG	NE-CZ-NH2	-5.25	117.68	120.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	98	ASP	CB-CG-OD1	5.14	122.92	118.30

There are no chirality outliers.

All planar outliers are listed below.

Mol	Chain	Res	Type	Group
2	B	59	TYR	Sidechain
2	B	100	TYR	Sidechain

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
2	B	1956	1875	1874	1
All	All	2034	1943	1939	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
2:B:161:CYS:SG	2:B:226:CYS:SG	0.53	3.07

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	8/10 (80%)	2 (25%)	5 (62%)	1 (12%)	<b>1</b>	<b>7</b>
2	B	255/257 (99%)	207 (81%)	42 (16%)	6 (2%)	<b>12</b>	49

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	263/267 (99%)	209 (79%)	47 (18%)	7 (3%)	10	45

All 7 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
2	B	131	GLY
2	B	125	GLY
2	B	55	GLY
2	B	109	TYR
2	B	168	ASN
2	B	127	GLY
1	A	69	PRO

### 6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	6/6 (100%)	6 (100%)	0 (0%)	100	100
2	B	213/213 (100%)	204 (96%)	9 (4%)	41	84
All	All	219/219 (100%)	210 (96%)	9 (4%)	42	84

All 9 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
2	B	142	LEU
2	B	98	ASP
2	B	220	ASP
2	B	75	LYS
2	B	6	GLU
2	B	242	GLU
2	B	1	GLN
2	B	193	GLN
2	B	178	LEU

### 6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	NLE	A	76	1	5,8,8	0.49	0 (0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	NLE	A	76	1	3,9,9	0.24	0 (0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	NLE	A	76	1	-	0,4,8,8	0,0,0,0

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided