



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:34 AM GMT

PDB ID : 3F7T  
Title : Structure of active IspH shows a novel fold with a [3Fe-4S] cluster in the catalytic centre  
Authors : Graewert, T.; Eppinger, J.; Rohdich, F.; Bacher, A.; Eisenreich, W.; Groll, M.  
Deposited on : 2008-11-10  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	
1	B	328	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PO4	A	901	-	-	X	X
4	PO4	B	906	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5280 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 4-hydroxy-3-methylbut-2-enyl diphosphate reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	310	Total	C	N	O	S	0	0	0
			2390	1492	428	460	10			
1	B	310	Total	C	N	O	S	0	0	0
			2390	1492	428	460	10			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	EXPRESSION TAG	UNP P62623
A	-10	ARG	-	EXPRESSION TAG	UNP P62623
A	-9	GLY	-	EXPRESSION TAG	UNP P62623
A	-8	SER	-	EXPRESSION TAG	UNP P62623
A	-7	HIS	-	EXPRESSION TAG	UNP P62623
A	-6	HIS	-	EXPRESSION TAG	UNP P62623
A	-5	HIS	-	EXPRESSION TAG	UNP P62623
A	-4	HIS	-	EXPRESSION TAG	UNP P62623
A	-3	HIS	-	EXPRESSION TAG	UNP P62623
A	-2	HIS	-	EXPRESSION TAG	UNP P62623
A	-1	GLY	-	EXPRESSION TAG	UNP P62623
A	0	SER	-	EXPRESSION TAG	UNP P62623
B	-11	MET	-	EXPRESSION TAG	UNP P62623
B	-10	ARG	-	EXPRESSION TAG	UNP P62623
B	-9	GLY	-	EXPRESSION TAG	UNP P62623
B	-8	SER	-	EXPRESSION TAG	UNP P62623
B	-7	HIS	-	EXPRESSION TAG	UNP P62623
B	-6	HIS	-	EXPRESSION TAG	UNP P62623
B	-5	HIS	-	EXPRESSION TAG	UNP P62623
B	-4	HIS	-	EXPRESSION TAG	UNP P62623
B	-3	HIS	-	EXPRESSION TAG	UNP P62623
B	-2	HIS	-	EXPRESSION TAG	UNP P62623
B	-1	GLY	-	EXPRESSION TAG	UNP P62623
B	0	SER	-	EXPRESSION TAG	UNP P62623

- Molecule 2 is FE3-S4 CLUSTER (three-letter code: F3S) (formula:  $\text{Fe}_3\text{S}_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			7	3	4		
2	B	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 3 is PYROPHOSPHATE 2- (three-letter code: POP) (formula:  $\text{H}_2\text{O}_7\text{P}_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			9	7	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	P	0	0
			9	7	2		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	213	Total 213	O 213	0	0
5	B	215	Total 215	O 215	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4-hydroxy-3-methylbut-2-enyl diphosphate reductase



- Molecule 1: 4-hydroxy-3-methylbut-2-enyl diphosphate reductase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.47Å 83.47Å 215.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	15.00 – 1.80 14.99 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.1 (15.00-1.80) 96.1 (14.99-1.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.53 (at 1.81Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.199 , 0.238 0.205 , 0.241	Depositor DCC
$R_{free}$ test set	3898 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.4	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 39.9	EDS
Estimated twinning fraction	0.031 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	1 of 78049 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5280	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, F3S, POP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	0/2425	0.91	8/3283 (0.2%)
1	B	0.71	0/2425	0.89	5/3283 (0.2%)
All	All	0.71	0/4850	0.90	13/6566 (0.2%)

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	9	ARG	NE-CZ-NH2	-11.11	114.74	120.30
1	B	81	ARG	NE-CZ-NH1	-9.87	115.36	120.30
1	A	9	ARG	NE-CZ-NH1	9.22	124.91	120.30
1	B	81	ARG	NE-CZ-NH2	7.54	124.07	120.30
1	A	230	ARG	NE-CZ-NH1	-6.92	116.84	120.30
1	A	281	ARG	NE-CZ-NH1	5.90	123.25	120.30
1	A	144	VAL	CG1-CB-CG2	5.81	120.20	110.90
1	A	118	LEU	CA-CB-CG	5.78	128.59	115.30
1	B	230	ARG	NE-CZ-NH1	-5.24	117.68	120.30
1	B	69	LEU	CA-CB-CG	5.19	127.23	115.30
1	A	144	VAL	CA-CB-CG1	5.14	118.62	110.90
1	A	244	LEU	CA-CB-CG	5.02	126.85	115.30
1	B	230	ARG	NE-CZ-NH2	5.02	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2390	0	2404	41	0
1	B	2390	0	2404	37	0
2	A	7	0	0	0	0
2	B	7	0	0	0	0
3	A	9	0	0	0	0
3	B	9	0	0	0	0
4	A	20	0	0	3	0
4	B	20	0	0	1	0
5	A	213	0	0	6	0
5	B	215	0	0	7	0
All	All	5280	0	4808	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:306:LYS:O	1:A:309:ARG:HG2	1.85	0.77
1:A:42:ASN:HD22	1:A:45:VAL:H	1.30	0.77
1:B:254:GLU:HG3	5:B:353:HOH:O	1.88	0.71
1:B:134:TYR:CB	1:B:141:MET:HG3	2.21	0.71
1:B:42:ASN:HB3	1:B:45:VAL:HG13	1.72	0.70
1:A:61:SER:HB3	5:A:444:HOH:O	1.91	0.70
1:B:103:HIS:HD2	5:B:362:HOH:O	1.74	0.69
1:B:171:VAL:H	1:B:204:GLN:HE22	1.39	0.68
1:B:42:ASN:HD22	1:B:45:VAL:H	1.42	0.68
1:A:42:ASN:HD21	1:A:44:TYR:HB3	1.60	0.65
1:A:9:ARG:NH2	4:A:901:PO4:O2	2.27	0.65
1:A:212:GLU:HB3	5:A:595:HOH:O	1.98	0.63
1:A:60:ILE:HD12	1:A:69:LEU:HD11	1.83	0.59
1:B:134:TYR:HB3	1:B:141:MET:HG3	1.83	0.59
1:A:151:TRP:HZ2	1:A:183:LYS:HD3	1.67	0.59
1:B:134:TYR:CG	1:B:141:MET:HG2	2.38	0.59
1:B:134:TYR:CB	1:B:141:MET:CG	2.83	0.56
1:A:103:HIS:HD2	5:A:416:HOH:O	1.88	0.56
1:B:157:ASN:HD21	1:B:159:GLU:HB2	1.71	0.55
1:A:1:MET:N	5:A:575:HOH:O	2.39	0.54
1:B:114:GLU:HG2	1:B:160:LYS:HB3	1.89	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:216:VAL:HG22	1:B:241:ARG:HB2	1.90	0.53
1:A:42:ASN:ND2	1:A:45:VAL:H	2.04	0.53
1:A:4:LEU:HD12	1:A:291:ILE:HB	1.90	0.53
1:A:4:LEU:HD22	1:A:261:CYS:SG	2.48	0.53
1:A:271:PRO:HD2	1:A:274:LEU:HD12	1.93	0.51
1:A:6:ALA:HB2	1:A:210:LEU:HD22	1.91	0.51
1:B:134:TYR:HB2	1:B:141:MET:HG3	1.92	0.51
1:A:4:LEU:HG	1:A:293:LEU:HD21	1.92	0.51
1:A:256:VAL:HG21	1:A:285:LEU:HB3	1.92	0.51
1:B:251:ILE:HD11	1:B:278:VAL:HG13	1.92	0.50
1:B:36:ARG:O	1:B:37:HIS:HB3	2.12	0.49
1:B:165:THR:CG2	1:B:166:GLN:O	2.60	0.48
1:A:256:VAL:HG11	1:A:285:LEU:HD23	1.96	0.48
1:A:68:ILE:HD13	1:A:90:THR:HB	1.94	0.48
1:A:251:ILE:HD11	1:A:278:VAL:HG13	1.95	0.48
1:A:136:ASN:C	1:A:136:ASN:HD22	2.17	0.47
1:B:1:MET:N	5:B:385:HOH:O	2.47	0.47
1:A:253:GLU:HG3	4:A:905:PO4:O2	2.16	0.46
1:A:151:TRP:CZ2	1:A:183:LYS:HD3	2.49	0.46
1:A:281:ARG:HD3	5:A:515:HOH:O	2.15	0.45
1:A:281:ARG:HH21	1:A:284:GLN:HG3	1.82	0.45
1:A:192:ARG:HH11	1:A:192:ARG:HD2	1.66	0.45
1:B:57:ILE:HG13	1:B:62:GLU:HB2	1.98	0.45
1:B:165:THR:HB	1:B:195:ASP:H	1.82	0.44
1:B:134:TYR:HB2	1:B:141:MET:CG	2.46	0.44
1:A:36:ARG:HD2	1:A:60:ILE:HD11	1.98	0.44
1:B:184:ARG:NE	5:B:406:HOH:O	2.35	0.44
1:B:165:THR:HG22	1:B:166:GLN:O	2.18	0.44
1:A:5:LEU:HD12	1:A:264:VAL:HG13	2.00	0.43
1:A:230:ARG:HD3	1:A:230:ARG:HA	1.69	0.43
1:B:252:GLN:NE2	5:B:387:HOH:O	2.48	0.43
1:A:292:PRO:HB3	4:A:901:PO4:O2	2.18	0.43
1:B:252:GLN:HB2	1:B:255:TRP:CE2	2.53	0.43
1:B:82:ASN:ND2	5:B:317:HOH:O	2.51	0.43
1:B:4:LEU:HD22	1:B:261:CYS:SG	2.57	0.43
1:B:159:GLU:OE2	1:B:187:LYS:HD2	2.19	0.42
1:B:231:LEU:HD13	5:B:369:HOH:O	2.18	0.42
1:A:117:ILE:O	1:A:163:PHE:HA	2.19	0.42
1:A:212:GLU:CB	5:A:595:HOH:O	2.61	0.42
1:A:8:PRO:HG2	1:A:202:ASN:HB3	2.01	0.42
1:B:292:PRO:HB3	4:B:906:PO4:O2	2.19	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:CYS:SG	1:A:15:VAL:HG23	2.60	0.42
1:B:203:ARG:HG3	1:B:203:ARG:NH2	2.35	0.42
1:A:15:VAL:HG11	1:A:269:SER:HA	2.01	0.41
1:A:234:LEU:O	1:A:238:MET:HG2	2.20	0.41
1:B:69:LEU:O	1:B:91:VAL:HA	2.21	0.41
1:A:42:ASN:ND2	1:A:44:TYR:HB3	2.33	0.41
1:B:236:GLN:HB2	1:B:236:GLN:HE21	1.66	0.41
1:B:42:ASN:HD21	1:B:44:TYR:HB3	1.86	0.41
1:A:136:ASN:ND2	1:A:138:GLU:H	2.18	0.41
1:B:270:ALA:HA	1:B:271:PRO:HD3	1.94	0.41
1:B:48:SER:O	1:B:51:GLU:HB3	2.21	0.41
1:B:306:LYS:HA	1:B:309:ARG:HG3	2.03	0.41
1:A:233:GLU:O	1:A:237:ARG:HD3	2.20	0.41
1:A:270:ALA:HA	1:A:271:PRO:HD3	1.90	0.40
1:A:257:LYS:HB3	1:A:258:GLU:H	1.42	0.40
1:B:36:ARG:O	1:B:37:HIS:CB	2.69	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/328 (94%)	303 (98%)	5 (2%)	0	100	100
1	B	308/328 (94%)	303 (98%)	5 (2%)	0	100	100
All	All	616/656 (94%)	606 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/274 (94%)	236 (92%)	22 (8%)	13	3
1	B	258/274 (94%)	232 (90%)	26 (10%)	9	2
All	All	516/548 (94%)	468 (91%)	48 (9%)	11	3

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LEU
1	A	5	LEU
1	A	16	ASP
1	A	69	LEU
1	A	91	VAL
1	A	98	LEU
1	A	103	HIS
1	A	118	LEU
1	A	136	ASN
1	A	144	VAL
1	A	153	LEU
1	A	161	LEU
1	A	213	GLN
1	A	230	ARG
1	A	231	LEU
1	A	236	GLN
1	A	238	MET
1	A	240	LYS
1	A	244	LEU
1	A	264	VAL
1	A	285	LEU
1	A	310	VAL
1	B	4	LEU
1	B	5	LEU
1	B	45	VAL
1	B	57	ILE
1	B	69	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	81	ARG
1	B	91	VAL
1	B	98	LEU
1	B	103	HIS
1	B	108	ARG
1	B	118	LEU
1	B	136	ASN
1	B	141	MET
1	B	143	LEU
1	B	153	LEU
1	B	157	ASN
1	B	230	ARG
1	B	231	LEU
1	B	238	MET
1	B	240	LYS
1	B	246	ASP
1	B	258	GLU
1	B	273	ILE
1	B	285	LEU
1	B	296	ARG
1	B	304	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	ASN
1	A	103	HIS
1	A	136	ASN
1	A	229	ASN
1	A	236	GLN
1	A	252	GLN
1	A	277	ASN
1	A	283	GLN
1	B	24	ASN
1	B	42	ASN
1	B	82	ASN
1	B	103	HIS
1	B	136	ASN
1	B	157	ASN
1	B	202	ASN
1	B	204	GLN
1	B	213	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	229	ASN
1	B	236	GLN
1	B	283	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	PO4	A	901	-	4,4,4	0.52	0	6,6,6	0.26	0
4	PO4	A	902	-	4,4,4	0.43	0	6,6,6	0.28	0
4	PO4	A	904	-	4,4,4	0.35	0	6,6,6	0.27	0
4	PO4	A	905	-	4,4,4	0.44	0	6,6,6	0.27	0
3	POP	A	998	-	8,8,8	1.04	0	13,13,13	0.98	1 (7%)
2	F3S	A	999	1	0,9,9	0.00	-	0,15,15	0.00	-
4	PO4	B	903	-	4,4,4	0.58	0	6,6,6	0.28	0
4	PO4	B	906	-	4,4,4	0.51	0	6,6,6	0.28	0
4	PO4	B	907	-	4,4,4	0.43	0	6,6,6	0.29	0
4	PO4	B	908	-	4,4,4	0.30	0	6,6,6	0.29	0



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	POP	B	998	-	8,8,8	0.94	0	13,13,13	1.38	1 (7%)
2	F3S	B	999	1	0,9,9	0.00	-	0,15,15	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PO4	A	901	-	-	0/0/0/0	0/0/0/0
4	PO4	A	902	-	-	0/0/0/0	0/0/0/0
4	PO4	A	904	-	-	0/0/0/0	0/0/0/0
4	PO4	A	905	-	-	0/0/0/0	0/0/0/0
3	POP	A	998	-	-	0/6/6/6	0/0/0/0
2	F3S	A	999	1	-	0/0/24/24	0/0/3/3
4	PO4	B	903	-	-	0/0/0/0	0/0/0/0
4	PO4	B	906	-	-	0/0/0/0	0/0/0/0
4	PO4	B	907	-	-	0/0/0/0	0/0/0/0
4	PO4	B	908	-	-	0/0/0/0	0/0/0/0
3	POP	B	998	-	-	0/6/6/6	0/0/0/0
2	F3S	B	999	1	-	0/0/24/24	0/0/3/3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	998	POP	O6-P2-O	-3.04	93.62	106.09
3	A	998	POP	O3-P1-O	2.34	115.69	106.09

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	901	PO4	2	0
4	A	905	PO4	1	0
4	B	906	PO4	1	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	310/328 (94%)	-0.05	6 (1%) 70 66	26, 33, 43, 52	0
1	B	310/328 (94%)	0.04	7 (2%) 64 59	25, 33, 44, 55	0
All	All	620/656 (94%)	-0.00	13 (2%) 67 62	25, 33, 44, 55	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	310	VAL	4.4
1	A	309	ARG	3.4
1	B	310	VAL	3.2
1	B	309	ARG	2.9
1	B	289	GLU	2.6
1	A	258	GLU	2.5
1	A	58	GLU	2.4
1	B	65	ASP	2.2
1	B	258	GLU	2.2
1	A	289	GLU	2.1
1	A	308	LEU	2.1
1	B	260	LYS	2.1
1	B	306	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	PO4	A	901	5/5	0.97	0.24	8.59	63,63,63,64	0
4	PO4	B	906	5/5	0.97	0.23	7.54	72,72,72,73	0
2	F3S	A	999	7/7	0.92	0.11	1.34	18,21,23,23	0
4	PO4	A	902	5/5	0.86	0.24	1.18	73,73,74,74	0
4	PO4	B	903	5/5	0.98	0.11	0.37	42,43,46,48	0
2	F3S	B	999	7/7	0.94	0.09	-0.33	19,22,23,23	0
3	POP	A	998	9/9	0.99	0.04	-2.19	24,27,31,31	0
3	POP	B	998	9/9	0.99	0.04	-2.40	20,26,28,31	0
4	PO4	A	905	5/5	0.90	0.30	-	81,82,82,82	0
4	PO4	B	907	5/5	0.96	0.25	-	62,63,64,64	0
4	PO4	B	908	5/5	0.94	0.31	-	75,77,77,77	0
4	PO4	A	904	5/5	0.91	0.31	-	99,99,99,100	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.