



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 29, 2016 – 01:40 PM EDT

PDB ID : 5FA2  
Title : Crystal structure of 426c.TM4deltaV1-3 p120  
Authors : Scharf, L.; Bjorkman, P.J.  
Deposited on : 2015-12-10  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027107  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027107







Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	601	-	-	-	X
2	NAG	A	604	-	-	-	X
2	NAG	B	601	-	-	-	X
2	NAG	B	602	-	-	-	X
5	FLC	A	611	-	-	-	X
5	FLC	A	612	-	-	-	X
5	FLC	B	611	-	-	-	X
6	SCN	A	613	-	-	-	X



## 2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 5875 atoms, of which 15 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called gp120.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	336	Total	C	N	O	S	0	0	0
			2570	1614	443	491	22			
1	B	336	Total	C	N	O	S	0	0	0
			2554	1605	439	488	22			

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		


*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 3 is BETA-D-MANNOSE (three-letter code: BMA) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		


- Molecule 4 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	B	1	Total	C	O	0	0
			11	6	5		


- Molecule 5 is CITRATE ANION (three-letter code: FLC) (formula: C<sub>6</sub>H<sub>5</sub>O<sub>7</sub>).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			13	6	7		
5	A	1	Total	C	H	O	0
			18	6	5	7	
5	B	1	Total	C	O	0	0
			13	6	7		
5	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 6 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	230	Total	H	O	0	0
			232	2	230		
7	B	186	Total	H	O	0	0
			194	8	186		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: gp120



#### • Molecule 1: gp120





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	144.85Å 85.91Å 90.04Å 90.00° 104.77° 90.00°	Depositor
Resolution (Å)	35.80 – 2.00 38.52 – 1.81	Depositor EDS
% Data completeness (in resolution range)	97.4 (35.80-2.00) 80.5 (38.52-1.81)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.88 (at 1.82Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.207 , 0.232 0.205 , 0.232	Depositor DCC
$R_{free}$ test set	1451 reflections (2.25%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.3	Xtriage
Anisotropy	0.357	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 45.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 88277 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5875	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FLC, SCN, NAG, BMA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.58	0/2622	0.72	0/3568
1	B	0.57	0/2605	0.72	0/3546
All	All	0.57	0/5227	0.72	0/7114

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2570	0	2462	29	0
1	B	2554	0	2438	36	0
2	A	98	0	89	2	0
2	B	112	0	102	2	0
3	A	11	0	8	0	0
3	B	11	0	9	0	0
4	A	22	0	20	2	0
4	B	11	0	10	0	0
5	A	26	5	10	2	0
5	B	26	0	10	4	0
6	A	3	0	0	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	230	2	0	3	0
7	B	186	8	0	2	0
All	All	5860	15	5158	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 70 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:602:NAG:O7	2:B:602:NAG:O3	1.98	0.81
1:A:65:CYS:SG	1:A:115:CYS:HB3	2.20	0.80
1:A:65:CYS:HB2	1:A:115:CYS:SG	2.21	0.80
1:B:65:CYS:HB2	1:B:115:CYS:SG	2.22	0.80
1:A:432:LYS:NZ	5:A:611:FLC:OG1	2.13	0.76

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	332/353 (94%)	324 (98%)	8 (2%)	0	100	100
1	B	332/353 (94%)	325 (98%)	7 (2%)	0	100	100
All	All	664/706 (94%)	649 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	283/313 (90%)	283 (100%)	0	100	100
1	B	279/313 (89%)	279 (100%)	0	100	100
All	All	562/626 (90%)	562 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

25 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	601	1	14,14,15	0.28	0	15,19,21	0.53	0
2	NAG	A	602	1	14,14,15	0.35	0	15,19,21	0.55	0
2	NAG	A	603	1	14,14,15	0.28	0	15,19,21	0.54	0
2	NAG	A	604	1	14,14,15	0.24	0	15,19,21	0.59	1 (6%)
2	NAG	A	605	1	14,14,15	0.27	0	15,19,21	0.57	0
2	NAG	A	606	1,2	14,14,15	0.19	0	15,19,21	0.71	1 (6%)
2	NAG	A	607	3,2	14,14,15	0.23	0	15,19,21	0.60	1 (6%)
3	BMA	A	608	2,4	11,11,12	0.79	1 (9%)	15,15,17	0.98	1 (6%)
4	MAN	A	609	3	11,11,12	0.26	0	15,15,17	0.58	0
4	MAN	A	610	3	11,11,12	0.65	0	15,15,17	0.96	1 (6%)
5	FLC	A	611	-	3,12,12	1.04	0	3,17,17	2.83	2 (66%)
5	FLC	A	612	-	3,12,12	1.04	0	3,17,17	2.34	2 (66%)
6	SCN	A	613	-	2,2,2	0.37	0	1,1,1	0.28	0
2	NAG	B	601	1	14,14,15	0.38	0	15,19,21	0.34	0
2	NAG	B	602	1	14,14,15	0.37	0	15,19,21	0.69	0
2	NAG	B	603	1	14,14,15	0.28	0	15,19,21	0.53	0
2	NAG	B	604	1	14,14,15	0.49	0	15,19,21	0.80	0
2	NAG	B	605	1	14,14,15	0.35	0	15,19,21	0.61	0
2	NAG	B	606	1	14,14,15	0.21	0	15,19,21	0.38	0
2	NAG	B	607	1,2	14,14,15	0.20	0	15,19,21	0.68	1 (6%)
2	NAG	B	608	3,2	14,14,15	0.22	0	15,19,21	0.55	0
3	BMA	B	609	2,4	11,11,12	0.56	0	15,15,17	0.78	0
4	MAN	B	610	3	11,11,12	0.26	0	15,15,17	0.58	0
5	FLC	B	611	-	3,12,12	1.01	0	3,17,17	2.43	2 (66%)
5	FLC	B	612	-	3,12,12	1.48	1 (33%)	3,17,17	1.81	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	601	1	-	0/6/23/26	0/1/1/1
2	NAG	A	602	1	-	0/6/23/26	0/1/1/1
2	NAG	A	603	1	-	0/6/23/26	0/1/1/1
2	NAG	A	604	1	-	0/6/23/26	0/1/1/1
2	NAG	A	605	1	-	0/6/23/26	0/1/1/1
2	NAG	A	606	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	607	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	608	2,4	-	0/2/19/22	0/1/1/1

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MAN	A	609	3	-	0/2/19/22	0/1/1/1
4	MAN	A	610	3	-	0/2/19/22	0/1/1/1
5	FLC	A	611	-	-	0/6/16/16	0/0/0/0
5	FLC	A	612	-	-	0/6/16/16	0/0/0/0
6	SCN	A	613	-	-	0/0/0/0	0/0/0/0
2	NAG	B	601	1	-	0/6/23/26	0/1/1/1
2	NAG	B	602	1	-	0/6/23/26	0/1/1/1
2	NAG	B	603	1	-	0/6/23/26	0/1/1/1
2	NAG	B	604	1	-	0/6/23/26	0/1/1/1
2	NAG	B	605	1	-	0/6/23/26	0/1/1/1
2	NAG	B	606	1	-	0/6/23/26	0/1/1/1
2	NAG	B	607	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	608	3,2	-	0/6/23/26	0/1/1/1
3	BMA	B	609	2,4	-	0/2/19/22	0/1/1/1
4	MAN	B	610	3	-	0/2/19/22	0/1/1/1
5	FLC	B	611	-	-	0/6/16/16	0/0/0/0
5	FLC	B	612	-	-	0/6/16/16	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	608	BMA	O5-C1	-2.11	1.40	1.43
5	B	612	FLC	OHB-CB	2.47	1.46	1.43

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	611	FLC	CB-CG-CGC	-3.47	109.53	114.95
5	B	611	FLC	CB-CG-CGC	-3.44	109.57	114.95
5	A	611	FLC	CB-CA-CAC	-3.43	109.59	114.95
5	A	612	FLC	CB-CA-CAC	-3.26	109.86	114.95
5	B	611	FLC	CB-CA-CAC	-2.40	111.20	114.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	NAG	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	606	NAG	1	0
2	A	607	NAG	1	0
4	A	609	MAN	2	0
5	A	611	FLC	2	0
6	A	613	SCN	1	0
2	B	602	NAG	2	0
5	B	611	FLC	3	0
5	B	612	FLC	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	336/353 (95%)	-0.25	10 (2%) 54 55	28, 42, 74, 89	0
1	B	336/353 (95%)	-0.11	16 (4%) 34 36	32, 48, 73, 94	0
All	All	672/706 (95%)	-0.18	26 (3%) 43 45	28, 45, 73, 94	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	462	THR	7.7
1	B	461	THR	4.1
1	A	462	THR	4.1
1	B	48	ALA	3.9
1	A	88	GLN	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	FLC	A	612	13/13	0.82	0.20	8.94	53,58,70,72	18
5	FLC	B	611	13/13	0.78	0.32	5.98	40,46,56,59	13
2	NAG	A	601	14/15	0.84	0.22	5.53	61,67,71,79	0
5	FLC	A	611	13/13	0.72	0.26	5.17	34,43,53,53	13
2	NAG	B	601	14/15	0.72	0.23	3.65	68,77,85,89	0
6	SCN	A	613	3/3	0.61	0.24	3.11	36,36,44,59	0
2	NAG	B	602	14/15	0.78	0.19	3.09	43,60,71,75	0
2	NAG	A	604	14/15	0.86	0.15	2.15	55,59,65,68	0
2	NAG	B	606	14/15	0.76	0.35	1.90	85,91,100,102	0
5	FLC	B	612	13/13	0.83	0.16	1.65	48,56,59,66	13
2	NAG	A	603	14/15	0.88	0.12	0.06	42,45,50,52	0
2	NAG	B	607	14/15	0.90	0.10	-0.01	41,48,57,59	0
2	NAG	A	606	14/15	0.92	0.11	-0.44	37,38,44,52	0
2	NAG	B	603	14/15	0.76	0.21	-	74,84,88,90	0
2	NAG	A	605	14/15	0.86	0.31	-	65,72,80,81	0
3	BMA	B	609	11/12	0.68	0.29	-	89,95,107,114	0
4	MAN	A	609	11/12	0.70	0.44	-	98,101,111,120	0
2	NAG	A	602	14/15	0.89	0.16	-	56,60,65,67	0
2	NAG	A	607	14/15	0.93	0.08	-	46,48,57,63	0
3	BMA	A	608	11/12	0.77	0.20	-	72,87,96,98	0
4	MAN	B	610	11/12	0.63	0.56	-	118,125,132,136	0
4	MAN	A	610	11/12	0.17	0.41	-	104,109,112,113	0
2	NAG	B	608	14/15	0.85	0.15	-	57,64,74,81	0
2	NAG	B	605	14/15	0.86	0.40	-	71,74,81,86	0
2	NAG	B	604	14/15	0.81	0.28	-	84,90,97,103	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.