



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:34 PM GMT

PDB ID : 4FCS  
Title : The crystal structures of several mutants of pleurotus eryngii versatile peroxidase  
Authors : Mate, M.J.; Romero, A.; Ruiz-Duenas, F.J.; Martinez, A.T.  
Deposited on : 2012-05-25  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	<div> <div>4%</div> <div>83%</div> <div>15%</div> <div>.</div> </div>

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2763 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Versatile peroxidase VPL2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	315	2338	1471	394	461	12	0	3	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	176	GLY	LYS	ENGINEERED MUTATION	UNP O94753
A	191	GLU	GLY	ENGINEERED MUTATION	UNP O94753

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ca	0	0
			2	2		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	363	Total	O	0	0
			363	363		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Versatile peroxidase VPL2



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 41	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.27Å 96.27Å 98.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.14 – 1.50 48.14 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.14-1.50) 99.9 (48.14-1.50)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.71 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.150 , 0.176 0.149 , 0.175	Depositor DCC
$R_{free}$ test set	3635 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	12.0	Xtriage
Anisotropy	0.205	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 47.7	EDS
Estimated twinning fraction	0.006 for l,-k,h 0.015 for -l,-k,-h 0.012 for -h,-l,-k 0.007 for -h,l,k 0.032 for -h,k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 71863 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2763	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, SO4, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.64	19/2411 (0.8%)	1.52	35/3288 (1.1%)

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	24	ILE	CB-CG1	-7.57	1.32	1.54
1	A	43	ARG	CZ-NH2	7.45	1.42	1.33
1	A	12	ALA	CA-CB	7.41	1.68	1.52
1	A	161	GLU	CD-OE2	-7.09	1.17	1.25
1	A	225	GLU	CD-OE1	6.67	1.32	1.25
1	A	170	SER	CB-OG	6.45	1.50	1.42
1	A	140	GLU	CD-OE1	6.17	1.32	1.25
1	A	40[A]	GLU	CB-CG	6.15	1.63	1.52
1	A	40[B]	GLU	CB-CG	6.15	1.63	1.52
1	A	272	LYS	CB-CG	-6.13	1.35	1.52
1	A	263	SER	CB-OG	6.11	1.50	1.42
1	A	24	ILE	CA-CB	-6.07	1.40	1.54
1	A	242	CYS	CB-SG	5.98	1.92	1.82
1	A	217	GLU	CD-OE2	5.88	1.32	1.25
1	A	14	CYS	CB-SG	-5.62	1.72	1.81
1	A	281	VAL	CB-CG2	5.37	1.64	1.52
1	A	43	ARG	CD-NE	5.20	1.55	1.46
1	A	304	GLU	CD-OE2	5.07	1.31	1.25
1	A	122	PHE	CE2-CZ	5.07	1.47	1.37

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	227	ARG	NE-CZ-NH1	11.33	125.96	120.30
1	A	280	ASP	CB-CG-OD1	9.61	126.95	118.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	236	ARG	NE-CZ-NH1	-9.25	115.68	120.30
1	A	101	ASP	CB-CG-OD1	9.14	126.53	118.30
1	A	75	PHE	CB-CG-CD2	-9.12	114.41	120.80
1	A	43	ARG	NE-CZ-NH2	8.15	124.37	120.30
1	A	175	ASP	CB-CG-OD1	7.74	125.26	118.30
1	A	151	ARG	NE-CZ-NH2	-7.70	116.45	120.30
1	A	187	ASP	CB-CG-OD1	7.64	125.18	118.30
1	A	24	ILE	CB-CG1-CD1	-7.63	92.55	113.90
1	A	69	ASP	CB-CG-OD1	7.42	124.97	118.30
1	A	236	ARG	NE-CZ-NH2	7.07	123.83	120.30
1	A	122	PHE	CB-CG-CD2	-6.70	116.11	120.80
1	A	69	ASP	CB-CG-OD2	-6.46	112.49	118.30
1	A	272	LYS	CD-CE-NZ	6.39	126.41	111.70
1	A	257[A]	ARG	NE-CZ-NH1	6.36	123.48	120.30
1	A	257[B]	ARG	NE-CZ-NH1	6.36	123.48	120.30
1	A	187	ASP	CB-CG-OD2	-6.33	112.61	118.30
1	A	178	ASP	CB-CG-OD2	-6.21	112.71	118.30
1	A	151	ARG	NE-CZ-NH1	6.03	123.31	120.30
1	A	217	GLU	OE1-CD-OE2	-5.95	116.16	123.30
1	A	22	ASP	CB-CG-OD1	5.91	123.61	118.30
1	A	258	PHE	CB-CG-CD1	-5.90	116.67	120.80
1	A	37	GLU	CG-CD-OE1	5.78	129.86	118.30
1	A	82	ASP	CB-CG-OD1	5.76	123.49	118.30
1	A	140	GLU	OE1-CD-OE2	-5.68	116.48	123.30
1	A	68	PHE	CB-CG-CD1	-5.59	116.89	120.80
1	A	220	SER	N-CA-CB	-5.46	102.31	110.50
1	A	30	ASP	CB-CG-OD1	-5.46	113.39	118.30
1	A	271	ASP	CB-CG-OD2	-5.43	113.41	118.30
1	A	68	PHE	CB-CG-CD2	5.31	124.52	120.80
1	A	154	ASP	CB-CG-OD1	-5.23	113.59	118.30
1	A	119	ARG	NE-CZ-NH2	-5.17	117.71	120.30
1	A	215	LYS	N-CA-CB	-5.14	101.34	110.60
1	A	146	ASP	CB-CG-OD1	5.04	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2338	0	2237	15	0
2	A	43	0	30	3	0
3	A	2	0	0	0	0
4	A	5	0	0	0	0
5	A	12	0	16	0	0
6	A	363	0	0	4	0
All	All	2763	0	2283	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:229:GLN:HE22	1:A:232:HIS:HD2	1.24	0.85
1:A:36:GLU:OE1	6:A:684:HOH:O	2.06	0.74
1:A:25:GLN:HE21	1:A:25:GLN:HA	1.53	0.72
1:A:229:GLN:NE2	1:A:232:HIS:HD2	1.89	0.70
2:A:401:HEM:HBB2	2:A:401:HEM:HHC	1.75	0.68
1:A:16:ILE:HD12	1:A:16:ILE:C	2.26	0.56
1:A:40[B]:GLU:OE2	2:A:401:HEM:O1D	2.23	0.55
1:A:229:GLN:HA	1:A:229:GLN:HE21	1.79	0.47
1:A:173:ALA:HA	1:A:185:PRO:HA	1.97	0.46
1:A:24:ILE:CD1	1:A:29:PHE:CE2	2.99	0.46
1:A:43:ARG:NE	6:A:840:HOH:O	2.38	0.45
1:A:295:PRO:HB2	1:A:298:PHE:CD1	2.52	0.45
1:A:43:ARG:NH1	6:A:840:HOH:O	2.47	0.43
1:A:229:GLN:NE2	1:A:232:HIS:CD2	2.79	0.43
1:A:97:ILE:HG22	6:A:701:HOH:O	2.18	0.43
1:A:165:LEU:HB3	2:A:401:HEM:HMC3	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	316/315 (100%)	311 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	252/249 (101%)	247 (98%)	5 (2%)	63	29

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	ILE
1	A	25	GLN
1	A	55	THR
1	A	81	ILE
1	A	229	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	A	25	GLN
1	A	88	GLN
1	A	104	GLN
1	A	113	ASN
1	A	229	GLN
1	A	232	HIS
1	A	249	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HEM	A	401	1,6	30,50,50	2.88	13 (43%)	24,82,82	2.85	10 (41%)
4	SO4	A	404	-	4,4,4	0.70	0	6,6,6	0.57	0
5	GOL	A	405	-	5,5,5	1.30	1 (20%)	5,5,5	1.82	3 (60%)
5	GOL	A	406	-	5,5,5	1.20	1 (20%)	5,5,5	1.61	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	401	1,6	-	0/10/54/54	0/0/8/8
4	SO4	A	404	-	-	0/0/0/0	0/0/0/0
5	GOL	A	405	-	-	0/4/4/4	0/0/0/0
5	GOL	A	406	-	-	0/4/4/4	0/0/0/0

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	HEM	C3B-C4B	-8.84	1.44	1.51
2	A	401	HEM	C3D-C4D	-4.37	1.46	1.51
2	A	401	HEM	C2B-C1B	-4.07	1.38	1.51
2	A	401	HEM	C2D-C3D	-3.30	1.44	1.54
2	A	401	HEM	C2D-C1D	-3.26	1.41	1.51
2	A	401	HEM	CAD-C3D	-2.73	1.48	1.54
2	A	401	HEM	CMB-C2B	2.05	1.57	1.53
5	A	405	GOL	C3-C2	2.07	1.60	1.52
2	A	401	HEM	C3C-CAC	2.08	1.55	1.51
5	A	406	GOL	O1-C1	2.14	1.51	1.42
2	A	401	HEM	CMA-C3A	2.70	1.57	1.51
2	A	401	HEM	C1C-NC	2.84	1.39	1.36
2	A	401	HEM	FE-NB	2.84	2.12	1.97
2	A	401	HEM	FE-NC	3.31	2.08	1.95
2	A	401	HEM	C4C-NC	6.74	1.44	1.36

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	HEM	CMA-C3A-C4A	-5.70	118.94	128.36
2	A	401	HEM	C2C-C1C-NC	-3.56	104.20	110.21
2	A	401	HEM	C4B-CHC-C1C	-3.22	120.44	125.82
2	A	401	HEM	CAA-CBA-CGA	-2.74	107.72	112.75
5	A	405	GOL	C3-C2-C1	-2.44	101.56	111.12
5	A	405	GOL	O2-C2-C3	-2.02	99.37	108.65
5	A	405	GOL	O3-C3-C2	2.01	119.95	110.18
2	A	401	HEM	CMD-C2D-C3D	2.17	123.93	114.35
2	A	401	HEM	C3B-CAB-CBB	2.95	128.98	124.46
5	A	406	GOL	O2-C2-C3	2.97	122.25	108.65
2	A	401	HEM	CMB-C2B-C3B	4.65	128.14	116.53
2	A	401	HEM	CMC-C2C-C3C	4.80	128.52	116.53
2	A	401	HEM	CAD-C3D-C4D	4.99	130.06	112.47
2	A	401	HEM	CAD-C3D-C2D	5.83	129.99	113.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	HEM	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	315/315 (100%)	-0.08	14 (4%) 38 40	6, 11, 26, 46	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	ASP	5.9
1	A	2	THR	4.0
1	A	3	CYS	3.8
1	A	213	ASP	3.8
1	A	1	ALA	3.4
1	A	308	ALA	3.2
1	A	309	ALA	3.2
1	A	6	GLY	3.1
1	A	5	ASP	3.0
1	A	96	ASN	3.0
1	A	311	PRO	2.6
1	A	16	ILE	2.4
1	A	314	ALA	2.3
1	A	8	THR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	A	406	6/6	0.94	0.11	1.64	17,23,26,27	0
5	GOL	A	405	6/6	0.94	0.08	0.81	13,20,22,23	0
3	CA	A	402	1/1	1.00	0.07	0.30	7,7,7,7	0
2	HEM	A	401	43/43	0.98	0.08	-0.05	6,8,13,20	0
3	CA	A	403	1/1	1.00	0.08	-0.30	7,7,7,7	0
4	SO4	A	404	5/5	0.99	0.06	-1.31	11,12,13,15	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.