



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2016 – 07:48 PM EDT

PDB ID : 5FKA
Title : Crystal structure of staphylococcal enterotoxin E in complex with a T cell receptor
Authors : Rodstrom, K.E.J.; Regenthal, P.; Lindkvist-Petersson, K.
Deposited on : 2015-10-15
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027674
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	206	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>8%</div> <div>• •</div> </div> </div>
2	B	243	<div> <div>87%</div> <div>11%</div> <div>• •</div> </div>
3	C	233	<div> <div>4%</div> <div> <div></div> <div>84%</div> <div>11%</div> <div>5%</div> </div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5197 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T CELL RECEPTOR ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	197	Total	C	N	O	S	0	0	0
			1482	935	245	295	7			

- Molecule 2 is a protein called T CELL RECEPTOR BETA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	240	Total	C	N	O	S	0	0	0
			1885	1187	336	356	6			

- Molecule 3 is a protein called STAPHYLOCOCCAL ENTEROTOXIN E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	221	Total	C	N	O	S	0	0	0
			1735	1106	298	328	3			

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	1	Total	Zn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	28	Total	O	0	0
			28	28		
5	B	36	Total	O	0	0
			36	36		
5	C	30	Total	O	0	0
			30	30		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: T CELL RECEPTOR ALPHA CHAIN



• Molecule 2: T CELL RECEPTOR BETA CHAIN



• Molecule 3: STAPHYLOCOCCAL ENTEROTOXIN E



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	113.29 Å 150.14 Å 39.16 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.78 – 2.40 45.78 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.7 (45.78-2.40) 98.8 (45.78-2.40)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.39 Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.212 , 0.242 0.222 , 0.258	Depositor DCC
R_{free} test set	1336 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	37.2	Xtriage
Anisotropy	0.140	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5197	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/1511	0.63	0/2057
2	B	0.34	0/1936	0.62	0/2636
3	C	0.37	0/1771	0.63	0/2397
All	All	0.36	0/5218	0.63	0/7090

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1482	0	1368	10	0
2	B	1885	0	1769	12	0
3	C	1735	0	1659	8	0
4	C	1	0	0	0	0
5	A	28	0	0	0	0
5	B	36	0	0	0	0
5	C	30	0	0	0	0
All	All	5197	0	4796	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:164:HIS:HD2	3:C:170:TYR:H	1.35	0.75
1:A:30:VAL:HG21	1:A:90:VAL:HB	1.77	0.66
1:A:50:ILE:HG12	1:A:64:THR:HG23	1.84	0.60
2:B:155:GLU:HB3	2:B:212:GLN:HB3	1.87	0.56
2:B:222:TRP:HB2	2:B:228:LYS:HD3	1.89	0.55
3:C:163:LEU:HD22	3:C:230:LEU:HD11	1.90	0.53
2:B:65:ARG:NH2	2:B:87:ASP:OD2	2.41	0.53
2:B:31:ASN:HD21	2:B:98:GLY:H	1.55	0.53
2:B:228:LYS:HG3	2:B:230:VAL:HG13	1.91	0.52
1:A:50:ILE:O	1:A:64:THR:HG21	2.11	0.50
3:C:111:VAL:H	3:C:216:ASN:HD21	1.61	0.48
2:B:210:GLN:HG3	2:B:233:ILE:HG23	1.94	0.48
2:B:8:ASN:HD21	2:B:23:ARG:HD3	1.77	0.48
3:C:12:LEU:HD21	3:C:183:LEU:HG	1.94	0.48
1:A:143:GLN:HE21	1:A:143:GLN:H	1.63	0.47
1:A:7:GLN:HE21	1:A:103:GLY:HA3	1.80	0.46
2:B:152:ASP:HB2	2:B:175:PRO:HG2	1.97	0.46
1:A:32:ASN:HD21	1:A:34:GLN:HE21	1.64	0.46
1:A:34:GLN:HE22	2:B:102:GLN:H	1.64	0.46
1:A:14:LEU:HD11	1:A:20:SER:HB2	1.99	0.45
1:A:153:ASP:HB2	1:A:180:LYS:HE2	1.98	0.45
3:C:72:GLY:HA2	3:C:214:ARG:HD3	2.00	0.43
2:B:85:GLN:HE22	2:B:114:GLU:HG3	1.84	0.43
2:B:111:THR:HG21	2:B:151:PRO:HB3	2.01	0.42
2:B:126:VAL:HG23	2:B:236:ALA:HB3	2.02	0.42
3:C:38:THR:HG21	3:C:57:PHE:HE1	1.84	0.42
3:C:179:VAL:HG11	3:C:230:LEU:HD22	2.02	0.41
3:C:145:THR:HG21	3:C:150:VAL:HG21	2.01	0.41
1:A:30:VAL:CG2	1:A:90:VAL:HB	2.48	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	189/206 (92%)	186 (98%)	2 (1%)	1 (0%)	34	48
2	B	238/243 (98%)	226 (95%)	12 (5%)	0	100	100
3	C	217/233 (93%)	211 (97%)	6 (3%)	0	100	100
All	All	644/682 (94%)	623 (97%)	20 (3%)	1 (0%)	52	69

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	92	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/185 (86%)	148 (92%)	12 (8%)	17	26
2	B	199/213 (93%)	191 (96%)	8 (4%)	38	58
3	C	179/207 (86%)	167 (93%)	12 (7%)	20	31
All	All	538/605 (89%)	506 (94%)	32 (6%)	24	38

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	SER
1	A	30	VAL
1	A	32	ASN
1	A	46	ASN
1	A	61	SER
1	A	64	THR
1	A	107	ARG
1	A	134	VAL
1	A	143	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	150	LYS
1	A	167	MET
1	A	182	ASP
2	B	31	ASN
2	B	32	ARG
2	B	37	ARG
2	B	58	LYS
2	B	85	GLN
2	B	97	LEU
2	B	192	ARG
2	B	241	ARG
3	C	45	ASP
3	C	53	LEU
3	C	68	LEU
3	C	83	LYS
3	C	124	LYS
3	C	183	LEU
3	C	184	ILE
3	C	187	HIS
3	C	192	SER
3	C	193	THR
3	C	198	LEU
3	C	220	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	GLN
1	A	7	GLN
1	A	32	ASN
1	A	34	GLN
1	A	46	ASN
1	A	116	ASN
1	A	143	GLN
1	A	172	ASN
2	B	8	ASN
2	B	31	ASN
2	B	51	GLN
2	B	52	ASN
2	B	55	GLN
2	B	85	GLN
2	B	183	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	78	ASN
3	C	114	HIS
3	C	158	GLN
3	C	164	HIS
3	C	180	GLN
3	C	204	GLN
3	C	216	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	197/206 (95%)	-0.09	5 (2%) 61 60	22, 35, 60, 71	0
2	B	240/243 (98%)	-0.23	1 (0%) 93 93	22, 32, 53, 67	4 (1%)
3	C	221/233 (94%)	0.15	10 (4%) 37 38	26, 42, 62, 70	0
All	All	658/682 (96%)	-0.06	16 (2%) 62 61	22, 36, 61, 71	4 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	49	GLU	6.7
3	C	226	ILE	3.6
3	C	137	THR	3.1
3	C	127	ILE	2.9
3	C	191	GLY	2.5
1	A	181	SER	2.5
1	A	182	ASP	2.5
3	C	116	ASN	2.4
2	B	225	ASP	2.3
3	C	131	ILE	2.2
1	A	96	GLY	2.1
3	C	47	PHE	2.1
3	C	48	LEU	2.1
1	A	151	ASP	2.0
3	C	231	TYR	2.0
1	A	168	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	ZN	C	300	1/1	0.96	0.15	-	66,66,66,66	1

6.5 Other polymers [i](#)

There are no such residues in this entry.