



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:05 AM GMT

PDB ID : 2FLB  
Title : Discovery of a Novel Hydroxy Pyrazole Based Factor IXa Inhibitor  
Authors : Katz, B.A.  
Deposited on : 2006-01-05  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	152	<div> <div>6%</div> <div> <div>47%</div> <div>13%</div> <div>•</div> <div>37%</div> </div> </div>
2	H	254	<div> <div>5%</div> <div> <div>79%</div> <div>17%</div> <div>•</div> </div> </div>
3	T	218	<div> <div>10%</div> <div> <div>48%</div> <div>19%</div> <div>•</div> <div>31%</div> </div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4278 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coagulation factor VII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	96	Total	C	N	O	S	0	2	0
			727	439	126	149	13			


- Molecule 2 is a protein called Coagulation factor VII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	254	Total	C	N	O	S	0	3	0
			1988	1263	353	359	13			

- Molecule 3 is a protein called Tissue factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	T	151	Total	C	N	O	S	0	0	0
			1218	774	198	244	2			

- Molecule 4 is 2-(4-HYDROXY-5-PHENYL-1H-PYRAZOL-3-YL)-1H-BENZOIMIDAZOLE-5-CARBOXAMIDINE (three-letter code: 6NH) (formula: C<sub>17</sub>H<sub>14</sub>N<sub>6</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	H	1	Total	C	N	O	0	0
			24	17	6	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	161	Total	O	0	0
			161	161		
5	L	65	Total	O	0	0
			65	65		
5	T	95	Total	O	0	0
			95	95		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Coagulation factor VII



#### • Molecule 2: Coagulation factor VII



#### • Molecule 3: Tissue factor



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.45Å 69.03Å 78.76Å 90.00° 90.21° 90.00°	Depositor
Resolution (Å)	7.00 – 1.95 7.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	(Not available) (7.00-1.95) 93.3 (7.00-1.95)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.29 (at 1.95Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, $R_{free}$	0.247 , 0.280 0.236 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	20.3	Xtriage
Anisotropy	0.962	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 98.8	EDS
Estimated twinning fraction	0.000 for l,k,-h 0.024 for h,-k,-l 0.016 for l,-k,h	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 56089 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4278	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 6NH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	L	0.81	0/748	1.40	8/1009 (0.8%)
2	H	0.94	0/2046	1.62	45/2784 (1.6%)
3	T	0.81	0/1244	1.51	15/1693 (0.9%)
All	All	0.88	0/4038	1.55	68/5486 (1.2%)

There are no bond length outliers.

All (68) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	134	ARG	NE-CZ-NH2	-11.16	114.72	120.30
3	T	25	TRP	CD1-CG-CD2	9.18	113.64	106.30
1	L	110	ARG	NE-CZ-NH2	-8.99	115.81	120.30
2	H	237	TRP	CD1-CG-CD2	8.47	113.08	106.30
2	H	107	ARG	NE-CZ-NH1	8.44	124.52	120.30
2	H	51	TRP	CG-CD2-CE3	8.28	141.35	133.90
3	T	14	TRP	CD1-CG-CD2	8.25	112.90	106.30
2	H	134	ARG	CA-CB-CG	-8.09	95.60	113.40
2	H	61	TRP	CD1-CG-CD2	8.02	112.72	106.30
2	H	29	TRP	CD1-CG-CD2	7.98	112.68	106.30
2	H	51	TRP	CE2-CD2-CG	-7.95	100.94	107.30
1	L	118	TYR	CB-CG-CD2	-7.88	116.27	121.00
3	T	45	TRP	CD1-CG-CD2	7.84	112.58	106.30
2	H	51	TRP	CD1-CG-CD2	7.78	112.52	106.30
2	H	61	TRP	CE2-CD2-CG	-7.71	101.13	107.30
2	H	141	TRP	CD1-CG-CD2	7.71	112.47	106.30
2	H	29	TRP	CE2-CD2-CG	-7.68	101.16	107.30
2	H	237	TRP	CE2-CD2-CG	-7.65	101.18	107.30
3	T	135	ARG	NE-CZ-NH1	-7.50	116.55	120.30
2	H	230	ARG	NE-CZ-NH1	7.44	124.02	120.30
3	T	45	TRP	CE2-CD2-CG	-7.34	101.43	107.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	T	157	TYR	N-CA-C	-7.26	91.39	111.00
3	T	14	TRP	CE2-CD2-CG	-7.07	101.64	107.30
2	H	77	ASP	CA-C-N	6.99	130.19	116.20
2	H	215	TRP	CE2-CD2-CG	-6.95	101.74	107.30
2	H	253	ARG	NE-CZ-NH2	-6.95	116.83	120.30
2	H	141	TRP	CE2-CD2-CG	-6.87	101.81	107.30
2	H	51	TRP	CB-CG-CD1	-6.82	118.14	127.00
2	H	215	TRP	CD1-CG-CD2	6.72	111.68	106.30
2	H	207	TRP	CD1-CG-CD2	6.58	111.56	106.30
2	H	68	LEU	CA-CB-CG	-6.55	100.24	115.30
3	T	25	TRP	CE2-CD2-CG	-6.53	102.08	107.30
2	H	170(C)	ARG	NE-CZ-NH2	6.20	123.40	120.30
2	H	134	ARG	NE-CZ-NH1	6.19	123.39	120.30
3	T	25	TRP	CG-CD1-NE1	-6.15	103.95	110.10
2	H	61	TRP	CG-CD2-CE3	6.00	139.30	133.90
1	L	108	THR	CA-CB-CG2	5.97	120.75	112.40
2	H	215	TRP	CG-CD2-CE3	5.86	139.17	133.90
2	H	77	ASP	O-C-N	-5.85	113.25	123.20
1	L	113	ARG	NE-CZ-NH2	-5.83	117.39	120.30
1	L	101	TYR	CB-CG-CD2	-5.80	117.52	121.00
2	H	138[A]	VAL	CA-CB-CG2	-5.80	102.20	110.90
2	H	138[B]	VAL	CA-CB-CG2	-5.80	102.20	110.90
2	H	230	ARG	NE-CZ-NH2	-5.74	117.43	120.30
2	H	237	TRP	CB-CG-CD1	-5.74	119.54	127.00
3	T	34	TYR	CB-CG-CD1	-5.74	117.56	121.00
2	H	29	TRP	CG-CD1-NE1	-5.73	104.37	110.10
1	L	113	ARG	NE-CZ-NH1	5.70	123.15	120.30
2	H	243	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	L	92	VAL	CG1-CB-CG2	5.67	119.97	110.90
2	H	204	ARG	NE-CZ-NH1	5.61	123.10	120.30
2	H	184	TYR	CB-CG-CD2	-5.53	117.68	121.00
3	T	153	TYR	CB-CG-CD1	-5.52	117.69	121.00
1	L	92	VAL	N-CA-CB	-5.51	99.38	111.50
2	H	204	ARG	NE-CZ-NH2	-5.51	117.55	120.30
2	H	68	LEU	CD1-CG-CD2	5.47	126.91	110.50
3	T	83	VAL	CA-CB-CG2	-5.29	102.96	110.90
2	H	207	TRP	CE2-CD2-CG	-5.26	103.09	107.30
3	T	86	THR	N-CA-CB	-5.23	100.36	110.30
2	H	29	TRP	CG-CD2-CE3	5.22	138.60	133.90
2	H	79	ASP	N-CA-C	5.21	125.08	111.00
2	H	61	TRP	CB-CG-CD1	-5.21	120.22	127.00
3	T	83	VAL	N-CA-CB	-5.13	100.21	111.50

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	T	14	TRP	CG-CD1-NE1	-5.11	105.00	110.10
2	H	237	TRP	CG-CD1-NE1	-5.09	105.01	110.10
2	H	164	MET	CG-SD-CE	-5.07	92.08	100.20
2	H	129(B)	ARG	NE-CZ-NH1	5.03	122.81	120.30
2	H	215	TRP	CB-CG-CD1	-5.02	120.47	127.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	727	0	663	9	0
2	H	1988	0	1968	23	0
3	T	1218	0	1181	20	0
4	H	24	0	14	6	0
5	H	161	0	0	3	0
5	L	65	0	0	0	0
5	T	95	0	0	0	0
All	All	4278	0	3826	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:T:152:ILE:HG13	3:T:192:VAL:HG23	1.68	0.73
2:H:195:SER:OG	4:H:1246:6NH:O4'	2.08	0.71
3:T:86:THR:HG23	3:T:90:GLY:HA2	1.79	0.65
2:H:187:GLY:HA2	2:H:221(A):ALA:O	1.98	0.64
4:H:1246:6NH:N3	5:H:1321:HOH:O	2.29	0.60
1:L:121[A]:LEU:HD21	1:L:128:THR:HB	1.85	0.58
2:H:59:PHE:HA	2:H:60(B):ILE:HG12	1.87	0.56
4:H:1246:6NH:H2B	5:H:1354:HOH:O	2.07	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:99:GLU:HA	2:H:204:ARG:HD3	1.89	0.54
2:H:244:SER:HB3	2:H:247:ARG:HH12	1.72	0.54
3:T:154:THR:O	3:T:189:VAL:HA	2.08	0.53
2:H:236:GLU:CD	2:H:236:GLU:H	2.14	0.52
2:H:244:SER:HB3	2:H:247:ARG:NH1	2.25	0.51
2:H:101:HIS:HA	2:H:234:TYR:OH	2.11	0.50
2:H:84:ARG:H	2:H:84:ARG:HD2	1.77	0.49
1:L:115:HIS:O	1:L:118:TYR:HB2	2.13	0.49
2:H:68:LEU:HD13	2:H:112:VAL:HG11	1.93	0.49
2:H:236:GLU:HG3	5:H:1383:HOH:O	2.13	0.48
2:H:50:ILE:HG13	2:H:51:TRP:CD1	2.48	0.48
3:T:16:SER:HA	3:T:20:LYS:O	2.14	0.47
2:H:143:GLN:NE2	2:H:147:ARG:H	2.12	0.47
1:L:105:HIS:HB3	1:L:108:THR:HG22	1.96	0.47
2:H:35:VAL:HG21	2:H:60(B):ILE:HD12	1.97	0.47
3:T:192:VAL:HA	3:T:200:ARG:O	2.15	0.46
2:H:68:LEU:HD22	2:H:118:VAL:HG13	1.96	0.46
2:H:92:SER:OG	2:H:255:PRO:HA	2.16	0.46
3:T:109:GLY:O	3:T:202:SER:HB2	2.15	0.46
3:T:193:ILE:HB	3:T:196:ARG:HD2	1.97	0.46
1:L:112:CYS:HB2	1:L:124:GLY:O	2.16	0.46
3:T:132:THR:HG22	3:T:143:LEU:HD12	1.99	0.45
2:H:195:SER:OG	4:H:1246:6NH:N3	2.40	0.45
3:T:86:THR:CG2	3:T:90:GLY:HA2	2.44	0.45
2:H:213:VAL:HG12	4:H:1246:6NH:H3	2.00	0.44
2:H:195:SER:HG	4:H:1246:6NH:HN3	1.62	0.44
3:T:45:TRP:HH2	3:T:72:LEU:HD12	1.81	0.44
2:H:170(I):PRO:HD2	2:H:215:TRP:CZ3	2.52	0.44
1:L:71:PHE:CE2	3:T:131:ARG:HD3	2.53	0.44
3:T:136:ARG:HB3	3:T:141:LEU:HD21	1.99	0.44
2:H:48:ASN:HD21	2:H:242:MET:CE	2.31	0.44
3:T:79:PRO:HD2	3:T:85:SER:HB2	2.00	0.43
3:T:152:ILE:HD11	3:T:194:PRO:HG3	1.99	0.43
1:L:62:LYS:O	1:L:68:TYR:HA	2.19	0.43
3:T:36:VAL:HG22	3:T:75:VAL:HG22	2.00	0.42
3:T:48:LYS:HE3	3:T:48:LYS:HB3	1.94	0.42
3:T:130:GLU:O	3:T:142:SER:HB2	2.20	0.42
3:T:79:PRO:HD2	3:T:85:SER:CB	2.50	0.41
2:H:124:PRO:O	2:H:235:ILE:HD13	2.20	0.41
1:L:129:PRO:HB3	1:L:134:PRO:HG3	2.03	0.41
3:T:131:ARG:HA	3:T:142:SER:HA	2.02	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:T:64:VAL:HB	3:T:134:VAL:HA	2.02	0.40
1:L:105:HIS:HB3	1:L:108:THR:CG2	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	96/152 (63%)	85 (88%)	10 (10%)	1 (1%)	19	8
2	H	255/254 (100%)	239 (94%)	15 (6%)	1 (0%)	39	27
3	T	145/218 (66%)	133 (92%)	7 (5%)	5 (3%)	5	0
All	All	496/624 (80%)	457 (92%)	32 (6%)	7 (1%)	14	4

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	50	CYS
3	T	156	TYR
3	T	88	SER
3	T	200	ARG
2	H	170(D)	LYS
3	T	150	ASP
3	T	199	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	85/132 (64%)	77 (91%)	8 (9%)	11	2
2	H	219/216 (101%)	206 (94%)	13 (6%)	24	10
3	T	139/199 (70%)	131 (94%)	8 (6%)	25	10
All	All	443/547 (81%)	414 (94%)	29 (6%)	22	8

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	49	GLN
1	L	66	GLN
1	L	74	PRO
1	L	88	GLN
1	L	92	VAL
1	L	103	SER
1	L	110	ARG
1	L	128	THR
2	H	29	TRP
2	H	68	LEU
2	H	75	GLU
2	H	84	ARG
2	H	129(B)	ARG
2	H	159	ASN
2	H	170(C)	ARG
2	H	190[A]	SER
2	H	190[B]	SER
2	H	217	GLN
2	H	233	GLN
2	H	236	GLU
2	H	247	ARG
3	T	6	THR
3	T	27	PRO
3	T	79	PRO
3	T	83	VAL
3	T	86	THR
3	T	129	ASP
3	T	188	SER
3	T	195	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	80	ASN
2	H	48	ASN
2	H	100	ASN
2	H	143	GLN
2	H	159	ASN
2	H	217	GLN
2	H	233	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	6NH	H	1246	-	27,27,27	2.09	7 (25%)	20,39,39	2.23	7 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	6NH	H	1246	-	-	0/8/12/12	0/4/4/4

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	1246	6NH	C1B-C3'	-6.54	1.41	1.49
4	H	1246	6NH	C1-C7	-4.11	1.40	1.47
4	H	1246	6NH	C5'-C8	-3.82	1.42	1.49
4	H	1246	6NH	C4-N3	-2.19	1.31	1.39
4	H	1246	6NH	C6-C1	2.00	1.40	1.37
4	H	1246	6NH	C7-N2	2.05	1.35	1.28
4	H	1246	6NH	N1'-N2'	3.69	1.45	1.37

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	1246	6NH	C1B-C3'-C4'	-4.40	121.95	130.48
4	H	1246	6NH	C1-C6-C5	-4.37	116.27	120.80
4	H	1246	6NH	C2-C3-C4	-3.47	117.12	120.88
4	H	1246	6NH	C2-C1-C7	-2.35	117.70	120.67
4	H	1246	6NH	C6B-C1B-C2B	2.11	121.54	117.55
4	H	1246	6NH	C2-C1-C6	3.16	123.38	119.29
4	H	1246	6NH	C1B-C3'-N2'	3.77	125.57	120.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	1246	6NH	6	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	96/152 (63%)	0.32	9 (9%) 11 17	9, 26, 53, 59	17 (17%)
2	H	254/254 (100%)	0.01	13 (5%) 32 43	7, 19, 54, 70	22 (8%)
3	T	151/218 (69%)	0.82	22 (14%) 3 5	10, 31, 65, 87	25 (16%)
All	All	501/624 (80%)	0.31	44 (8%) 12 20	7, 23, 58, 87	64 (12%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	170(F)	GLY	18.1
2	H	170(E)	VAL	15.5
3	T	205	SER	13.6
3	T	137	ASN	11.0
3	T	188	SER	10.5
1	L	66	GLN	10.4
3	T	149	LYS	10.1
3	T	156	TYR	9.5
3	T	150	ASP	8.8
3	T	81	GLY	7.9
2	H	170(G)	ASP	7.3
1	L	53	SER	5.6
2	H	60(C)	LYS	5.1
3	T	157	TYR	5.1
3	T	151	LEU	4.6
3	T	88	SER	4.4
3	T	129	ASP	4.2
3	T	155	LEU	4.1
1	L	49	GLN	3.9
1	L	52	SER	3.9
2	H	247	ARG	3.7
2	H	62	ARG	3.7
2	H	170(H)	SER	3.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	L	48	ASP	3.5
3	T	148	GLY	3.5
3	T	152	ILE	3.4
3	T	87	GLY	3.3
1	L	50	CYS	3.3
3	T	198	VAL	3.2
3	T	203	THR	2.9
1	L	67	SER	2.9
2	H	170(D)	LYS	2.7
3	T	138	ASN	2.7
2	H	245	GLU	2.7
2	H	60(B)	ILE	2.6
2	H	248	PRO	2.5
3	T	153	TYR	2.3
1	L	51	ALA	2.3
2	H	246	PRO	2.2
3	T	109	GLY	2.2
3	T	197	THR	2.2
2	H	244	SER	2.1
3	T	199	ASN	2.0
1	L	47	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	6NH	H	1246	24/24	0.92	0.11	0.62	16,21,27,31	0



## 6.5 Other polymers [i](#)

There are no such residues in this entry.