



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:06 AM GMT

PDB ID : 2FMY
Title : CO-dependent transcription factor CooA from Carboxydotherrmus hydrogenoformans (Imidazole-bound form)
Authors : Higuchi, Y.; Komori, H.
Deposited on : 2006-01-10
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	220	<div> <div>5%</div> <div>93%</div> <div>6%</div> </div>
1	B	220	<div> <div>12%</div> <div>88%</div> <div>11%</div> </div>
1	C	220	<div> <div>5%</div> <div>90%</div> <div>9%</div> </div>
1	D	220	<div> <div>15%</div> <div>94%</div> <div>5%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7446 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called carbon monoxide oxidation system transcription regulator CooA-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	218	Total	C	N	O	S	0	0	0
			1750	1122	295	326	7			
1	B	218	Total	C	N	O	S	0	0	0
			1750	1122	295	326	7			
1	C	218	Total	C	N	O	S	0	0	0
			1750	1122	295	326	7			
1	D	218	Total	C	N	O	S	0	0	0
			1750	1122	295	326	7			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	-	CLONING ARTIFACT	GB 77996136
A	3	THR	-	CLONING ARTIFACT	GB 77996136
A	4	GLN	-	CLONING ARTIFACT	GB 77996136
B	1002	ALA	-	CLONING ARTIFACT	GB 77996136
B	1003	THR	-	CLONING ARTIFACT	GB 77996136
B	1004	GLN	-	CLONING ARTIFACT	GB 77996136
C	2002	ALA	-	CLONING ARTIFACT	GB 77996136
C	2003	THR	-	CLONING ARTIFACT	GB 77996136
C	2004	GLN	-	CLONING ARTIFACT	GB 77996136
D	3002	ALA	-	CLONING ARTIFACT	GB 77996136
D	3003	THR	-	CLONING ARTIFACT	GB 77996136
D	3004	GLN	-	CLONING ARTIFACT	GB 77996136

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is IMIDAZOLE (three-letter code: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N 5 3 2	0	0
3	B	1	Total C N 5 3 2	0	0
3	C	1	Total C N 5 3 2	0	0
3	D	1	Total C N 5 3 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	87	Total O 87 87	0	0
4	B	57	Total O 57 57	0	0
4	C	66	Total O 66 66	0	0
4	D	44	Total O 44 44	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: carbon monoxide oxidation system transcription regulator CoxA-1




- Molecule 1: carbon monoxide oxidation system transcription regulator CoxA-1



- Molecule 1: carbon monoxide oxidation system transcription regulator CoxA-1



- Molecule 1: carbon monoxide oxidation system transcription regulator CoxA-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.65Å 93.93Å 91.90Å 90.00° 104.73° 90.00°	Depositor
Resolution (Å)	32.33 – 2.20 32.32 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (32.33-2.20) 99.4 (32.32-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.07 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.207 , 0.247 0.200 , 0.241	Depositor DCC
R_{free} test set	2563 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	35.4	Xtriage
Anisotropy	0.955	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	2 of 51311 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7446	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.30 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 8.1831e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/1771	0.57	0/2387
1	B	0.46	1/1771 (0.1%)	0.56	0/2387
1	C	0.51	1/1771 (0.1%)	0.58	0/2387
1	D	0.42	0/1771	0.55	0/2387
All	All	0.46	2/7084 (0.0%)	0.57	0/9548

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	2091	GLU	CD-OE2	8.87	1.35	1.25
1	B	1035	LYS	CE-NZ	5.67	1.63	1.49

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1750	0	1825	9	0
1	B	1750	0	1825	17	0
1	C	1750	0	1825	13	0
1	D	1750	0	1825	4	0
2	A	43	0	30	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	43	0	30	2	0
2	C	43	0	30	1	0
2	D	43	0	30	4	0
3	A	5	0	4	2	0
3	B	5	0	4	0	0
3	C	5	0	4	0	0
3	D	5	0	4	2	0
4	A	87	0	0	0	0
4	B	57	0	0	0	0
4	C	66	0	0	1	0
4	D	44	0	0	0	0
All	All	7446	0	7436	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:2181:THR:HG22	1:C:2183:GLN:H	1.46	0.80
1:B:1143:ARG:HH21	1:B:1180:THR:HG23	1.54	0.71
1:C:2125:LEU:HD23	2:D:3300:HEM:CMD	2.24	0.67
1:D:3048:LEU:HD13	1:D:3097:TYR:HB2	1.78	0.65
1:A:117:MET:HE1	1:B:1118:VAL:HG11	1.83	0.61
1:C:2005:MET:HB2	3:D:3301:IMD:C2	2.33	0.59
4:C:4067:HOH:O	1:D:3194:LYS:HE3	2.02	0.59
1:A:158:LYS:HE3	1:A:159:VAL:O	2.06	0.55
1:B:1143:ARG:HH21	1:B:1180:THR:CG2	2.20	0.54
1:C:2125:LEU:HD23	2:D:3300:HEM:HMD2	1.90	0.54
1:B:1161:GLN:HB3	1:B:1207:LEU:HD21	1.91	0.53
1:A:121:LEU:HA	1:A:124:LEU:HD12	1.90	0.53
1:A:72:ILE:HD12	1:A:90:MET:SD	2.50	0.52
1:C:2077:ASP:OD1	1:C:2119:LYS:NZ	2.44	0.50
1:B:1171:THR:HG22	1:B:1182:ARG:HD2	1.94	0.50
1:A:104:GLN:HA	2:A:300:HEM:HBB1	1.93	0.50
1:C:2152:ALA:HA	1:C:2165:LEU:CD2	2.41	0.50
1:C:2051:LEU:HB2	1:C:2078:ILE:HD13	1.94	0.49
1:C:2181:THR:HG22	1:C:2183:GLN:N	2.22	0.48
3:A:301:IMD:H5	1:B:1118:VAL:O	2.14	0.48
1:B:1200:ARG:HD3	1:B:1202:ASN:O	2.13	0.48
1:B:1007:LEU:HD11	1:B:1118:VAL:HG11	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1142:LEU:O	1:B:1146:GLU:HG3	2.15	0.47
1:A:7:LEU:HD22	2:B:1300:HEM:HBB2	1.96	0.47
1:C:2152:ALA:HA	1:C:2165:LEU:HD21	1.96	0.47
1:B:1143:ARG:NH2	1:B:1180:THR:CG2	2.78	0.47
1:C:2125:LEU:HD23	2:D:3300:HEM:HMD1	1.97	0.46
1:B:1191:ASP:O	1:B:1195:MET:HG3	2.17	0.45
1:C:2197:ILE:HG21	1:C:2214:LEU:HD13	1.99	0.44
1:B:1140:ALA:CB	1:B:1180:THR:HG21	2.48	0.44
1:B:1140:ALA:HB2	1:B:1180:THR:HG21	2.01	0.43
1:A:125:LEU:HD23	2:B:1300:HEM:HMD2	2.01	0.43
1:A:66:LYS:CG	1:A:67:GLU:H	2.32	0.42
1:D:3199:GLU:HG3	1:D:3209:LYS:HD2	2.00	0.42
3:A:301:IMD:N1	1:B:1005:MET:HG3	2.35	0.42
1:C:2100:ILE:HD11	2:C:2300:HEM:C2B	2.55	0.42
1:B:1195:MET:CE	1:B:1217:PHE:CZ	3.03	0.42
1:B:1159:VAL:HB	1:B:1160:PRO:HD2	2.03	0.41
1:D:3035:LYS:HG2	1:D:3036:LYS:HG2	2.02	0.41
2:D:3300:HEM:CHD	3:D:3301:IMD:H4	2.51	0.41
1:C:2142:LEU:HD11	1:C:2219:SER:HB2	2.02	0.41
1:A:165:LEU:HD22	1:A:208:LEU:HD13	2.03	0.41
1:B:1056:ARG:HG2	1:B:1074:GLU:OE1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	216/220 (98%)	211 (98%)	5 (2%)	0	100	100
1	B	216/220 (98%)	212 (98%)	4 (2%)	0	100	100
1	C	216/220 (98%)	212 (98%)	4 (2%)	0	100	100
1	D	216/220 (98%)	211 (98%)	5 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	864/880 (98%)	846 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/196 (100%)	195 (100%)	0	100	100
1	B	195/196 (100%)	193 (99%)	2 (1%)	82	91
1	C	195/196 (100%)	192 (98%)	3 (2%)	72	84
1	D	195/196 (100%)	191 (98%)	4 (2%)	61	74
All	All	780/784 (100%)	771 (99%)	9 (1%)	78	88

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1009	ASP
1	B	1124	LEU
1	C	2008	THR
1	C	2124	LEU
1	C	2218	SER
1	D	3024	VAL
1	D	3044	THR
1	D	3155	THR
1	D	3200	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	1104	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	HEM	A	300	1,3	30,50,50	2.24	9 (30%)	24,82,82	2.59	12 (50%)
3	IMD	A	301	2	3,5,5	0.45	0	4,5,5	0.58	0
2	HEM	B	1300	1,3	30,50,50	2.08	8 (26%)	24,82,82	2.60	11 (45%)
3	IMD	B	1301	2	3,5,5	0.54	0	4,5,5	0.40	0
2	HEM	C	2300	1,3	30,50,50	2.21	11 (36%)	24,82,82	2.63	10 (41%)
3	IMD	C	2301	2	3,5,5	0.49	0	4,5,5	0.28	0
2	HEM	D	3300	1,3	30,50,50	2.20	10 (33%)	24,82,82	2.65	13 (54%)
3	IMD	D	3301	2	3,5,5	0.45	0	4,5,5	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	300	1,3	-	0/10/54/54	0/0/8/8

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IMD	A	301	2	-	0/0/0/0	0/1/1/1
2	HEM	B	1300	1,3	-	0/10/54/54	0/0/8/8
3	IMD	B	1301	2	-	0/0/0/0	0/1/1/1
2	HEM	C	2300	1,3	-	0/10/54/54	0/0/8/8
3	IMD	C	2301	2	-	0/0/0/0	0/1/1/1
2	HEM	D	3300	1,3	-	0/10/54/54	0/0/8/8
3	IMD	D	3301	2	-	0/0/0/0	0/1/1/1

All (38) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3300	HEM	C3B-C4B	-7.12	1.45	1.51
2	A	300	HEM	C3B-C4B	-6.73	1.45	1.51
2	C	2300	HEM	C3B-C4B	-6.58	1.46	1.51
2	B	1300	HEM	C3B-C4B	-6.33	1.46	1.51
2	C	2300	HEM	C3D-C4D	-5.64	1.44	1.51
2	A	300	HEM	C3D-C4D	-5.59	1.44	1.51
2	D	3300	HEM	C3D-C4D	-5.56	1.44	1.51
2	B	1300	HEM	C3D-C4D	-5.20	1.44	1.51
2	C	2300	HEM	C2C-C1C	-4.19	1.44	1.52
2	B	1300	HEM	C2C-C1C	-3.96	1.45	1.52
2	A	300	HEM	C2C-C1C	-3.58	1.45	1.52
2	D	3300	HEM	C2C-C1C	-3.54	1.45	1.52
2	D	3300	HEM	C2D-C1D	-2.20	1.44	1.51
2	C	2300	HEM	C2D-C1D	-2.14	1.44	1.51
2	B	1300	HEM	C2D-C1D	-2.09	1.45	1.51
2	A	300	HEM	C2B-C1B	-2.07	1.45	1.51
2	B	1300	HEM	C2B-C1B	-2.01	1.45	1.51
2	D	3300	HEM	C2B-C1B	-2.01	1.45	1.51
2	B	1300	HEM	C1C-NC	2.01	1.38	1.36
2	C	2300	HEM	C4C-NC	2.05	1.38	1.36
2	D	3300	HEM	CAA-C2A	2.06	1.55	1.52
2	D	3300	HEM	FE-NC	2.08	2.04	1.95
2	B	1300	HEM	C3B-CAB	2.08	1.55	1.51
2	C	2300	HEM	C1C-NC	2.09	1.38	1.36
2	D	3300	HEM	C3C-CAC	2.10	1.55	1.51
2	A	300	HEM	C3C-CAC	2.11	1.55	1.51
2	C	2300	HEM	CAA-C2A	2.11	1.55	1.52
2	C	2300	HEM	C3B-CAB	2.13	1.55	1.51
2	B	1300	HEM	C3C-CAC	2.21	1.55	1.51
2	A	300	HEM	CMA-C3A	2.26	1.56	1.51
2	D	3300	HEM	C4C-NC	2.28	1.38	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2300	HEM	CMA-C3A	2.28	1.56	1.51
2	C	2300	HEM	FE-ND	2.34	2.09	1.97
2	A	300	HEM	C4C-NC	2.37	1.38	1.36
2	C	2300	HEM	FE-NC	2.56	2.05	1.95
2	D	3300	HEM	FE-ND	2.59	2.11	1.97
2	A	300	HEM	FE-NB	2.97	2.13	1.97
2	A	300	HEM	FE-NC	3.31	2.08	1.95

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2300	HEM	CBA-CAA-C2A	-4.45	104.56	112.53
2	C	2300	HEM	CBD-CAD-C3D	-4.34	100.92	113.55
2	D	3300	HEM	CBD-CAD-C3D	-4.25	101.18	113.55
2	A	300	HEM	CBA-CAA-C2A	-4.11	105.17	112.53
2	D	3300	HEM	C3B-CAB-CBB	-3.92	118.45	124.46
2	B	1300	HEM	CBA-CAA-C2A	-3.72	105.87	112.53
2	B	1300	HEM	CBD-CAD-C3D	-3.21	104.20	113.55
2	A	300	HEM	CBD-CAD-C3D	-3.19	104.28	113.55
2	C	2300	HEM	C3C-CAC-CBC	-3.06	119.77	124.46
2	B	1300	HEM	C3C-CAC-CBC	-3.01	119.84	124.46
2	A	300	HEM	C3C-CAC-CBC	-2.88	120.04	124.46
2	D	3300	HEM	CBA-CAA-C2A	-2.58	107.91	112.53
2	D	3300	HEM	C3C-CAC-CBC	-2.40	120.77	124.46
2	B	1300	HEM	CMA-C3A-C4A	-2.37	124.44	128.36
2	D	3300	HEM	C3B-C4B-NB	-2.29	107.25	111.63
2	A	300	HEM	C3B-C4B-NB	-2.22	107.38	111.63
2	C	2300	HEM	CAA-CBA-CGA	2.25	116.87	112.75
2	D	3300	HEM	C2C-C1C-CHC	2.33	127.22	123.68
2	B	1300	HEM	C2D-C3D-C4D	2.43	105.61	101.50
2	B	1300	HEM	C3B-C4B-CHC	2.47	126.64	123.16
2	D	3300	HEM	C2D-C3D-C4D	2.53	105.79	101.50
2	A	300	HEM	C2D-C3D-C4D	2.53	105.79	101.50
2	A	300	HEM	C2C-C1C-CHC	2.70	127.78	123.68
2	A	300	HEM	C3B-C4B-CHC	2.71	126.98	123.16
2	A	300	HEM	CMD-C2D-C3D	2.84	126.91	114.35
2	D	3300	HEM	C3B-C4B-CHC	2.94	127.30	123.16
2	C	2300	HEM	CMD-C2D-C3D	2.95	127.39	114.35
2	C	2300	HEM	C2D-C3D-C4D	2.99	106.57	101.50
2	D	3300	HEM	CMD-C2D-C3D	3.04	127.78	114.35
2	B	1300	HEM	CMD-C2D-C3D	3.12	128.13	114.35
2	C	2300	HEM	CAD-C3D-C4D	3.97	126.49	112.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	HEM	CMC-C2C-C3C	4.02	126.56	116.53
2	D	3300	HEM	CAD-C3D-C4D	4.06	126.79	112.47
2	D	3300	HEM	CMB-C2B-C3B	4.15	126.89	116.53
2	C	2300	HEM	CMB-C2B-C3B	4.16	126.92	116.53
2	D	3300	HEM	CMC-C2C-C3C	4.23	127.09	116.53
2	B	1300	HEM	CAD-C3D-C4D	4.26	127.51	112.47
2	A	300	HEM	CAD-C3D-C4D	4.27	127.55	112.47
2	B	1300	HEM	CMC-C2C-C3C	4.43	127.59	116.53
2	C	2300	HEM	CMC-C2C-C3C	4.61	128.04	116.53
2	B	1300	HEM	CMB-C2B-C3B	4.64	128.12	116.53
2	A	300	HEM	CAD-C3D-C2D	4.67	126.64	113.22
2	B	1300	HEM	CAD-C3D-C2D	4.74	126.85	113.22
2	C	2300	HEM	CAD-C3D-C2D	4.75	126.88	113.22
2	A	300	HEM	CMB-C2B-C3B	4.80	128.51	116.53
2	D	3300	HEM	CAD-C3D-C2D	4.92	127.38	113.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	300	HEM	1	0
3	A	301	IMD	2	0
2	B	1300	HEM	2	0
2	C	2300	HEM	1	0
2	D	3300	HEM	4	0
3	D	3301	IMD	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	218/220 (99%)	0.22	10 (4%) 36 35	35, 40, 51, 59	0
1	B	218/220 (99%)	0.53	26 (11%) 6 6	30, 41, 51, 62	0
1	C	218/220 (99%)	0.37	11 (5%) 32 32	35, 41, 50, 70	0
1	D	218/220 (99%)	0.63	33 (15%) 3 3	32, 41, 53, 57	0
All	All	872/880 (99%)	0.44	80 (9%) 11 10	30, 41, 52, 70	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	2002	ALA	10.1
1	B	1002	ALA	8.4
1	D	3019	GLU	6.6
1	D	3002	ALA	6.6
1	D	3026	LYS	6.3
1	C	2003	THR	5.8
1	C	2219	SER	5.4
1	D	3016	LEU	4.6
1	D	3018	SER	4.5
1	D	3022	SER	4.5
1	B	1020	GLU	4.3
1	B	1021	TYR	4.1
1	B	1019	GLU	4.0
1	D	3024	VAL	4.0
1	D	3023	GLY	3.9
1	D	3065	ASP	3.9
1	A	219	SER	3.8
1	B	1018	SER	3.7
1	D	3219	SER	3.6
1	A	2	ALA	3.5
1	D	3017	ASN	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	3020	GLU	3.5
1	D	3021	TYR	3.3
1	B	1022	SER	3.3
1	B	1017	ASN	3.2
1	D	3162	GLY	3.2
1	D	3045	GLU	3.1
1	B	1023	GLY	3.1
1	D	3064	GLU	3.1
1	A	65	ASP	3.1
1	C	2121	LEU	3.0
1	B	1065	ASP	3.0
1	D	3099	ASP	2.9
1	D	3101	ARG	2.9
1	D	3218	SER	2.8
1	D	3118	VAL	2.8
1	A	118	VAL	2.8
1	B	1024	VAL	2.8
1	D	3029	ARG	2.7
1	D	3121	LEU	2.7
1	A	121	LEU	2.7
1	B	1122	GLY	2.7
1	C	2118	VAL	2.7
1	B	1026	LYS	2.6
1	D	3046	ARG	2.6
1	B	1120	VAL	2.6
1	D	3015	VAL	2.6
1	B	1007	LEU	2.6
1	B	1025	LEU	2.5
1	B	1028	PHE	2.5
1	A	176	LEU	2.5
1	C	2125	LEU	2.4
1	D	3013	LEU	2.4
1	D	3014	GLU	2.4
1	B	1003	THR	2.4
1	B	1158	LYS	2.4
1	B	1160	PRO	2.4
1	D	3160	PRO	2.4
1	D	3102	ASN	2.3
1	B	1006	ARG	2.3
1	D	3043	ASN	2.2
1	C	2117	MET	2.2
1	A	49	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1027	GLU	2.2
1	A	117	MET	2.2
1	A	142	LEU	2.2
1	C	2005	MET	2.2
1	C	2064	GLU	2.2
1	D	3125	LEU	2.1
1	B	1124	LEU	2.1
1	B	1161	GLN	2.1
1	C	2004	GLN	2.1
1	D	3122	GLY	2.1
1	B	1016	LEU	2.1
1	B	1106	ILE	2.1
1	D	3110	PHE	2.1
1	D	3007	LEU	2.1
1	A	106	ILE	2.1
1	B	1217	PHE	2.0
1	C	2045	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	HEM	D	3300	43/43	0.95	0.13	-1.05	40,45,51,55	0
2	HEM	B	1300	43/43	0.97	0.10	-1.55	33,35,42,47	0
3	IMD	B	1301	5/5	0.98	0.07	-2.11	31,32,33,33	0
3	IMD	D	3301	5/5	0.98	0.07	-2.19	42,42,43,43	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	IMD	C	2301	5/5	0.98	0.10	-2.24	31,32,32,32	0
2	HEM	C	2300	43/43	0.97	0.11	-2.37	29,32,39,44	0
2	HEM	A	300	43/43	0.96	0.11	-2.38	26,34,45,51	0
3	IMD	A	301	5/5	0.98	0.12	-2.58	33,34,34,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.