



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:41 AM GMT

PDB ID : 3FNM  
Title : Crystal structure of acivicin-inhibited gamma-glutamyltranspeptidase reveals critical roles for its C-terminus in autoprocessing and catalysis  
Authors : Williams, K.; Cullati, S.; Sand, A.; Biterova, E.I.; Barycki, J.J.  
Deposited on : 2008-12-25  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	<div> <div>5%</div> <div>85% 6% 9%</div> </div>
1	C	377	<div> <div>2%</div> <div>86% 6% 8%</div> </div>
2	B	188	<div> <div>4%</div> <div>90% 9%</div> </div>
2	D	188	<div> <div>4%</div> <div>90% 9%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8684 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-glutamyltranspeptidase (Ggt) Large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	342	Total	C	N	O	S	0	1	0
			2600	1657	445	489	9			
1	C	348	Total	C	N	O	S	0	2	0
			2654	1686	456	503	9			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	MET	-	EXPRESSION TAG	UNP O25743
A	4	GLY	-	EXPRESSION TAG	UNP O25743
A	5	SER	-	EXPRESSION TAG	UNP O25743
A	6	SER	-	EXPRESSION TAG	UNP O25743
A	7	HIS	-	EXPRESSION TAG	UNP O25743
A	8	HIS	-	EXPRESSION TAG	UNP O25743
A	9	HIS	-	EXPRESSION TAG	UNP O25743
A	10	HIS	-	EXPRESSION TAG	UNP O25743
A	11	HIS	-	EXPRESSION TAG	UNP O25743
A	12	HIS	-	EXPRESSION TAG	UNP O25743
A	13	SER	-	EXPRESSION TAG	UNP O25743
A	14	SER	-	EXPRESSION TAG	UNP O25743
A	15	GLY	-	EXPRESSION TAG	UNP O25743
A	16	LEU	-	EXPRESSION TAG	UNP O25743
A	17	VAL	-	EXPRESSION TAG	UNP O25743
A	18	PRO	-	EXPRESSION TAG	UNP O25743
A	19	ARG	-	EXPRESSION TAG	UNP O25743
A	20	GLY	-	EXPRESSION TAG	UNP O25743
A	21	SER	-	EXPRESSION TAG	UNP O25743
A	22	HIS	-	EXPRESSION TAG	UNP O25743
A	23	MET	-	EXPRESSION TAG	UNP O25743
A	24	ALA	-	EXPRESSION TAG	UNP O25743
C	3	MET	-	EXPRESSION TAG	UNP O25743
C	4	GLY	-	EXPRESSION TAG	UNP O25743
C	5	SER	-	EXPRESSION TAG	UNP O25743

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	6	SER	-	EXPRESSION TAG	UNP O25743
C	7	HIS	-	EXPRESSION TAG	UNP O25743
C	8	HIS	-	EXPRESSION TAG	UNP O25743
C	9	HIS	-	EXPRESSION TAG	UNP O25743
C	10	HIS	-	EXPRESSION TAG	UNP O25743
C	11	HIS	-	EXPRESSION TAG	UNP O25743
C	12	HIS	-	EXPRESSION TAG	UNP O25743
C	13	SER	-	EXPRESSION TAG	UNP O25743
C	14	SER	-	EXPRESSION TAG	UNP O25743
C	15	GLY	-	EXPRESSION TAG	UNP O25743
C	16	LEU	-	EXPRESSION TAG	UNP O25743
C	17	VAL	-	EXPRESSION TAG	UNP O25743
C	18	PRO	-	EXPRESSION TAG	UNP O25743
C	19	ARG	-	EXPRESSION TAG	UNP O25743
C	20	GLY	-	EXPRESSION TAG	UNP O25743
C	21	SER	-	EXPRESSION TAG	UNP O25743
C	22	HIS	-	EXPRESSION TAG	UNP O25743
C	23	MET	-	EXPRESSION TAG	UNP O25743
C	24	ALA	-	EXPRESSION TAG	UNP O25743

- Molecule 2 is a protein called Gamma-glutamyltranspeptidase (Ggt) Small subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	188	Total	C	N	O	S	0	2	0
			1440	910	243	280	7			
2	D	188	Total	C	N	O	S	0	1	0
			1436	909	242	278	7			

- Molecule 3 is (2S)-AMINO[(5S)-3-CHLORO-4,5-DIHYDROISOXAZOL-5-YL]ACETIC ACID (three-letter code: AVN) (formula: C<sub>5</sub>H<sub>7</sub>ClN<sub>2</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			10	5	2	3		
3	D	1	Total	C	N	O	0	0
			10	5	2	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	131	Total	O	0	0
			131	131		
4	B	90	Total	O	0	0
			90	90		
4	C	191	Total	O	0	0
			191	191		
4	D	122	Total	O	0	0
			122	122		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Gamma-glutamyltranspeptidase (Ggt) Large subunit




- Molecule 1: Gamma-glutamyltranspeptidase (Ggt) Large subunit



- Molecule 2: Gamma-glutamyltranspeptidase (Ggt) Small subunit



- Molecule 2: Gamma-glutamyltranspeptidase (Ggt) Small subunit



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.71Å 105.38Å 91.93Å 90.00° 91.69° 90.00°	Depositor
Resolution (Å)	29.20 – 1.70 26.42 – 1.70	Depositor EDS
% Data completeness (in resolution range)	96.4 (29.20-1.70) 96.4 (26.42-1.70)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.88 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.178 , 0.211 0.197 , 0.223	Depositor DCC
$R_{free}$ test set	10906 reflections (9.90%)	DCC
Wilson B-factor (Å <sup>2</sup> )	26.7	Xtriage
Anisotropy	0.253	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 45.3	EDS
Estimated twinning fraction	0.022 for h,-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 110128 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8684	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AVN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	0/2646	0.66	0/3562
1	C	0.72	2/2700 (0.1%)	0.72	0/3633
2	B	0.64	0/1470	0.74	1/1999 (0.1%)
2	D	0.70	0/1466	0.77	2/1992 (0.1%)
All	All	0.67	2/8282 (0.0%)	0.71	3/11186 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	118	LYS	C-O	10.31	1.43	1.23
1	C	119	GLN	CD-NE2	5.06	1.45	1.32

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	448	ARG	NE-CZ-NH1	6.18	123.39	120.30
2	B	502	ARG	NE-CZ-NH2	-5.45	117.57	120.30
2	D	448	ARG	NE-CZ-NH2	-5.21	117.69	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2600	0	2661	12	0
1	C	2654	0	2703	20	0
2	B	1440	0	1415	11	0
2	D	1436	0	1421	10	0
3	B	10	0	6	2	0
3	D	10	0	6	0	0
4	A	131	0	0	0	0
4	B	90	0	0	0	0
4	C	191	0	0	2	0
4	D	122	0	0	0	0
All	All	8684	0	8212	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:275:LYS:HE2	1:A:277:ILE:HD11	1.53	0.91
1:C:49:GLU:HG3	1:C:53:LYS:HE2	1.57	0.86
1:C:49:GLU:HG3	1:C:53:LYS:CE	2.19	0.73
2:B:380:THR:HB	2:B:471:PRO:CB	2.24	0.68
2:B:380:THR:HB	2:B:471:PRO:HB3	1.76	0.67
2:B:380:THR:HG21	3:B:1:AVN:N2	2.14	0.63
2:D:519:MET:HG2	2:D:524:LYS:HG3	1.80	0.62
2:B:380:THR:HG23	2:B:398:THR:HB	1.83	0.60
2:D:380:THR:N	2:D:398:THR:HG1	2.00	0.59
1:A:351:LYS:HD3	1:A:371:MET:HE3	1.85	0.58
1:C:49:GLU:HG3	1:C:53:LYS:NZ	2.19	0.58
1:A:53:LYS:O	1:A:57:GLU:HG3	2.04	0.57
1:C:164:LYS:HG3	4:C:455:HOH:O	2.07	0.54
1:C:53:LYS:N	1:C:53:LYS:HD3	2.23	0.54
2:B:476:ILE:HD11	3:B:1:AVN:N2	2.24	0.53
1:C:289:LEU:HD12	2:D:481:LEU:HB3	1.91	0.52
1:C:40:ALA:HB1	1:C:68:ILE:HD11	1.91	0.52
1:C:351:LYS:O	1:C:355:THR:HG23	2.09	0.52
2:D:380:THR:HG23	2:D:380:THR:O	2.10	0.51
1:C:75:VAL:HA	1:C:170:TYR:CZ	2.46	0.50
1:A:352:ILE:HG13	1:A:371:MET:HE2	1.93	0.50
1:C:289:LEU:CD1	2:D:481:LEU:HB3	2.43	0.49
2:D:461:ASN:O	2:D:462:ASN:HB2	2.13	0.48
1:C:79:ALA:HB1	2:D:380:THR:N	2.28	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:49:GLU:CG	1:C:53:LYS:HE2	2.37	0.48
2:D:551:LYS:HE2	2:D:553:LYS:O	2.14	0.48
1:A:289:LEU:HD12	2:B:481:LEU:HB3	1.97	0.46
2:B:470:SER:HB2	2:B:543:VAL:HG22	1.98	0.46
2:D:519:MET:HG2	2:D:524:LYS:CG	2.47	0.45
1:A:200:LYS:O	1:A:203:LEU:HG	2.17	0.45
1:A:40:ALA:HB1	1:A:68:ILE:HD11	1.97	0.44
1:C:112:LYS:HE2	1:C:112:LYS:HB3	1.73	0.44
2:B:519:MET:SD	2:B:524:LYS:HG2	2.58	0.44
2:D:470:SER:HB2	2:D:543:VAL:HG22	2.00	0.43
2:B:399:ILE:HB	2:B:417:ASN:HA	1.99	0.43
1:C:167[A]:GLU:HG3	1:C:217:ALA:HB1	1.99	0.43
1:C:188:LEU:O	1:C:194:LYS:HE2	2.19	0.42
1:A:75:VAL:HA	1:A:170:TYR:CZ	2.54	0.42
2:B:382:HIS:CE1	2:B:469:GLY:HA3	2.55	0.42
1:C:75:VAL:HG11	1:C:165:LEU:HD13	2.02	0.42
1:A:164:LYS:HG3	1:A:168:ASN:ND2	2.35	0.42
2:B:519:MET:HG2	2:B:524:LYS:HG3	2.02	0.42
1:C:310:LYS:NZ	4:C:440:HOH:O	2.49	0.42
1:C:75:VAL:CG1	1:C:165:LEU:HD13	2.50	0.41
1:A:351:LYS:HB3	1:A:371:MET:HE1	2.02	0.41
1:A:262:LYS:HD3	1:A:264:ARG:NH2	2.35	0.41
1:C:49:GLU:HG3	1:C:53:LYS:HZ1	1.85	0.41
1:C:108:LEU:HD11	1:C:254:GLU:HG2	2.02	0.40
1:A:178:GLU:O	1:A:182:GLU:HG3	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	341/377 (90%)	335 (98%)	6 (2%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	348/377 (92%)	343 (99%)	5 (1%)	0	100	100
2	B	188/188 (100%)	177 (94%)	10 (5%)	1 (0%)	34	15
2	D	187/188 (100%)	179 (96%)	7 (4%)	1 (0%)	34	15
All	All	1064/1130 (94%)	1034 (97%)	28 (3%)	2 (0%)	52	32

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	400	ASN
2	D	400	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	270/297 (91%)	269 (100%)	1 (0%)	93	90
1	C	276/297 (93%)	273 (99%)	3 (1%)	80	69
2	B	159/159 (100%)	155 (98%)	4 (2%)	55	34
2	D	159/159 (100%)	153 (96%)	6 (4%)	40	17
All	All	864/912 (95%)	850 (98%)	14 (2%)	70	54

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	LYS
2	B	380	THR
2	B	397	TYR
2	B	475	ARG
2	B	522	ASP
1	C	53	LYS
1	C	119	GLN
1	C	176	GLN
2	D	380	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	D	397	TYR
2	D	475	ARG
2	D	522	ASP
2	D	533	GLN
2	D	552	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	168	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	AVN	B	1	2	5,10,11	3.94	3 (60%)	2,13,15	3.89	1 (50%)
3	AVN	D	1	2	5,10,11	1.25	1 (20%)	2,13,15	2.81	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AVN	B	1	2	-	0/3/15/17	0/1/1/1
3	AVN	D	1	2	-	0/3/15/17	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1	AVN	O3-C3	-6.74	1.37	1.46
3	B	1	AVN	O3-N2	-4.43	1.35	1.42
3	B	1	AVN	C3-C2	-2.95	1.49	1.53
3	D	1	AVN	O3-C3	-2.23	1.43	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1	AVN	O3-C3-C5	-5.48	100.84	104.49
3	D	1	AVN	O3-C3-C5	-3.94	101.86	104.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1	AVN	2	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	342/377 (90%)	0.22	18 (5%) 30 32	17, 23, 31, 40	0
1	C	348/377 (92%)	0.01	9 (2%) 59 64	14, 20, 27, 32	0
2	B	188/188 (100%)	0.29	8 (4%) 39 43	10, 19, 26, 33	0
2	D	188/188 (100%)	0.06	7 (3%) 45 50	14, 18, 25, 31	0
All	All	1066/1130 (94%)	0.13	42 (3%) 43 47	10, 21, 28, 40	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	552	THR	5.3
2	D	552	THR	5.2
1	C	379	ASN	4.5
1	C	377	GLY	4.5
1	A	31	PRO	4.4
1	C	378	SER	4.3
1	C	376	GLU	4.0
2	B	565	LYS	4.0
1	A	200	LYS	3.8
1	C	375	HIS	3.5
1	A	118	LYS	3.3
1	C	118	LYS	3.3
2	D	565	LYS	3.2
1	A	119	GLN	3.1
1	A	201	GLY	2.9
2	B	399	ILE	2.9
2	D	551	LYS	2.8
2	D	567	PHE	2.7
1	A	71	ALA	2.7
1	A	202	HIS	2.6
2	D	550	PRO	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	553	LYS	2.5
1	A	120	GLY	2.5
1	A	36	LYS	2.5
1	A	82	ILE	2.5
2	B	415	LEU	2.5
1	A	292	ILE	2.4
1	A	68	ILE	2.4
1	A	372	GLY	2.4
1	A	122	VAL	2.4
1	A	83	GLY	2.3
2	D	554	GLY	2.3
2	B	567	PHE	2.3
1	A	229	LYS	2.3
1	C	292	ILE	2.2
2	B	393	VAL	2.2
1	C	119	GLN	2.2
2	D	481	LEU	2.2
1	C	245	LYS	2.1
1	A	359	ASP	2.1
2	B	416	LEU	2.1
1	A	73	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	AVN	B	1	10/11	0.88	0.10	-0.17	24,28,39,40	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	AVN	D	1	10/11	0.93	0.09	-0.38	20,27,32,33	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.