



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 29, 2016 – 11:22 AM EDT

PDB ID : 5FOI
Title : Crystal structure of mycinamicin VIII C21 methyl hydroxylase MycCI from *Micromonospora griseorubida* bound to mycinamicin VIII
Authors : Demars, M.; Sheng, F.; Podust, L.M.; Sherman, D.H.
Deposited on : 2015-11-22
Resolution : 2.21 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939

i

X-RAY DIFFRACTION

A.

Metric	Percentile Rank	Value
--------	-----------------	-------



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	4405 (2.24-2.20)
Clashscore	102246	5146 (2.24-2.20)
Ramachandran outliers	100387	5065 (2.24-2.20)
Sidechain outliers	100360	5066 (2.24-2.20)
RSRZ outliers	91569	4414 (2.24-2.20)

Mol	Chain	Length	Quality of chain
1	A	408	<div><div></div><div></div><div></div></div> 78%16%5%
1	B	408	<div><div></div><div></div><div></div></div> %77%17%5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	1390	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6369 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MYCINAMICIN VIII C21 METHYL HYDROXYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	389	Total	C	N	O	S	0	0	0
			3004	1903	542	547	12			
1	B	388	Total	C	N	O	S	0	1	0
			3018	1909	549	547	13			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP Q83WF5
A	-18	GLY	-	EXPRESSION TAG	UNP Q83WF5
A	-17	SER	-	EXPRESSION TAG	UNP Q83WF5
A	-16	SER	-	EXPRESSION TAG	UNP Q83WF5
A	-15	HIS	-	EXPRESSION TAG	UNP Q83WF5
A	-14	HIS	-	EXPRESSION TAG	UNP Q83WF5
A	-13	HIS	-	EXPRESSION TAG	UNP Q83WF5
A	-12	HIS	-	EXPRESSION TAG	UNP Q83WF5
A	-11	HIS	-	EXPRESSION TAG	UNP Q83WF5
A	-10	HIS	-	EXPRESSION TAG	UNP Q83WF5
A	-9	SER	-	EXPRESSION TAG	UNP Q83WF5
A	-8	SER	-	EXPRESSION TAG	UNP Q83WF5
A	-7	GLY	-	EXPRESSION TAG	UNP Q83WF5
A	-6	LEU	-	EXPRESSION TAG	UNP Q83WF5
A	-5	VAL	-	EXPRESSION TAG	UNP Q83WF5
A	-4	PRO	-	EXPRESSION TAG	UNP Q83WF5
A	-3	ARG	-	EXPRESSION TAG	UNP Q83WF5
A	-2	GLY	-	EXPRESSION TAG	UNP Q83WF5
A	-1	SER	-	EXPRESSION TAG	UNP Q83WF5
A	0	HIS	-	EXPRESSION TAG	UNP Q83WF5
B	-19	MET	-	EXPRESSION TAG	UNP Q83WF5
B	-18	GLY	-	EXPRESSION TAG	UNP Q83WF5
B	-17	SER	-	EXPRESSION TAG	UNP Q83WF5
B	-16	SER	-	EXPRESSION TAG	UNP Q83WF5
B	-15	HIS	-	EXPRESSION TAG	UNP Q83WF5

Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	EXPRESSION TAG	UNP Q83WF5
B	-13	HIS	-	EXPRESSION TAG	UNP Q83WF5
B	-12	HIS	-	EXPRESSION TAG	UNP Q83WF5
B	-11	HIS	-	EXPRESSION TAG	UNP Q83WF5
B	-10	HIS	-	EXPRESSION TAG	UNP Q83WF5
B	-9	SER	-	EXPRESSION TAG	UNP Q83WF5
B	-8	SER	-	EXPRESSION TAG	UNP Q83WF5
B	-7	GLY	-	EXPRESSION TAG	UNP Q83WF5
B	-6	LEU	-	EXPRESSION TAG	UNP Q83WF5
B	-5	VAL	-	EXPRESSION TAG	UNP Q83WF5
B	-4	PRO	-	EXPRESSION TAG	UNP Q83WF5
B	-3	ARG	-	EXPRESSION TAG	UNP Q83WF5
B	-2	GLY	-	EXPRESSION TAG	UNP Q83WF5
B	-1	SER	-	EXPRESSION TAG	UNP Q83WF5
B	0	HIS	-	EXPRESSION TAG	UNP Q83WF5

- # HEM
-
- The diagram illustrates the chemical structure of Hemoglobin (HEM). It features a central iron atom (Fe) coordinated by four nitrogen atoms (N) in a porphyrin-like ring. The ring is substituted with various side chains, including vinyl groups (CH=CH₂), methyl groups (CH₃), and other functional groups. The structure is labeled with numerous atoms and bonds, indicating a complex molecular structure.


- Molecule 3 is MYCINAMICIN VIII (three-letter code: MY8) (formula: $\text{C}_{29}\text{H}_{49}\text{NO}_6$).



- # GOL
-
- The diagram shows the chemical structure of glycerol (1,2,3-propanetriol). It consists of a three-carbon chain. The first carbon (C1) is bonded to a hydroxyl group (HO) with oxygen O1. The second carbon (C2) is bonded to two hydroxyl groups (OH) with oxygens O2 and O3. The third carbon (C3) is bonded to a hydroxyl group (OH) with oxygen O4. The atoms are labeled with green text: C1, C2, C3 for carbons and O1, O2, O3, O4 for oxygens. The hydroxyl groups are shown in red text.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0

- Molecule 5 is SPERMIDINE (three-letter code: SPD) (formula: C₇H₁₉N₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	N	0	0
			10	7	3		
5	A	1	Total	C	N	0	0
			10	7	3		
5	A	1	Total	C	N	0	0
			10	7	3		


- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	77	Total	O	0	0
			77	77		
6	B	70	Total	O	0	0
			70	70		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: MYCINAMICIN VIII C21 METHYL HYDROXYLASE

Chain A: 



• Molecule 1: MYCINAMICIN VIII C21 METHYL HYDROXYLASE

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.78Å 59.52Å 74.62Å 83.20° 72.15° 62.61°	Depositor
Resolution (Å)	71.00 – 2.21 71.01 – 2.21	Depositor EDS
% Data completeness (in resolution range)	90.8 (71.00-2.21) 90.3 (71.01-2.21)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.182 , 0.252 0.189 , 0.251	Depositor DCC
R_{free} test set	1838 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	26.0	Xtriage
Anisotropy	0.500	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 40.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.010 for -h,-h+k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6369	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SPD, MY8, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	0/3070	0.87	3/4188 (0.1%)
1	B	0.68	0/3083	0.88	4/4201 (0.1%)
All	All	0.69	0/6153	0.88	7/8389 (0.1%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	254	ARG	NE-CZ-NH1	9.03	124.81	120.30
1	A	245	ARG	NE-CZ-NH1	6.34	123.47	120.30
1	A	52	VAL	CB-CA-C	-5.94	100.12	111.40
1	B	275	ASP	CB-CG-OD1	5.74	123.47	118.30
1	A	91	ARG	NE-CZ-NH1	5.48	123.04	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3004	0	2974	42	0
1	B	3018	0	3004	53	0
2	A	43	0	30	2	0
2	B	43	0	30	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	36	0	0	1	0
3	B	36	0	0	0	0
4	A	12	0	16	3	0
5	A	30	0	57	3	0
6	A	77	0	0	5	0
6	B	70	0	0	2	0
All	All	6369	0	6111	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 102 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:ASP:OD2	1:A:132:THR:OG1	2.06	0.73
1:A:340:GLN:HE21	1:A:344:ARG:HH12	1.38	0.70
1:B:138:MET:CE	1:B:142:VAL:HG21	2.24	0.68
1:B:195:ASP:OD2	1:B:196:THR:O	2.11	0.68
1:B:231:THR:HG21	2:B:413:HEM:CHC	2.25	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	387/408 (95%)	369 (95%)	18 (5%)	0	100	100
1	B	387/408 (95%)	372 (96%)	14 (4%)	1 (0%)	46	50
All	All	774/816 (95%)	741 (96%)	32 (4%)	1 (0%)	56	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	27	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/347 (90%)	304 (97%)	9 (3%)	50	61
1	B	316/347 (91%)	306 (97%)	10 (3%)	46	57
All	All	629/694 (91%)	610 (97%)	19 (3%)	48	60

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	371	ARG
1	B	34	ASP
1	B	171	ARG
1	A	365	GLN
1	B	186	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	141	GLN
1	B	168	ASN
1	B	326	HIS
1	A	340	GLN
1	B	335	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	1389	-	5,5,5	0.24	0	5,5,5	0.57	0
4	GOL	A	1390	-	5,5,5	0.51	0	5,5,5	0.66	0
5	SPD	A	1391	-	9,9,9	0.45	0	8,8,8	1.59	2 (25%)
5	SPD	A	1392	-	9,9,9	0.41	0	8,8,8	1.36	2 (25%)
5	SPD	A	1393	-	9,9,9	0.29	0	8,8,8	1.21	1 (12%)
2	HEM	A	413	1	24,50,50	1.08	2 (8%)	16,82,82	1.73	4 (25%)
3	MY8	A	414	-	35,37,37	1.98	4 (11%)	41,51,51	3.62	27 (65%)
2	HEM	B	413	1	24,50,50	0.80	0	16,82,82	1.30	3 (18%)
3	MY8	B	414	-	35,37,37	2.05	4 (11%)	41,51,51	3.35	24 (58%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1389	-	-	0/4/4/4	0/0/0/0
4	GOL	A	1390	-	-	0/4/4/4	0/0/0/0
5	SPD	A	1391	-	-	0/7/7/7	0/0/0/0
5	SPD	A	1392	-	-	0/7/7/7	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SPD	A	1393	-	-	0/7/7/7	0/0/0/0
2	HEM	A	413	1	-	0/6/54/54	0/0/8/8
3	MY8	A	414	-	-	0/46/62/62	0/1/2/2
2	HEM	B	413	1	-	0/6/54/54	0/0/8/8
3	MY8	B	414	-	-	0/46/62/62	0/1/2/2

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	414	MY8	O2-C10	-10.19	1.22	1.43
3	A	414	MY8	O2-C10	-10.11	1.22	1.43
3	B	414	MY8	O1-C1	-3.33	1.40	1.46
3	A	414	MY8	C14-C16	-2.85	1.48	1.53
2	A	413	HEM	C3C-C2C	-2.83	1.36	1.40

The worst 5 of 63 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	414	MY8	C15-C14-C16	-9.26	95.46	111.28
3	A	414	MY8	C5-C4-C6	-7.40	91.49	110.16
3	A	414	MY8	C1-O1-C29	-7.40	106.13	117.46
3	A	414	MY8	C12-C11-C13	-6.93	100.23	110.69
3	B	414	MY8	O1-C1-C2	-6.02	96.33	106.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1390	GOL	3	0
5	A	1391	SPD	3	0
5	A	1392	SPD	3	0
2	A	413	HEM	2	0
3	A	414	MY8	1	0
2	B	413	HEM	8	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	389/408 (95%)	-0.51	1 (0%) 94 94	15, 26, 44, 59	0
1	B	388/408 (95%)	-0.48	3 (0%) 87 87	15, 26, 43, 61	1 (0%)
All	All	777/816 (95%)	-0.49	4 (0%) 91 91	15, 26, 44, 61	1 (0%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	25	ALA	3.5
1	B	313	GLY	2.8
1	B	26	THR	2.8
1	B	25	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	GOL	A	1390	6/6	0.74	0.22	8.34	45,50,52,53	0
2	HEM	B	413	43/43	0.96	0.12	0.59	12,17,19,23	0
3	MY8	A	414	36/36	0.94	0.12	0.33	16,19,24,25	0
2	HEM	A	413	43/43	0.96	0.10	-0.13	12,16,18,19	0
4	GOL	A	1389	6/6	0.96	0.10	-0.40	32,37,40,42	0
3	MY8	B	414	36/36	0.94	0.10	-0.92	18,21,24,24	0
5	SPD	A	1391	10/10	0.91	0.16	-	29,31,34,34	0
5	SPD	A	1392	10/10	0.83	0.15	-	39,42,47,47	0
5	SPD	A	1393	10/10	0.75	0.20	-	48,55,56,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.