



# Full wwPDB X-ray Structure Validation Report ⓘ

May 12, 2016 – 04:08 PM EDT

PDB ID : 5FXU  
Title : Crystal Structure of Puumala virus Gn glycoprotein ectodomain  
Authors : Li, S.; Rissanen, I.; Zeltina, A.; Hepojoki, J.; Raghwani, J.; Harlos, K.; Pybus, O.G.; Huiskonen, J.T.; Bowden, T.A.  
Deposited on : 2016-03-02  
Resolution : 2.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027457  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	5193 (2.30-2.26)
Clashscore	102246	5929 (2.30-2.26)
Ramachandran outliers	100387	5851 (2.30-2.26)
Sidechain outliers	100360	5850 (2.30-2.26)
RSRZ outliers	91569	5204 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	367	<div> <div>8%</div> <div> <div></div> <div>83%</div> <div>7%</div> <div>10%</div> </div> </div>
1	B	367	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>6%</div> <div>10%</div> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5551 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENVELOPE POLYPROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	331	Total	C	N	O	S	0	0	0
			2517	1599	409	488	21			
1	B	332	Total	C	N	O	S	0	4	0
			2551	1623	416	491	21			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	GLU	-	EXPRESSION TAG	UNP Q9WJ31
A	27	THR	-	EXPRESSION TAG	UNP Q9WJ31
A	28	GLY	-	EXPRESSION TAG	UNP Q9WJ31
A	384	GLY	-	EXPRESSION TAG	UNP Q9WJ31
A	385	THR	-	EXPRESSION TAG	UNP Q9WJ31
A	386	LYS	-	EXPRESSION TAG	UNP Q9WJ31
A	387	HIS	-	EXPRESSION TAG	UNP Q9WJ31
A	388	HIS	-	EXPRESSION TAG	UNP Q9WJ31
A	389	HIS	-	EXPRESSION TAG	UNP Q9WJ31
A	390	HIS	-	EXPRESSION TAG	UNP Q9WJ31
A	391	HIS	-	EXPRESSION TAG	UNP Q9WJ31
A	392	HIS	-	EXPRESSION TAG	UNP Q9WJ31
B	26	GLU	-	EXPRESSION TAG	UNP Q9WJ31
B	27	THR	-	EXPRESSION TAG	UNP Q9WJ31
B	28	GLY	-	EXPRESSION TAG	UNP Q9WJ31
B	384	GLY	-	EXPRESSION TAG	UNP Q9WJ31
B	385	THR	-	EXPRESSION TAG	UNP Q9WJ31
B	386	LYS	-	EXPRESSION TAG	UNP Q9WJ31
B	387	HIS	-	EXPRESSION TAG	UNP Q9WJ31
B	388	HIS	-	EXPRESSION TAG	UNP Q9WJ31
B	389	HIS	-	EXPRESSION TAG	UNP Q9WJ31
B	390	HIS	-	EXPRESSION TAG	UNP Q9WJ31
B	391	HIS	-	EXPRESSION TAG	UNP Q9WJ31
B	392	HIS	-	EXPRESSION TAG	UNP Q9WJ31

- Molecule 2 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	4	Total	C	N	O	0	0
			50	28	2	20		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	B	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 4 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	3	Total	C	N	O	0	0
			39	22	2	15		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	138	Total	O	0	0
			138	138		
5	B	200	Total	O	0	0
			200	200		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ENVELOPE POLYPROTEIN



#### • Molecule 1: ENVELOPE POLYPROTEIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.57Å 66.79Å 77.35Å 107.30° 93.55° 100.94°	Depositor
Resolution (Å)	73.26 – 2.28 62.23 – 2.28	Depositor EDS
% Data completeness (in resolution range)	97.9 (73.26-2.28) 88.1 (62.23-2.28)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.10 (at 2.27Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.189 , 0.219 0.192 , 0.222	Depositor DCC
$R_{free}$ test set	2171 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	39.7	Xtriage
Anisotropy	0.117	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 39.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5551	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	0/2567	0.80	3/3488 (0.1%)
1	B	0.62	0/2611	0.78	4/3548 (0.1%)
All	All	0.61	0/5178	0.79	7/7036 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	282	ARG	NE-CZ-NH1	9.96	125.28	120.30
1	B	217	ARG	NE-CZ-NH2	-6.43	117.08	120.30
1	B	131	ARG	NE-CZ-NH1	5.81	123.21	120.30
1	A	282	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	A	131	ARG	NE-CZ-NH1	5.63	123.12	120.30
1	B	258	LEU	CB-CG-CD2	5.13	119.71	111.00
1	B	217	ARG	NE-CZ-NH1	5.06	122.83	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2517	0	2512	12	1
1	B	2551	0	2556	14	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	50	0	43	0	0
3	A	28	0	25	0	0
3	B	28	0	25	0	0
4	B	39	0	34	0	0
5	A	138	0	0	0	0
5	B	200	0	0	2	0
All	All	5551	0	5195	26	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:224:LYS:NZ	1:B:343:ASP:OD1	2.27	0.67
1:A:241:VAL:HG12	1:A:241:VAL:O	2.02	0.58
1:A:269:PRO:HB3	1:A:280:LEU:HD11	1.86	0.57
1:A:36:HIS:NE2	1:A:183:CYS:O	2.37	0.57
1:B:269:PRO:HB3	1:B:280:LEU:HD11	1.87	0.56
1:B:195[B]:ASN:HD22	1:B:379:GLU:HG2	1.72	0.54
1:A:193:CYS:O	1:A:380:ILE:O	2.25	0.54
1:A:282:ARG:HH11	1:A:282:ARG:HG2	1.72	0.53
1:A:338:LEU:HG	1:A:347:ILE:HD11	1.92	0.52
1:B:338:LEU:HG	1:B:347:ILE:HD11	1.93	0.51
1:A:241:VAL:CG1	1:A:241:VAL:O	2.59	0.50
1:B:80:GLN:HG2	1:B:113:LEU:O	2.13	0.49
1:A:80:GLN:HG2	1:A:113:LEU:O	2.14	0.47
1:A:189:VAL:HG11	1:A:214:VAL:HG12	1.97	0.46
1:A:36:HIS:CD2	1:A:183:CYS:O	2.70	0.45
1:B:70:PHE:CE1	1:B:72:LEU:HD13	2.52	0.45
1:B:232:LYS:HE3	5:B:2131:HOH:O	2.17	0.44
1:A:42:GLN:HB2	1:A:264:GLU:OE2	2.18	0.43
1:B:70:PHE:HE1	1:B:72:LEU:HD13	1.83	0.43
1:B:268:ILE:HA	1:B:269:PRO:HA	1.86	0.43
1:B:51:LEU:HD11	1:B:174:ILE:HD11	2.01	0.42
1:B:321:ILE:HD12	1:B:347:ILE:HD13	2.01	0.42
1:B:118:LEU:HD12	1:B:368[A]:LEU:HD21	2.02	0.42
1:A:282:ARG:HH11	1:A:282:ARG:CG	2.32	0.41
1:B:131:ARG:HH11	1:B:131:ARG:HG3	1.86	0.40
1:B:117:CYS:HB2	5:B:2067:HOH:O	2.21	0.40



All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:272:ASP:OD1	1:B:203:SER:OG[1_545]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	323/367 (88%)	316 (98%)	7 (2%)	0	100	100
1	B	328/367 (89%)	319 (97%)	9 (3%)	0	100	100
All	All	651/734 (89%)	635 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/323 (90%)	281 (97%)	10 (3%)	44	58
1	B	296/323 (92%)	287 (97%)	9 (3%)	48	63
All	All	587/646 (91%)	568 (97%)	19 (3%)	45	61

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	LYS
1	A	70	PHE
1	A	80	GLN
1	A	173	ARG
1	A	209	TYR
1	A	224	LYS
1	A	243	LYS
1	A	258	LEU
1	A	282	ARG
1	A	291	ASP
1	B	63	LYS
1	B	70	PHE
1	B	72	LEU
1	B	80	GLN
1	B	209	TYR
1	B	217	ARG
1	B	376	LEU
1	B	379	GLU
1	B	382	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

11 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	1382	1,2	14,14,15	0.40	0	15,19,21	1.26	2 (13%)
2	NAG	A	1383	2	14,14,15	0.55	0	15,19,21	0.84	0
2	BMA	A	1384	2	11,11,12	0.68	0	15,15,17	1.50	5 (33%)
2	MAN	A	1385	2	11,11,12	0.82	0	15,15,17	2.29	4 (26%)
3	NAG	A	1386	1,3	14,14,15	0.37	0	15,19,21	1.40	3 (20%)
3	NAG	A	1387	3	14,14,15	0.43	0	15,19,21	1.05	1 (6%)
3	NAG	B	1383	1,3	14,14,15	0.39	0	15,19,21	1.25	2 (13%)
3	NAG	B	1384	3	14,14,15	0.33	0	15,19,21	1.07	1 (6%)
4	NAG	B	1385	1,4	14,14,15	0.50	0	15,19,21	0.87	0
4	NAG	B	1386	4	14,14,15	0.50	0	15,19,21	1.02	0
4	BMA	B	1387	4	11,11,12	0.66	0	15,15,17	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1382	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	1383	2	-	0/6/23/26	0/1/1/1
2	BMA	A	1384	2	-	0/2/19/22	0/1/1/1
2	MAN	A	1385	2	-	0/2/19/22	0/1/1/1
3	NAG	A	1386	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1387	3	-	0/6/23/26	0/1/1/1
3	NAG	B	1383	1,3	-	0/6/23/26	0/1/1/1
3	NAG	B	1384	3	-	0/6/23/26	0/1/1/1
4	NAG	B	1385	1,4	-	0/6/23/26	0/1/1/1
4	NAG	B	1386	4	-	0/6/23/26	0/1/1/1
4	BMA	B	1387	4	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1382	NAG	O4-C4-C3	-3.24	103.05	110.36
2	A	1384	BMA	O5-C5-C4	-2.26	106.39	110.13
2	A	1384	BMA	C2-C3-C4	2.03	114.59	111.05
2	A	1385	MAN	O4-C4-C5	2.14	114.86	109.23
3	A	1386	NAG	C2-N2-C7	2.14	125.89	123.11
2	A	1382	NAG	C4-C3-C2	2.17	114.71	111.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1384	BMA	O5-C1-C2	2.29	114.56	110.89
2	A	1384	BMA	C1-O5-C5	2.33	115.56	112.14
3	B	1383	NAG	C8-C7-N2	2.33	120.57	116.10
3	B	1383	NAG	C2-N2-C7	2.61	126.50	123.11
2	A	1385	MAN	O5-C5-C4	2.62	114.47	110.13
3	A	1386	NAG	C8-C7-N2	2.64	121.16	116.10
2	A	1385	MAN	O3-C3-C4	2.74	116.54	110.36
2	A	1384	BMA	C1-C2-C3	2.74	112.88	109.55
3	B	1384	NAG	C1-O5-C5	3.28	116.96	112.14
3	A	1386	NAG	C1-O5-C5	3.28	116.97	112.14
3	A	1387	NAG	C1-O5-C5	3.38	117.11	112.14
2	A	1385	MAN	C1-O5-C5	7.08	122.55	112.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	331/367 (90%)	0.43	29 (8%) 12 17	25, 46, 102, 138	0
1	B	332/367 (90%)	0.08	8 (2%) 62 70	23, 41, 76, 117	0
All	All	663/734 (90%)	0.25	37 (5%) 28 35	23, 43, 92, 138	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	40	LEU	7.7
1	A	182	TYR	7.2
1	A	200	MET	6.7
1	A	38	ILE	6.6
1	A	27	THR	6.0
1	A	201	ALA	5.6
1	A	203	SER	5.5
1	B	200	MET	5.5
1	A	199	THR	5.2
1	A	42	GLN	5.1
1	A	41	GLY	4.8
1	A	202	LEU	4.1
1	A	75	SER	4.1
1	B	201	ALA	3.8
1	A	185	SER	3.6
1	A	102	THR	3.3
1	B	203	SER	3.0
1	A	72	LEU	3.0
1	A	36	HIS	3.0
1	A	198	HIS	2.9
1	A	39	GLY	2.9
1	A	77	SER	2.8
1	A	381	GLU	2.7
1	A	209	TYR	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	91	SER	2.6
1	A	244	THR	2.5
1	B	209	TYR	2.5
1	A	74	THR	2.4
1	A	184	VAL	2.4
1	A	173	ARG	2.3
1	B	40	LEU	2.3
1	B	91	SER	2.3
1	A	191	GLY	2.3
1	A	183	CYS	2.3
1	A	181	THR	2.3
1	B	199	THR	2.2
1	B	39	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	B	1385	14/15	0.88	0.21	0.91	42,50,57,64	0
2	NAG	A	1382	14/15	0.91	0.12	-0.73	40,47,55,67	0
2	MAN	A	1385	11/12	0.73	0.27	-	90,103,108,110	0
4	NAG	B	1386	14/15	0.84	0.26	-	66,74,75,76	0
4	BMA	B	1387	11/12	0.76	0.19	-	62,75,78,79	0
2	NAG	A	1383	14/15	0.84	0.16	-	69,78,80,84	0
3	NAG	B	1383	14/15	0.90	0.11	-	50,57,63,76	0
2	BMA	A	1384	11/12	0.84	0.18	-	74,81,85,92	0
3	NAG	A	1387	14/15	0.76	0.15	-	76,87,89,90	0
3	NAG	B	1384	14/15	0.76	0.21	-	81,90,99,102	0
3	NAG	A	1386	14/15	0.89	0.14	-	56,62,68,81	0

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.