



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:31 PM GMT

PDB ID : 1FZP
Title : CRYSTAL STRUCTURES OF SARA: A PLEIOTROPIC REGULATOR OF
VIRULENCE GENES IN S. AUREUS
Authors : Schumacher, M.A.; Hurlburt, B.; Brennan, R.G.
Deposited on : 2000-10-03
Resolution : 2.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| Clashscore | 102246 | 2552 (3.00-2.92) |
| Ramachandran outliers | 100387 | 2468 (3.00-2.92) |
| Sidechain outliers | 100360 | 2471 (3.00-2.92) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--------------------|
| 1 | K | 7 | 100% |
| 1 | W | 7 | 100% |
| 2 | B | 123 | 7% 33% 34% 11% 15% |
| 2 | D | 123 | 9% 32% 31% 10% 19% |

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1997 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(P*AP*TP*AP*TP*AP*TP*A)-3'.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|----|----|----|---|---------|---------|-------|
| 1 | W | 7 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 144 | 70 | 26 | 41 | 7 | | | |
| 1 | K | 7 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 144 | 70 | 26 | 41 | 7 | | | |

- Molecule 2 is a protein called STAPHYLOCOCCAL ACCESSORY REGULATOR A.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 2 | D | 100 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 832 | 537 | 135 | 158 | 2 | | | |
| 2 | B | 105 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 875 | 563 | 145 | 165 | 2 | | | |

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 3 | B | 1 | Total | Ca | 0 | 0 |
| | | | 1 | 1 | | |
| 3 | D | 1 | Total | Ca | 0 | 0 |
| | | | 1 | 1 | | |

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: 5'-D(P*AP*TP*AP*TP*AP*TP*A)-3'

Chain W:  100%

A3
T4
A5
T6
A7
T8
A9

- Molecule 1: 5'-D(P*AP*TP*AP*TP*AP*TP*A)-3'

Chain K:  100%

A3
T4
A5
T6
A7
T8
A9

- Molecule 2: STAPHYLOCOCCAL ACCESSORY REGULATOR A

Chain D:  9% 32% 31% 10% 19%

A2 I3 T4 K5 I6 I7 D8 C9 F10 E11 L12 L13 L14 S14 M15 V16 V17 T17 Y18 A19 D20 D21 L22 L23 K23 S24 K28 K27 K26 L25 E86 H87 D88 E89 R90 F91 T91 Y92 L93 L94 L95 V96 E96 N97 F97 A98 Q99 Q100 R101 L40 T41 K103 Y42 I43 S44 E45 M46 R47 E48 K49 R110 V111 ASN Y51 Y52 L53 K54 D55 I56 ILE ASN HIS ASN

Tyr Lys Glu Glu Glu V67 V68 K69 A70 V71 W72 K72 L73 L74 L75 S75 Q76 E77 D78 Y79 F80 K83 R84 N85 E86 H87 D88 E89 R90 T91 Y92 L93 L94 L95 V96 E96 N97 F97 A98 Q99 Q100 R101 L40 T41 K103 Y42 I43 S44 E45 M46 R47 E48 K49 R110 V111 ASN Y51 Y52 L53 K54 D55 I56 ILE ASN ASN ASN ILE

GLU
LEU

- Molecule 2: STAPHYLOCOCCAL ACCESSORY REGULATOR A

Chain B:  7% 33% 34% 11% 15%

A2 I3 T4 K5 I6 I7 D8 C9 F10 E11 L12 L13 L14 S14 M15 V16 V17 T17 Y18 A19 D20 D21 L22 L23 K23 S24 K28 K27 K26 L25 E86 H87 D88 E89 R90 T91 Y92 L93 L94 L95 V96 E96 N97 F97 A98 Q99 Q100 R101 L40 T41 K103 Y42 I43 S44 E45 M46 R47 E48 K49 R110 V111 Y51 Y52 L53 K54 D55 I56 ILE ASN HIS ASN

Tyr Lys Glu Glu Glu V67 V68 K69 L73 L74 L75 S75 Q76 E77 D78 Y79 F80 D81 K82 K83 R84 N85 E86 H87 D88 E89 R90 T91 Y92 L93 L94 L95 V96 E96 N97 F97 A98 Q99 Q100 R101 L40 T41 K103 Y42 I43 S44 E45 M46 R47 E48 K49 R110 V111 N112 K113 R114 T115 T116 ALA ASN ASN ASN ASN ASN ILE GLU LEU GLU

LEU

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

| Property | Value | Source |
|--|--|-----------|
| Space group | P 1 21 1 | Depositor |
| Cell constants a, b, c, α , β , γ | 54.50 Å 65.20 Å 57.80 Å 90.00° 118.00° 90.00° | Depositor |
| Resolution (Å) | 10.00 – 2.95 | Depositor |
| % Data completeness (in resolution range) | 89.0 (10.00-2.95) | Depositor |
| R_{merge} | 0.08 | Depositor |
| R_{sym} | (Not available) | Depositor |
| Refinement program | TNT | Depositor |
| R, R_{free} | 0.220 , 0.308 | Depositor |
| Estimated twinning fraction | No twinning to report. | Xtriage |
| Total number of atoms | 1997 | wwPDB-VP |
| Average B, all atoms (Å ²) | 43.0 | wwPDB-VP |

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|----------------|-------------|----------------|
| | | RMSZ | $\# Z > 5$ | RMSZ | $\# Z > 5$ |
| 1 | K | 1.85 | 4/161 (2.5%) | 2.87 | 17/246 (6.9%) |
| 1 | W | 1.93 | 5/161 (3.1%) | 2.82 | 18/246 (7.3%) |
| 2 | B | 0.87 | 1/884 (0.1%) | 1.51 | 16/1181 (1.4%) |
| 2 | D | 0.99 | 1/841 (0.1%) | 1.51 | 19/1124 (1.7%) |
| All | All | 1.14 | 11/2047 (0.5%) | 1.82 | 70/2797 (2.5%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 2 | B | 1 | 0 |
| 2 | D | 1 | 0 |
| All | All | 2 | 0 |

The worst 5 of 11 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|---------|------|-------------|----------|
| 2 | D | 9 | CYS | CB-SG | 8.80 | 1.97 | 1.82 |
| 1 | W | 8 | DT | C5'-C4' | 7.80 | 1.59 | 1.51 |
| 1 | W | 4 | DT | C5-C7 | 7.73 | 1.54 | 1.50 |
| 1 | K | 4 | DT | C5-C7 | 7.56 | 1.54 | 1.50 |
| 1 | K | 8 | DT | C5-C7 | 6.30 | 1.53 | 1.50 |

The worst 5 of 70 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-------------|--------|-------------|----------|
| 1 | W | 8 | DT | O4'-C4'-C3' | -18.45 | 94.93 | 106.00 |
| 1 | K | 3 | DA | O5'-P-OP2 | -18.20 | 88.86 | 110.70 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-------------|--------|-------------|----------|
| 1 | K | 3 | DA | OP1-P-OP2 | 16.08 | 143.72 | 119.60 |
| 1 | K | 8 | DT | O4'-C4'-C3' | -11.99 | 98.81 | 106.00 |
| 1 | W | 7 | DA | O4'-C4'-C3' | -11.91 | 98.86 | 106.00 |

All (2) chirality outliers are listed below:

| Mol | Chain | Res | Type | Atom |
|-----|-------|-----|------|------|
| 2 | D | 43 | ILE | CA |
| 2 | B | 13 | LEU | CA |

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | K | 144 | 0 | 81 | 43 | 0 |
| 1 | W | 144 | 0 | 81 | 46 | 0 |
| 2 | B | 875 | 0 | 916 | 210 | 0 |
| 2 | D | 832 | 0 | 866 | 235 | 0 |
| 3 | B | 1 | 0 | 0 | 0 | 0 |
| 3 | D | 1 | 0 | 0 | 0 | 0 |
| All | All | 1997 | 0 | 1944 | 489 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 125.

The worst 5 of 489 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|---------------|----------------|--------------------------|-------------------|
| 2:D:8:ASP:HB2 | 2:D:13:LEU:CD2 | 1.24 | 1.59 |
| 1:W:5:DA:H61 | 1:K:5:DA:N6 | 1.17 | 1.41 |
| 1:W:5:DA:N6 | 1:K:5:DA:H61 | 1.17 | 1.34 |
| 1:W:3:DA:N1 | 1:K:7:DA:N6 | 1.83 | 1.27 |
| 2:D:8:ASP:CB | 2:D:13:LEU:CD2 | 2.20 | 1.18 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|---------------|-----------|----------|----------|-------------|---|
| 2 | B | 101/123 (82%) | 73 (72%) | 19 (19%) | 9 (9%) | 1 | 4 |
| 2 | D | 96/123 (78%) | 72 (75%) | 13 (14%) | 11 (12%) | 0 | 2 |
| All | All | 197/246 (80%) | 145 (74%) | 32 (16%) | 20 (10%) | 1 | 2 |

5 of 20 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | D | 3 | ILE |
| 2 | D | 30 | PHE |
| 2 | D | 31 | SER |
| 2 | D | 100 | GLN |
| 2 | B | 3 | ILE |

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|---------------|-----------|----------|-------------|---|
| 2 | B | 100/117 (86%) | 50 (50%) | 50 (50%) | 0 | 0 |
| 2 | D | 95/117 (81%) | 52 (55%) | 43 (45%) | 0 | 0 |
| All | All | 195/234 (83%) | 102 (52%) | 93 (48%) | 0 | 0 |

5 of 93 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | D | 111 | VAL |
| 2 | B | 28 | LYS |
| 2 | B | 107 | LEU |
| 2 | B | 3 | ILE |
| 2 | B | 11 | GLU |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | B | 7 | ASN |
| 2 | B | 99 | GLN |
| 2 | B | 76 | GLN |
| 2 | D | 99 | GLN |
| 2 | B | 85 | ASN |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.